

10-21-00

A

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b> <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	BB1043 US NA DIV
	First Named Inventor or Application Identifier	
	Jonathan Edward Lightner et al.	
	Express Mail Label No.	EL073740691US
	Express Mailing Date	October 26, 2000

10/26/00 10:58:39 PTO

10/26/00

<b>APPLICATION ELEMENTS</b> See MPEP chapter 600 concerning utility patent application contents.		<b>ADDRESS TO:</b> Assistant Commissioner for Patents Box Patent Application Washington, DC 20231	
1. <input checked="" type="checkbox"/> Fee (Authority to charge deposit account below.) <i>(Submit an original, and a duplicate for fee processing)</i> 2. <input checked="" type="checkbox"/> Specification [Total Pages <b>161</b> ] <i>(preferred arrangement set forth below)</i> - Descriptive title of the invention - Cross References to Related Applications <i>(if needed)</i> - Statement Regarding Fed sponsored R & D <i>(if needed)</i> - Reference to Microfiche Appendix <i>(if filed)</i> - Background of the Invention - Brief Summary of the Invention - Brief Description of the Drawings <i>(if filed)</i> - Detailed Description - Claim(s) - Abstract of the Disclosure 3. <input type="checkbox"/> Drawing(s) (35 USC 113) [Total Sheets <b>0</b> ] 4. <input checked="" type="checkbox"/> Oath or Declaration [Total Pages <b>1</b> ] a. <input type="checkbox"/> Newly executed (original or copy) b. <input checked="" type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 14 completed)</i> i. <input type="checkbox"/> <b>DELETION OF INVENTORS</b> Signed Statement below at 15 deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b). 5. <input checked="" type="checkbox"/> Incorporation by Reference <i>(useable if Box 4b is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.	6. <input type="checkbox"/> Microfiche Computer Program (Appendix) 7. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission <i>(if applicable, all necessary)</i> a. <input checked="" type="checkbox"/> Computer Readable Copy b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy) c. <input checked="" type="checkbox"/> Statement verifying identity of above copies		
<b>ACCOMPANYING APPLICATION PARTS</b>			
8. <input type="checkbox"/> Power of Attorney 9. <input type="checkbox"/> Information Disclosure Statement (IDS)/Cover Letter plus PTO-1449 <input type="checkbox"/> Copies of IDS Citations 10. <input checked="" type="checkbox"/> Preliminary Amendment 11. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i> 12. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i> 13. <input type="checkbox"/> Other:			
14. If a <b>CONTINUING APPLICATION</b> , check appropriate box and supply the requisite information: <input type="checkbox"/> Continuation <input checked="" type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior Application No.: <u>09/133,962</u>			
15. <input type="checkbox"/> <b>DELETION OF INVENTOR(S) STATEMENT:</b> This application is being filed by less than all the inventors named in the prior application. In accordance with 37 CFR 1.63(d)(2) and 1.33(b), the Assistant Commissioner is requested to delete the name(s) of the following person or persons who are not inventors of the invention being claimed in this application:			
16. <input checked="" type="checkbox"/> Amend the specification by inserting before the first line the sentence: -- This is a division of Application No. 09/133,962 filed August 14, 1998, now pending. --			
17. <input type="checkbox"/> Cancel in this application original claims ____ of the prior application before calculating the filing. (At least one original independent claim must be retained for filing purposes.)			
18. <input type="checkbox"/> Priority of foreign Application No. _____ filed on _____ in _____ _____ is claimed under 35 U.S.C. 119. (country)			

CLAIMS	(1) FOR	(2) NUMBER FILED	(3) NUMBER EXTRA	(4) RATE	(5) CALCULATIONS
	<b>TOTAL CLAIMS</b> (37 CFR 1.16(e))	67 - 20 =	47	x \$ 18 =	\$ 846.00
	<b>INDEPENDENT CLAIMS</b> (37 CFR 1.16(b))	18 - 3 =	15	x \$ 80 =	\$ 1200.00
	<b>MULTIPLE DEPENDENT CLAIM(S) (if applicable)</b>			+ \$ 270 =	270.00
				<b>BASIC FEE</b> (37 CFR 1.16(a))	+ \$ 710.00
				<b>TOTAL =</b>	<b>\$ 3026.00</b>

19. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 04-1928:

a. ☒ Fees required under 37 CFR 1.16.

b. ☒ Fees required under 37 CFR 1.17.

20. ☐ Other:

#### 21. CORRESPONDENCE ADDRESS

NAME	Lynne M. Christenbury					
ADDRESS	E. I. du Pont de Nemours and Company					
	Legal - Patents					
	1007 Market Street					
CITY	Wilmington	STATE	Delaware	ZIP CODE	19898	
COUNTRY	U.S.A.	TELEPHONE	302-992-5481	FAX	302-892-1026	

#### 22. SIGNATURE OF ATTORNEY OR AGENT REQUIRED

NAME	Lynne M. Christenbury	REG. NO.: 30,971
SIGNATURE	<i>Lynne M. Christenbury</i>	
DATE	26 October 2000	

EXPRESS MAIL LABEL NO. EL073740691US

PATENT  
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN THE APPLICATION OF:

JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY

CASE NO.: BB-1043 US NA DIV

APPLN. NO.: UNKNOWN

GROUP ART UNIT: UNKNOWN

FILED: CONCURRENTLY HEREWITH

EXAMINER: UNKNOWN

FOR: GENES FOR MICROSOMAL DELTA-12  
FATTY ACID DESATURASES AND  
RELATED ENZYMES FROM PLANTS

Date: OCTOBER 26, 2000

Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

Preliminary Amendment

This is submitted to facilitate prosecution of the above-identified application.

In the Claims

Kindly cancel claims 2-8, 11, 12, 14 and 21-23.

Kindly amend the following claims:

9. (amended) Oil obtained from the seeds of [the plants of Claim 8] a  
soybean plant having a chimeric gene comprising a nucleic acid fragment encoding a  
fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide  
encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15.

10. (amended) A method of producing seed oil containing altered levels of  
[unsaturated] hydroxylated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a  
chimeric gene [of Claim 5] comprising a nucleic acid fragment encoding a delta-12  
hydroxylase;
- (b) growing fertile plants from the transformed plant cells of step  
(a);

(c) screening progeny seeds from the fertile plants of step (b) for the desired levels of [unsaturated] hydroxylated fatty acids; and

(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of [unsaturated] hydroxylated fatty acids.

15. (amended) A method for altering fatty acids composition in seeds comprising:

(a) making a cross between a mutant line with altered fatty acid composition with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acids.

09697379.102600

16. (amended) A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant a delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and  
(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

18. (amended) A method for reducing saturated fatty acids in rapeseed seeds comprising:

09697379.102600

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a delta-12 desaturase or a delta-12 hydroxylase wherein said nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42 °C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

(b) growing fertile plants from seeds obtained from the cross; and  
(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

19. (amended) A method for reducing polyunsaturated fatty acids in soybean oil comprising:

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a

transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant a delta-12 desaturase or a delta-12 hydroxylase wherein said isolated nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42 °C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

20. (amended) A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant [containing] comprising [the] a chimeric gene [of Claim 7] capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment selected from the group consisting of:

09697379.102600

(i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding a plant enzyme encoding a delta-12 desaturase or a delta-12 hydroxylase wherein said nucleic acid fragment hybridizes to one of the nucleotide sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, and 15 under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50 °C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42 °C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each;

(ii) the isolated nucleic acid fragment of (i) wherein the encoded polypeptide has an amino acid identity of 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15; or

(iii) the isolated nucleic acid fragment of (i) wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15;

- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

27. (amended) Oil obtained from the plants of Claims [21-26] 24-26.

Please add the following new claims:

28. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

- (a) transforming a plant oil-producing species with a chimeric gene comprising the nucleic acid fragment encoding all or a portion of a plant delta-12 desaturase;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids;



(d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

29. The product of the method of Claim 28.

30. Canola oil obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed canola plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

31. The soybean oil of Claim 9 obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed soybean plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

32. The canola oil of Claim 30 obtained from the seeds of a plant having a chimeric gene capable of causing altered levels of fatty acid in a transformed canola plant cell, said chimeric gene comprising an isolated nucleic acid fragment comprising a sequence encoding a fatty acid desaturase wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15, operably linked to a suitable regulatory sequence.

33. An isolated nucleic acid fragment comprising a nucleic acid sequence selected from the group consisting of:

(a) a nucleic acid sequence encoding a fatty acid desaturase or a fatty acid desaturase-related plant enzyme with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15, or

(b) a nucleic acid sequence or a part thereof which is useful in antisense inhibition or sense suppression of endogenous desaturase activity in a transformed plant wherein the nucleic acid has an identity of 80% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15 or a part thereof.

34. The isolated nucleic acid fragment of Claim 33 wherein the amino acid identity is 60% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.

35. The isolated nucleic acid fragment of Claim 33 wherein the nucleic acid identity is 90% or greater to any one of the sequences set forth in SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.

36. The isolated nucleic acid fragment of any of Claim 33, 34 or 35 wherein said fragment is isolated from an oil-producing plant species.

37. A chimeric gene comprising the nucleic acid fragment of Claim 33, 34 or 35 operably linked to suitable regulatory sequences.

38. A plant comprising in its genome the chimeric gene of Claim 37.

39. Seeds obtained from the plant of Claim 38.

40. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 37;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

41. The isolated nucleic acid fragment of Claim 33, 34, or 35 comprising a nucleic acid sequence encoding a plant microsomal delta-12 fatty acid desaturase.

42. A nucleic acid fragment isolated from an oil-producing plant species wherein said fragment is selected from the group consisting of:

- (i) an isolated nucleic acid fragment comprising a nucleic acid sequence encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered

09697379.102600

from the methyl end of an 18 carbon long fatty acyl chain, wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain and further wherein the amino acid sequence comprising said enzyme contains at least one of the following amino acid sequences selected from the group consisting of: AIPPHCF, AWXXYW, HECGH, LLVPY, WKYSHR, and SHRRHH;

(ii) an isolated nucleic acid fragment encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered from the methyl end of an 18 carbon long fatty acyl chain wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain wherein said isolated nucleic acid fragment encodes a protein comprising any one of the amino acid sequences set forth in SEQ ID NOS:2, 4, 6, 8, 10 or 12;

(iii) an isolated nucleic acid fragment encoding an enzyme which catalyzes a reaction at carbon positions 6 and 7 numbered from the methyl end of an 18 carbon long fatty acyl chain, wherein positions 6 and 7 correspond to carbon positions 12 and 13 numbered from the carbonyl carbon of an 18 carbon long fatty acyl chain wherein said isolated nucleic acid fragment hybridizes to the isolated nucleic acid fragment of (ii) under one of the following sets of conditions:

(a) hybridization in 50 mM Tris, pH 7.6, 6X SSC, 5X Denhardt's, 0.5% sodium dodecyl sulfate (SDS), 100 µg denatured calf thymus DNA and 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each;

(b) hybridization in 6X SSPE, 5X Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 5% dextran sulfate, 100 µg denatured calf thymus DNA at 50°C and wash twice with 2X SSC, 0.5% SDS at room temperature for 15 min each, then wash twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each and then wash twice with 0.2X SSC, 0.5% SDS at 50 °C for 15 min each; or

(c) hybridization in 50% formamide, 5X SSPE, 1% sodium dodecyl sulfate (SDS), 1% Denhardt's Reagent, 100 µg denatured salmon sperm DNA at 42 °C and wash twice with 2X SSPE, 0.2% SDS at 42° C for 15 min each, then wash twice with 0.2X SSPE, 0.2% SDS at 55 °C for 30 min each.

43. A chimeric gene comprising the nucleic acid fragment of Claim 42 operably linked to suitable regulatory sequences.

44. A plant comprising in its genome the chimeric gene of Claim 43.

00697379.102600

45. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 43;
- (b) growing fertile plants from the transformed plant cells of step (a);
- (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of unsaturated fatty acids.

46. A method for altering fatty acids composition in seeds comprising:

- (a) making a cross between a mutant line with altered fatty acid composition with a plant containing the chimeric gene of Claim 37;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acid levels.

47. A method for altering fatty acids composition in seeds comprising:

- (a) making a cross between a mutant line with altered fatty acid composition with a plant containing the chimeric gene of Claim 43;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acid levels.

48. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

- (a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 37;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

49. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

00697379.102600

(a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

50. The method of Claim 46 or 47 wherein the cross in (a) is between a progeny plant derived from a seed comprising a Brassica variety having an oleic acid content of about 69% to 77%, based upon total extractable oil and belonging to a line in which the said oleic acid content has been stabilized for both the generation to which the seed belongs and its parent generation.

51. A method for reducing saturated fatty acids in rapeseed seeds comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

52. A method for reducing saturated fatty acids in rapeseed seeds comprising:

(a) making a cross between a rapeseed variety with increased oleic acid content with a plant containing the chimeric gene of Claim 49;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

53. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

54. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

09697379, 102600

(a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

55. A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant containing the chimeric gene of Claim 37;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

56. A method for reducing saturated fatty acids in soybean seeds comprising:

(a) making a cross between a soybean variety with increased oleic acid content with a plant containing the chimeric gene of Claim 43;

(b) growing fertile plants from seeds obtained from the cross; and

(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

57. Oil obtained from the seeds of a soybean plant having an oleic acid content of at least 75%.

58. Oil obtained from the seeds of a soybean plant having an oleic acid content of at least 70% and a linoleic acid content less than 14%.

59. The oil of Claim 58 in which the linoleic acid content is less than 10%.--

In the Specification

Kindly amend the specification by adding after the title at line 3:

--This is a divisional of application number 09/133,962, pending, filed August 14, 1998, which is a continuation of application number 08/262,401, filed June 20, 1994, now abandoned, which was a national filing continuation-in-part of PCT/US93/09987, filed October 15, 1993, which was a continuation-in-part of 07/977,339, filed November 17, 1992, now abandoned.--

009201\*9262690

Kindly amend the specification as follows:

Page 13, line 5: please delete "1394" and substitute therefor --1426--.

Page 13, line 7: please delete "99 to 101" and substitute therefor --130 to 132--.

Page 13, lines 7-8: please delete "1248 to 1250" and substitute therefor --1282 to 1284--.

Page 13, line 10: please delete "99 to 1250" and substitute therefor --130 to 1284--.

Page 13, lines 10-11: please delete "1 to 98 and 1251 to 1394" and substitute therefor --1 to 129 and 1285 to 1426--.

Page 13, line 13: please delete "383" and substitute therefor --384--.

Page 13, lines 14-15: please delete "99 to 1250" and substitute therefor --130 to 1284--.

Page 36, line 8: please delete "1394" and substitute therefor --1426--.

Page 36, lines 10: please delete "99 to 101" and substitute therefor --130 to 132-- and delete "1248 to 1250" and substitute therefor --1282 to 1284--.

Page 36, line 13: please delete "99 to 1250" and substitute therefor --130 to 1284-- and delete "1 to 98 and 1251 to 1394" and substitute therefor --1 to 129 and 1285 to 1426--.

Page 36, lines 21: please delete "both"; after "nucleotide" insert --level--; and after "and" insert --90% at the--.

Page 36, line 22: please delete "levels" and insert --level--.

#### Remarks

This case is a divisional application under 37 CFR §1.53(b).

Claim 9 has been amended to recite oil obtained from the seeds of a soybean plant having a chimeric gene comprising a nucleic acid fragment encoding a fatty acid desaturase with an amino acid identity of 50% or greater to the polypeptide encoded by any one of the sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11 or 15. Support for this clarification can be found in the specification on pages 44 - 48 and in the claims as originally filed. Thus, no new matter has been added.

Claim 10 has been clarified by replacing the term "unsaturated" with --hydroxylated and it has been rewritten to be in independent form. Support for the term "hydroxylated" can be found in the specification on page 9 at lines 8-18 of the specification. Thus, no new matter has been added.

009201" 62E26960

Claims 15, 16, 18, 19 and 20 have been amended by reciting alternate hybridization conditions. Support for the three different sets of alternative hybridization and wash conditions can be found in the specification on page 35 at lines 27-32, on page 70, on page 73 at lines 5-11 and on pages 74 and 75. Thus, no new matter has been added.

New claims 28 - 59 have been added.

Claim 28 is similar to original claim 10 except that it is written in independent form.

Support for claim 30 can be found in the specification on pages 44-48 and in Example 3.

Support for claims 31, 32, 34 can be found in the specification on pages 44-48 and, in particular, on page 46.

Support for claim 33 can be found in claim 1 as originally filed and in the specification on pages 54-55.

Support for claim 35 can be found in claim 3 as originally filed.

Support for claims 37-39 can be found throughout the specification and claims as originally filed.

Support for claim 40 can be found in claim 10 as originally filed.

Support for claim 41 can be found on page 6 of the specification.

Support for claim 42 can be found in Table 7 on pages 47 and 48 of the specification and on page 35 at lines 27-32, on page 70, on page 73 at lines 5-11 and on pages 74 and 75.

Support for claims 43-56 can be found in the specification and claims as originally filed.

Support for claims 57-59 can be found in the specification in Table 13A on page 100-119.

Thus, it can be seen that no new matter has been added.

Substituted herewith pursuant to 37 CFR §1.825(a) are substitute sheets for the sequence listing provided as set forth in 37 CFR §1.821 (c). These sheets and a substitute copy of the computer readable form are enclosed herewith. The computer readable form corresponds identically to the substitute sequence listing pages submitted with this response. Support for this amendment can be found in the specification.

Specifically, the sequence listings are the same except for SEQ ID NOS: 3 and 4 which are being clarified due to sequencing errors. Support for this clarification can be found on page 36 at lines 24-27 which shows that the material has now been correctly sequenced, pCF2-165D, and was deposited with the ATCC on October 16,

009201" 62626960



Application No.: UNKNOWN  
Docket No.: BB1043 US NA DIV


1992 under the provisions of the Budapest Treaty and bears accession number ATCC 69094. Accordingly, no new matter has been added.

The clarifications in the specification were necessitated due to the clarifications in the sequence listings.

Enclosed herewith along with this Preliminary Amendment is an Information Disclosure Statement setting forth all references which had been cited by Applicants or the Examiner in connection with Serial No. 09/133,912 and some additional information as well as a petition for a three (3) month extension of time. Also, a copy of U.S. Patent No. 5,534,425 which issued to Fehr et al. on July 9, 1996 accompanies this preliminary amendment.

Please charge any fees or credit any overpayment of fees which are required in connection with the filing of this Preliminary Amendment, Information Disclosure Statement and Petition for Extension of Time to Deposit Account No. 04-1928 (E. I. du Pont de Nemours and Company).

Respectfully submitted,



Lynne M. Christenbury  
Attorney for Applicants  
Registration No. 30,971  
Telephone: 302-992-5481

Enclosures

009697379 102600

TITLE

GENES FOR MICROSOMAL DELTA-12 FATTY ACID  
DESATURASES AND RELATED ENZYMES FROM PLANTS

FIELD OF THE INVENTION

5       The invention relates to the preparation and use of  
nucleic acid fragments encoding fatty acid desaturase  
enzymes to modify plant lipid composition. Chimeric  
genes incorporating such nucleic acid fragments and  
suitable regulatory sequences may be used to create  
10       transgenic plants with altered levels of unsaturated  
fatty acids.

BACKGROUND OF THE INVENTION

Plant lipids have a variety of industrial and  
nutritional uses and are central to plant membrane  
15       function and climatic adaptation. These lipids  
represent a vast array of chemical structures, and these  
structures determine the physiological and industrial  
properties of the lipid. Many of these structures  
result either directly or indirectly from metabolic  
20       processes that alter the degree of unsaturation of the  
lipid. Different metabolic regimes in different plants  
produce these altered lipids, and either domestication  
of exotic plant species or modification of agronomically  
adapted species is usually required to economically  
25       produce large amounts of the desired lipid.

Plant lipids find their major use as edible oils in  
the form of triacylglycerols. The specific performance  
and health attributes of edible oils are determined  
largely by their fatty acid composition. Most vegetable  
30       oils derived from commercial plant varieties are  
composed primarily of palmitic (16:0), stearic (18:0),  
oleic (18:1), linoleic (18:2) and linolenic (18:3)  
acids. Palmitic and stearic acids are, respectively,  
16- and 18-carbon-long, saturated fatty acids. Oleic,  
35       linoleic, and linolenic acids are 18-carbon-long,

09697379.102600

unsaturated fatty acids containing one, two, and three double bonds, respectively. Oleic acid is referred to as a mono-unsaturated fatty acid, while linoleic and linolenic acids are referred to as poly-unsaturated fatty acids. The relative amounts of saturated and unsaturated fatty acids in commonly used, edible vegetable oils are summarized below (Table 1):

TABLE 1  
Percentages of Saturated and Unsaturated Fatty  
Acids in the Oils of Selected Oil Crops

	<u>Saturated</u>	<u>Mono- unsaturated</u>	<u>Poly- unsaturated</u>
<u>Canola</u>	6%	58%	36%
<u>Soybean</u>	15%	24%	61%
<u>Corn</u>	13%	25%	62%
<u>Peanut</u>	18%	48%	34%
<u>Safflower</u>	9%	13%	78%
<u>Sunflower</u>	9%	41%	51%
<u>Cotton</u>	30%	19%	51%

Many recent research efforts have examined the role that saturated and unsaturated fatty acids play in reducing the risk of coronary heart disease. In the past, it was believed that mono-unsaturates, in contrast to saturates and poly-unsaturates, had no effect on serum cholesterol and coronary heart disease risk. Several recent human clinical studies suggest that diets high in mono-unsaturated fat and low in saturated fat may reduce the "bad" (low-density lipoprotein) cholesterol while maintaining the "good" (high-density lipoprotein) cholesterol (Mattson et al., Journal of Lipid Research (1985) 26:194-202).

A vegetable oil low in total saturates and high in mono-unsaturates would provide significant health benefits to consumers as well as economic benefits to

oil processors. As an example, canola oil is considered a very healthy oil. However, in use, the high level of poly-unsaturated fatty acids in canola oil renders the oil unstable, easily oxidized, and susceptible to

- 5 development of disagreeable odors and flavors (Gailliard, 1980, Vol. 4, pp. 85-116 In: Stumpf, P. K., Ed., The Biochemistry of Plants, Academic Press, New York). The levels of poly-unsaturates may be reduced by hydrogenation, but the expense of this process and the  
10 concomitant production of nutritionally questionable trans isomers of the remaining unsaturated fatty acids reduces the overall desirability of the hydrogenated oil (Mensink et al., New England J. Medicine (1990) N323: 439-445). Similar problems exist with soybean and corn  
15 oils.

- For specialized uses, high levels of poly-unsaturates can be desirable. Linoleate and linolenate are essential fatty acids in human diets, and an edible oil high in these fatty acids can be used for  
20 nutritional supplements, for example in baby foods.

- Mutation-breeding programs have met with some success in altering the levels of poly-unsaturated fatty acid levels found in the edible oils of agronomic species. Examples of commercially grown varieties are  
25 high (85%) oleic sunflower and low (2%) linolenic flax (Knowles, (1980) pp. 35-38 In: Applewhite, T. H., Ed., World Conference on Biotechnology for the Fats and Oils Industry Proceedings, American Oil Chemists' Society). Similar commercial progress with the other plants shown  
30 in Table 1 has been largely elusive due to the difficult nature of the procedure and the pleiotropic effects of the mutational regime on plant hardiness and yield potential.

- The biosynthesis of the major plant lipids has been  
35 the focus of much research (Browse et al., Ann. Rev.

09697379.102600

Plant Physiol. Mol. Biol. (1991) 42:467-506). These studies show that, with the notable exception of the soluble stearyl-acyl carrier protein desaturase, the controlling steps in the production of unsaturated fatty acids are largely catalyzed by membrane-associated fatty acid desaturases. Desaturation reactions occur in plastids and in the endoplasmic reticulum using a variety of substrates including galactolipids, sulfolipids, and phospholipids. Genetic and physiological analyses of *Arabidopsis thaliana* nuclear mutants defective in various fatty acid desaturation reactions indicates that most of these reactions are catalyzed by enzymes encoded at single genetic loci in the plant. The analyses show further that the different defects in fatty acid desaturation can have profound and different effects on the ultra-structural morphology, cold sensitivity, and photosynthetic capacity of the plants (Ohlrogge, et al., Biochim. Biophys. Acta (1991) 1082:1-26). However, biochemical characterization of the desaturase reactions has been meager. The instability of the enzymes and the intractability of their proper assay has largely limited researchers to investigations of enzyme activities in crude membrane preparations. These investigations have, however, demonstrated the role of delta-12 desaturase and delta-15 desaturase activities in the production of linoleate and linolenate from 2-oleoyl-phosphatidylcholine and 2-linoleoyl-phosphatidylcholine, respectively (Wang et al., Plant Physiol. Biochem. (1988) 26:777-792). Thus, modification of the activities of these enzymes represents an attractive target for altering the levels of lipid unsaturation by genetic engineering.

Nucleotide sequences encoding microsomal delta-9 stearyl-coenzyme-A desaturases from yeast, rat, and

- mice have been described (Stukey, et al., J. Biol. Chem. (1990) 265:20144-20149; Thiede, et al., J. Biol. Chem. (1986) 261:13230-13235; Kaestner, et al., J. Biol. Chem. (1989) 264:14755-1476). Nucleotide sequences
- 5 encoding soluble delta-9 stearyl-acyl carrier protein desaturases from higher plants have also been described (Thompson, et al., Proc. Natl. Acad. Sci. U.S.A. (1991) 88:2578-2582; Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514). A nucleotide sequence from
- 10 coriander plant encoding a soluble fatty acid desaturase, whose deduced amino acid sequence is highly identical to that of the stearyl-acyl carrier protein desaturase and which is responsible for introducing the double bond in petroselinic fatty acid (18:1, 6c), has
- 15 also been described [Cahoon, et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:11184-11188]. Two fatty acid desaturase genes from the cyanobacterium, Synechocystis PCC6803, have been described: one encodes a fatty acid desaturase, designated *des A*, that catalyzes the
- 20 conversion of oleic acid at the sn-1 position of galactolipids to linoleic acid [Wada, et al., Nature (1990) 347:200-203]; another encodes a delta-6 fatty acid desaturase that catalyzes the conversion of linoleic acid at the sn-1 position of galactolipids to
- 25  $\gamma$ -linolenic acid (18:2, 6c,9c) [WO 9306712]. Nucleotide sequences encoding higher plant membrane-bound microsomal and plastid delta-15 fatty acid desaturases have also been described [WO 9311245]; Arondel, V. et. al. (1992) Science 258:1353-1355]. There is no report
- 30 of the isolation of higher plant genes encoding fatty acid desaturases other than the soluble delta-6 and delta-9 desaturases and the membrane-bound (microsomal and plastid) delta-15 desaturases. While there is extensive amino acid sequence identity between the
- 35 soluble desaturases and significant amino acid sequence

09697379.102600

identity between the higher plant microsomal and plastid delta-15 desaturases, there is no significant homology between the soluble and the membrane-bound desaturases. Sequence-dependent protocols based on the sequences encoding delta-15 desaturases have been unsuccessful in cloning sequences for microsomal delta-12 desaturase. For example, nucleotide sequences of microsomal or plastid delta-15 desaturases as hybridization probes have been unsuccessful in isolating a plant microsomal delta-12 desaturase clone. Furthermore, while we have used a set of degenerate oligomers made to a stretch of 12 amino acids, which is identical in all plant delta-15 desaturases and highly conserved (10/12) in the cyanobacterial *des A* desaturase, as a hybridization probe to isolate a higher plant nucleotide sequence encoding plastid delta-12 fatty acid desaturase, this method has been unsuccessful in isolating the microsomal delta-12 desaturase cDNAs. Furthermore, there has been no success in isolating the microsomal delta-12 desaturase by using the polymerase chain reaction products derived from plant DNA, plant RNA or plant cDNA library using PCR primers made to stretches of amino acids that are conserved between the higher plant delta-15 and *des A* desaturases. Thus, there are no teachings which enable the isolation of plant microsomal delta-12 fatty acid desaturases or plant fatty acid desaturase-related enzymes. Furthermore, there is no evidence for a method to control the the level of delta-12 fatty acid desaturation or hydroxylolation in plants using nucleic acids encoding delta-12 fatty acid desaturases or hydroxylases.

The biosynthesis of the minor plant lipids has been less well studied. While hundreds of different fatty acids have been found, many from the plant kingdom, only a tiny fraction of all plants have been surveyed for

09697379.102600

their lipid content (Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge). Accordingly, little is known about the biosynthesis of these unusual fatty acids and fatty acid derivatives.

- 5 Interesting chemical features found in such fatty acids include, for example, allenic and conjugated double bonds, acetylenic bonds, trans double bonds, multiple double bonds, and single double bonds in a wide number of positions and configurations along the fatty acid
- 10 chain. Similarly, many of the structural modifications found in unusual lipids (e.g., hydroxylation, epoxidation, cyclization, etc.) are probably produced via further metabolism following chemical activation of the fatty acid by desaturation or they involve a
- 15 chemical reaction that is mechanistically similar to desaturation. Many of these fatty acids and derivatives having such features within their structure could prove commercially useful if an agronomically viable species could be induced to synthesize them by introduction of a
- 20 gene encoding the appropriate desaturase. Of particular interest are vegetable oils rich in 12-hydroxyoctadeca-9-enoic acid (ricinoleic acid). Ricinoleic acid and its derivatives are widely used in the manufacture of lubricants, polymers, cosmetics, coatings and
- 25 pharmaceuticals (e.g., see Gunstone, et al., Eds., (1986) The Lipids Handbook, Chapman and Hall Ltd., Cambridge). The only commercial source of ricinoleic acid is castor oil and 100% of the castor oil used by the U.S. is derived from beans grown elsewhere in the
- 30 world, mainly Brazil. Ricinoleic acid in castor beans is synthesized by the addition of an hydroxyl group at the delta-12 position of oleic acid (Galliard & Stumpf (1966) J. Biol. Chem. 241: 5806-5812). This reaction resembles the initial reaction in a possible mechanism
- 35 for the desaturation of oleate at the delta-12 position

09697379, 102600



to linoleate since dehydration of 12-hydroxyoctadeca-9-  
 enoic acid, by an enzyme activity analogous to the  
 hydroxydecanoyl dehydrase of *E. coli* (Cronan et al.  
 (1988) J. Biol. Chem. 263:4641-4646), would result in  
 5 the formation of linoleic acid. Evidence for the  
 hydroxylation reaction being part of a general mechanism  
 of enzyme-catalyzed desaturation in eukaryotes has been  
 obtained by substituting a sulfur atom in the place of  
 carbon at the delta-9 position of stearic acid. When  
 10 incubated with yeast cell extracts the thiostearate was  
 converted to a 9-sulfoxide (Buist et al. (1987)  
 Tetrahedron Letters 28:857-860). This sulfoxidation was  
 specific for sulfur at the delta-9 position and did not  
 occur in a yeast delta-9-desaturase deficient mutant  
 15 (Buist & Marecak (1991) Tetrahedron Letters 32:891-894).  
 The 9-sulfoxide is the sulfur analogue of 9-hydroxyocta-  
 decastearate, the proposed intermediate of stearate  
 desaturation.

Hydroxylation of oleic acid to ricinoleic acid in  
 20 castor bean cells, like microsomal desaturation of  
 oleate in plants, occurs at the delta-12 position of the  
 fatty acid at the sn-2 position of phosphatidylcholine  
 in microsomes (Bafor et al. (1991) Plant Physiol  
 280:507-514). Furthermore, castor oleate delta-12  
 25 hydroxylation and plant oleate microsomal delta-12  
 desaturation are both inhibited by iron chelators and  
 require molecular oxygen [Moreau & Stumpf (1981) Plant  
 Physiology 67:672-676; Somerville, C. (1992) MSU-DOE  
 Plant Research Laboratory Annual Report]. These  
 30 biochemical similarities in conjunction with the  
 observation that antibodies raised against cytochrome b<sub>5</sub>  
 completely inhibit the activities of both oleate  
 delta-12 desaturation in safflower microsomes and oleate  
 delta-12 hydroxylase in castor microsomes [Somerville,  
 35 C. (1992) MSU-DOE Plant Research Laboratory Annual

09697379.102600

Report] comprise strong evidence that the hydroxylase and the desaturase are functionally related. It seems reasonable to assume, therefore, that the nucleotide sequence encoding a plant delta-12 desaturase would be

5 useful in cloning the oleate hydroxylase gene from castor by sequence-dependent protocols. For example, by screening a castor DNA library with oligomers based on amino acid regions conserved between delta-12

10 desaturases, or regions conserved between delta-12 and other desaturases, or with oligomers based on amino acids conserved between delta-12 desaturases and known membrane-associated hydroxylases. It would be more efficient to isolate the castor oleate hydroxylase cDNA by combining the sequence dependent protocols with a

15 "differential" library approach. One example of such a difference library would be based on different stages of castor seed development, since ricinoleic acid is not synthesized by very young castor seeds (less than 12 DAP, corresponding to stage I and stage II seeds in the scheme of Greenwood & Bewley, Can. J. Bot. (1982) 60:1751-1760), in the 20 days following these early stages the relative ricinoleate content increases from 0% to almost 90% of total seed fatty acids (James et al. Biochem. J. (1965) 95:448-452, Canvin. Can. J. Biochem. 25 Physiol. (1963) 41:1879-1885). Thus it would be possible to make a cDNA "difference" library made from mRNA present in a stage when ricinoleic acid was being synthesized at a high rate but from which mRNA present in earlier stages was removed. For the earlier stage

30 mRNA, a stage such as stage II (10 DAP) when ricinoleic acid is not being made but when other unsaturated fatty acids are, would be appropriate. The construction of libraries containing only differentially expressed genes is well known in the art (Sargent. Meth. Enzymol. (1987) 35 152:423-432). Assembly of the free ricinoleic acid, via

09697379.102600

ricinoleoyl-CoA, into triacylglycerol is readily catalyzed by canola and safflower seed microsomes (Bafor et al., Biochem J. (1991) 280:507-514, Wiberg et al. 10th International Symposium on the Metabolism, Structure & Function of Plant Lipids (1992), Jerba, Tunisia) and ricinoleic acid is removed from phosphatidylcholine by a lipase common to all oilseeds investigated. Thus, expression of the castor bean oleate hydroxylase gene in oil crops, such as canola seeds and soybeans, would be expected to result in an oil rich in triglycerides containing ricinoleic acid.

#### SUMMARY OF THE INVENTION

Applicants have discovered a means to control the nature and levels of unsaturated fatty acids in plants. Nucleic acid fragments from cDNAs or genes encoding fatty acid desaturases are used to create chimeric genes. The chimeric genes may be used to transform various plants to modify the fatty acid composition of the plant or the oil produced by the plant. More specifically, one embodiment of the invention is an isolated nucleic acid fragment comprising a nucleotide sequence encoding a fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50%, 60%, 90% or greater respectively to the identity of 50%, 60%, 90% or greater respectively to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15. Most specifically, the invention pertains to a gene sequence for plant microsomal delta-12 fatty acid desaturase or desaturase-related enzyme. The plant in this embodiment may more specifically be soybean, oilseed *Brassica* species, *Arabidopsis thaliana*, castor, and corn.

Another embodiment of this invention involves the use of these nucleic acid fragments in sequence-dependent protocols. Examples include use of the fragments as hybridization probes to isolate nucleotide

00697379.102600

sequences encoding other fatty acid desaturases or fatty acid desaturase-related enzymes. A related embodiment involves using the disclosed sequences for amplification of RNA or DNA fragments encoding other fatty acid desaturases or fatty acid desaturase-related enzymes.

Another aspect of this invention involves chimeric genes capable of modifying the fatty acid composition in the seed of a transformed plant, the gene comprising nucleic acid fragments related as defined to SEQ ID NOS:1, 3, 5, 7, 9, or 15 encoding fatty acid desaturases or SEQ ID NOS:11 encoding a desaturase or desaturase-related enzyme operably-linked in suitable orientation to suitable regulatory sequences. Preferred are those chimeric genes which incorporate nucleic acid fragments encoding microsomal delta-12 fatty acid desaturase or desaturase-related enzymes.

Yet another embodiment of the invention involves a method of producing seed oil containing altered levels of unsaturated fatty acids comprising: (a) transforming a plant cell with a chimeric gene described above; (b) growing sexually mature plants from the transformed plant cells of step (a); (c) screening progeny seeds from the sexually mature plants of step (b) for the desired levels of unsaturated fatty acids, and (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of the unsaturated fatty acids. Preferred plant cells and oils are derived from soybean, rapeseed, sunflower, cotton, cocoa, peanut, safflower, coconut, flax, oil palm, and corn. Preferred methods of transforming such plant cells would include the use of Ti and Ri plasmids of Agrobacterium, electroporation, and high-velocity ballistic bombardment.

The invention also is embodied in a method of RFLP breeding to obtain altered levels of oleic acids in the

00697379.102600

seed oil of oil producing plant species. This method involves (a) making a cross between two varieties of oil producing plant species differing in the oleic acid trait; (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross; and (c) hybridizing the Southern blot with the radiolabelled nucleic acid fragments encoding the fatty acid desaturases or desaturase-related enzymes.

The invention is also embodied in a method of RFLP mapping that uses the isolated microsomal delta-12 desaturase cDNA or related genomic fragments described herein.

The invention is also embodied in plants capable of producing altered levels of fatty acid desaturase by virtue of containing the chimeric genes described herein. Further, the invention is embodied by seed oil obtained from such plants.

#### BRIEF DESCRIPTION OF THE SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the Sequence Descriptions which form a part of this application. The Sequence Descriptions contain the three letter codes for amino acids as defined in 37 C.F.R. 1.822 which are incorporated herein by reference.

SEQ ID NO:1 shows the 5' to 3' nucleotide sequence of 1372 base pairs of the *Arabidopsis thaliana* cDNA which encodes microsomal delta-12 desaturase. Nucleotides 93-95 and nucleotides 1242-1244 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 93-1244). Nucleotides 1-92 and 1245-1372 are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:2 is the 383 amino acid protein sequence deduced from the open reading frame (nucleotides 93-1244 in SEQ ID NO:1.

- 5 *WU* SEQ ID NO:3 shows the 5' to 3' nucleotide sequence of 1394 base pairs of the Brassica napus cDNA which encodes microsomal delta-12 desaturase in plasmid pCF2-165d. Nucleotides 99 to 101 and nucleotides 1248 to 1250 are, respectively, the putative initiation codon and the termination codon of the open reading frame  
10 (nucleotides 99 to 1250). Nucleotides 1 to 98 and 1251 to 1394 are, respectively, the 5' and 3' untranslated nucleotides.

- SEQ ID NO:4 is the 383 amino acid protein sequence deduced from the open reading frame (nucleotides 99 to  
15 1250) in SEQ ID NO:3.

- SEQ ID NO:5 shows the 5' to 3' nucleotide sequence of 1369 base pairs of soybean (Glycine max) cDNA which encodes microsomal delta-12 desaturase in plasmid pSF2-169K. Nucleotides 108 to 110 and nucleotides 1245  
20 to 1247 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 108 to 1247). Nucleotides 1 to 107 and 1248 to 1369 are, respectively, the 5' and 3' untranslated nucleotides.

- 25 SEQ ID NO:6 is the 381 amino acid protein sequence deduced from the open reading frame (nucleotides 113 to 1258) in SEQ ID NO:5.

- SEQ ID NO:7 shows the 5' to 3' nucleotide sequence of 1790 base pairs of corn (Zea mays) cDNA which encodes  
30 microsomal delta-12 desaturase in plasmid pFad2#1. Nucleotides 165 to 167 and nucleotides 1326 to 1328 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 164 to 1328). Nucleotides 1 to 163 and 1329 to 1790

are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:8 is the 387 amino acid protein sequence deduced from the open reading frame (nucleotides 164 to 5 1328) in SEQ ID NO:7.

SEQ ID NO:9 shows the 5' to 3' nucleotide sequence of 673 base pairs of castor (*Ricinus communis*) incomplete cDNA which encodes part of a microsomal delta-12 desaturase in plasmid pRF2-1C. The sequence 10 encodes an open reading frame from base 1 to base 673..

SEQ ID NO:10 is the 219 amino acid protein sequence deduced from the open reading frame (nucleotides 1 to 657) in SEQ ID NO:9.

SEQ ID NO:11 shows the 5' to 3' nucleotide sequence 15 of 1369 base pairs of castor (*Ricinus communis*) cDNA which encodes part of a microsomal delta-12 desaturase or desaturase-related enzyme in plasmid pRF197C-42. Nucleotides 184 to 186 and nucleotides 1340 to 1342 are, respectively, the putative initiation codon and the 20 termination codon of the open reading frame (nucleotides 184 to 1347). Nucleotides 1 to 183 and 1348 to 1369 are, respectively, the 5' and 3' untranslated nucleotides.

SEQ ID NO:12 is the 387 amino acid protein sequence 25 deduced from the open reading frame (nucleotides 184 to 1342) in SEQ ID NO:11.

SEQ ID NO:13 is the sequence of a set of 64-fold degenerate 26 nucleotide-long oligomers, designated NS3, made to conserved amino acids 101-109 of SEQ ID NO:2, 30 designed to be used as sense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

SEQ ID NO:14 is the sequence of a set of 64-fold degenerate and 26 nucleotide-long oligomers, designated 35 NS9, which is made to conserved amino acids 313-321 of

09697379.102600

SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

5 SEQ ID NO:15 shows the 5' to 3' nucleotide sequence of 2973 bp of Arabidopsis thaliana genomic fragment containing the microsomal delta-12 desaturase gene contained in plasmid pAGF2-6. Its nucleotides 433 and 2938 correspond to the start and end, respectively, of SEQ ID NO:1. Its nucleotides 521 to 1654 are the 1134  
10 bp intron.

SEQ ID NO:16 is the sequence of a set of 256-fold degenerate and 25 nucleotide-long oligomers, designated RB5a, which is made to conserved amino acids 318-326 of  
15 SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

SEQ ID NO:17 is the sequence of a set of 128-fold degenerate and 25 nucleotide-long oligomers, designated RB5b, which is made to conserved amino acids 318-326 of  
20 SEQ ID NO:2 and designed to be used as antisense primers in PCR to isolate novel sequences encoding microsomal delta-12 desaturases or desaturase-like enzymes.

#### DETAILED DESCRIPTION OF THE INVENTION

Applicants have isolated nucleic acid fragments  
25 that encode plant fatty acid desaturases and that are useful in modifying fatty acid composition in oil-producing species by genetic transformation.

Thus, transfer of the nucleic acid fragments of the invention or a part thereof that encodes a functional  
30 enzyme, along with suitable regulatory sequences that direct the transcription of their mRNA, into a living cell will result in the production or over-production of plant fatty acid desaturases and will result in increased levels of unsaturated fatty acids in cellular  
35 lipids, including triacylglycerols.

09697379.102600



Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their antisense RNA, into plants will result in the inhibition of expression of the endogenous fatty acid desaturase that is substantially homologous with the transferred nucleic acid fragment and will result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

Transfer of the nucleic acid fragments of the invention or a part thereof, along with suitable regulatory sequences that direct the transcription of their mRNA, into plants may result in inhibition by cosuppression of the expression of the endogenous fatty acid desaturase gene that is substantially homologous with the transferred nucleic acid fragment and may result in decreased levels of unsaturated fatty acids in cellular lipids, including triacylglycerols.

The nucleic acid fragments of the invention can also be used as restriction fragment length polymorphism (RFLP) markers in plant genetic mapping and plant breeding programs.

The nucleic acid fragments of the invention or oligomers derived therefrom can also be used to isolate other related fatty acid desaturase genes using DNA, RNA, or a library of cloned nucleotide sequences from the same or different species by well known sequence-dependent protocols, including, for example, methods of nucleic acid hybridization and amplification by the polymerase chain reaction.

#### Definitions

In the context of this disclosure, a number of terms shall be used. Fatty acids are specified by the number of carbon atoms and the number and position of the double bond: the numbers before and after the colon

00697379.102600

- refer to the chain length and the number of double bonds, respectively. The number following the fatty acid designation indicates the position of the double bond from the carboxyl end of the fatty acid with the "c" affix for the *cis*-configuration of the double bond. For example, palmitic acid (16:0), stearic acid (18:0), oleic acid (18:1,9c), petroselinic acid (18:1, 6c), linoleic acid (18:2,9c,12c),  $\gamma$ -linolenic acid (18:3, 6c,9c,12c) and  $\alpha$ -linolenic acid (18:3, 9c,12c,15c).
- Unless otherwise specified 18:1, 18:2 and 18:3 refer to oleic, linoleic and linolenic fatty acids. Ricinoleic acid refers to an 18 carbon fatty acid with a *cis*-9 double bond and a 12-hydroxyl group. The term "fatty acid desaturase" used herein refers to an enzyme which catalyzes the breakage of a carbon-hydrogen bond and the introduction of a carbon-carbon double bond into a fatty acid molecule. The fatty acid may be free or esterified to another molecule including, but not limited to, acyl-carrier protein, coenzyme A, sterols and the glycerol moiety of glycerolipids. The term "glycerolipid desaturases" used herein refers to a subset of the fatty acid desaturases that act on fatty acyl moieties esterified to a glycerol backbone. "Delta-12 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 6 and 7 (numbered from the methyl end), (i.e., those that correspond to carbon positions 12 and 13 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. "Delta-15 desaturase" refers to a fatty acid desaturase that catalyzes the formation of a double bond between carbon positions 3 and 4 (numbered from the methyl end), (i.e., those that correspond to carbon positions 15 and 16 (numbered from the carbonyl carbon) of an 18 carbon-long fatty acyl chain. Examples of fatty acid desaturases include, but are not limited

09697379.102500

would be particularly useful. The consensus sequences will also take into account conservative substitutions known to one skilled in the art, such as Lys/Arg, Glu/Asp, Ile/Val/Leu/Met, Ala/Gly, Gln/Asn, and Ser/Thr.

5 Amino acid sequences as short as four amino acids long can successfully be used in PCR [Nunberg et. al. (1989) Journal of Virology 63:3240-3249]. Amino acid sequences conserved between delta-12 desaturases (SEQ ID NOS:2, 4, 6, 8, and 10) may also be used in sequence-dependent

10 protocols to isolate fatty acid desaturases and fatty acid desaturase-related enzymes expected to be more related to delta-12 desaturases, such as the oleate hydroxylase from castor bean. Particularly useful are conserved sequences in column 3 (Table 7), since they

15 are also conserved well with delta-15 desaturases (column 4, Table 7).

Determining the conserved amino acid sequences from diverse desaturases will also allow one to identify more and better consensus sequences that will further aid in

20 the isolation of novel fatty acid desaturases, including those from non-plant sources such as fungi, algae (including the desaturases involved in the desaturations of the long chain n-3 fatty acids), and even cyano-bacteria, as well as other membrane-associated

25 desaturases from other organisms.

The function of the diverse nucleotide fragments encoding fatty acid desaturases or desaturase-related enzymes that can be isolated using the present invention can be identified by transforming plants with the

30 isolated sequences, linked in sense or antisense orientation to suitable regulatory sequences required for plant expression, and observing the fatty acid phenotype of the resulting transgenic plants. Preferred target plants for the transformation are the same as the

35 source of the isolated nucleotide fragments when the

- goal is to obtain inhibition of the corresponding endogenous gene by antisense inhibition or cosuppression. Preferred target plants for use in expression or overexpression of the isolated nucleic acid fragments are wild type plants or plants with known mutations in desaturation reactions, such as the Arabidopsis mutants fadA, fadB, fadC, fadD, fad2, and fad3; mutant flax deficient in delta-15 desaturation; or mutant sunflower deficient in delta-12 desaturation.
- Alternatively, the function of the isolated nucleic acid fragments can be determined similarly via transformation of other organisms, such as yeast or cyanobacteria, with chimeric genes containing the nucleic acid fragment and suitable regulatory sequences followed by analysis of fatty acid composition and/or enzyme activity.

#### Overexpression of the Fatty Acid

##### Desaturase Enzymes in Transgenic Species

- The nucleic acid fragment(s) of the instant invention encoding functional fatty acid desaturase(s), with suitable regulatory sequences, can be used to overexpress the enzyme(s) in transgenic organisms. An example of such expression or overexpression is demonstrated by transformation of the Arabidopsis mutant lacking oleate desaturation. Such recombinant DNA constructs may include either the native fatty acid desaturase gene or a chimeric fatty acid desaturase gene isolated from the same or a different species as the host organism. For overexpression of fatty acid desaturase(s), it is preferable that the introduced gene be from a different species to reduce the likelihood of cosuppression. For example, overexpression of delta-12 desaturase in soybean, rapeseed, or other oil-producing species to produce altered levels of polyunsaturated fatty acids may be achieved by expressing RNA from the full-length cDNA found in p92103, pCF2-165D, and

09697379.102600

09697379.102600

pSF2-169K. Transgenic lines overexpressing the delta-12 desaturase, when crossed with lines overexpressing delta-15 desaturases, will result in ultrahigh levels of 18:3. Similarly, the isolated nucleic acid fragments  
5 encoding fatty acid desaturases from Arabidopsis, rapeseed, and soybean can also be used by one skilled in the art to obtain other substantially homologous full-length cDNAs, if not already obtained, as well as the corresponding genes as fragments of the invention.  
10 These, in turn, may be used to overexpress the corresponding desaturases in plants. One skilled in the art can also isolate the coding sequence(s) from the fragment(s) of the invention by using and/or creating sites for restriction endonucleases, as described in  
15 Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

One particularly useful application of the claimed inventions is to repair the agronomic performance of,  
20 plant mutants containing ultra high levels of oleate in seed oil. In Arabidopsis reduction in linoleate in phosphatidylcholine due to a mutation in microsomal delta-12 desaturase affected low temperature survival [Miquel, M. et. al. (1993) Proc. Natl Acad. Sci. USA  
25 90:6208-6212]. Furthermore, there is evidence that the poor agronomic performance of canola plants containing ultra high (>80%) levels of oleate in seed is due to mutations in the microsomal delta-12 desaturase genes that reduce the level of linoleate in phosphatidyl-  
30 choline of roots and leaves. That is, the mutations are not seed-specific. Thus, the root and/or leaf-specific expression (that is, with no expression in the seeds) of microsomal delta-12 desaturase activity in mutants of oilseeds with ultra-high levels of oleate in seed oil

will result in agronomically-improved mutant plants with ultra high levels of oleate in seed oil.

Inhibition of Plant Target  
Genes by Use of Antisense RNA

5           Antisense RNA has been used to inhibit plant target genes in a tissue-specific manner (see van der Krol et al., *Biotechniques* (1988) 6:958-976). Antisense inhibition has been shown using the entire cDNA sequence (Sheehy et al., *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809) as well as a partial cDNA sequence (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47). There is also evidence that the 3' non-coding sequences (Ch'ng et al., *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010) and fragments of 5' coding sequence, containing as few as 41 base-pairs of a 1.87 kb cDNA (Cannon et al., *Plant Molec. Biol.* (1990) 15:39-47), can play important roles in antisense inhibition.

          The use of antisense inhibition of the fatty acid desaturases may require isolation of the transcribed sequence for one or more target fatty acid desaturase genes that are expressed in the target tissue of the target plant. The genes that are most highly expressed are the best targets for antisense inhibition. These genes may be identified by determining their levels of transcription by techniques, such as quantitative analysis of mRNA levels or nuclear run-off transcription, known to one skilled in the art.

          The entire soybean microsomal delta-12 desaturase cDNA was cloned in the antisense orientation with respect to either soybean b-conglycinin, soybean KTi3, and bean phaseolin promoter and the chimeric gene transformed into soybean somatic embryos that were previously shown to serve as good model system for soybean zygotic embryos and are predictive of seed composition (Table 11). Transformed somatic embryos

009697379.102600

showed inhibition of linoleate biosynthesis. Similarly, the entire Brassica napus microsomal delta-12 desaturase cDNA was cloned in the antisense orientation with respect to a rapeseed napin promoter and the chimeric gene transformed into B. napus. Seeds of transformed B. napus plants showed inhibition of linoleate biosynthesis. Thus, antisense inhibition of delta-12 desaturase in oil-producing species, including corn, Brassica napus, and soybean resulting in altered levels of polyunsaturated fatty acids may be achieved by expressing antisense RNA from the entire or partial cDNA encoding microsomal delta-12 desaturase.

#### Inhibition of Plant

##### Target Genes by Cosuppression

The phenomenon of cosuppression has also been used to inhibit plant target genes in a tissue-specific manner. Cosuppression of an endogenous gene using the entire cDNA sequence (Napoli et al., The Plant Cell (1990) 2:279-289; van der Krol et al., The Plant Cell (1990) 2:291-299) as well as a partial cDNA sequence (730 bp of a 1770 bp cDNA) (Smith et al., Mol. Gen. Genetics (1990) 224:477-481) are known.

The nucleic acid fragments of the instant invention encoding fatty acid desaturases, or parts thereof, with suitable regulatory sequences, can be used to reduce the level of fatty acid desaturases, thereby altering fatty acid composition, in transgenic plants which contain an endogenous gene substantially homologous to the introduced nucleic acid fragment. The experimental procedures necessary for this are similar to those described above for the overexpression of the fatty acid desaturase nucleic acid fragments except that one may also use a partial cDNA sequence. For example, cosuppression of delta-12 desaturase in Brassica napus and soybean resulting in altered levels of

09697379.102600

polyunsaturated fatty acids may be achieved by expressing in the sense orientation the entire or partial seed delta-12 desaturase cDNA found in pCF2-165D and pSF2-165K, respectively. Endogenous genes can also  
 5 be inhibited by non-coding regions of an introduced copy of the gene [For example, Brusslan, J. A. et al. (1993) Plant Cell 5:667-677; Matzke, M. A. et al., Plant Molecular Biology 16:821-830]. We have shown that an Arabidopsis gene (SEQ ID NO:15) corresponding to the  
 10 cDNA (SEQ ID NO:1) can be isolated. One skilled in the art can readily isolate genomic DNA containing or flanking the genes and use the coding or non-coding regions in such transgene inhibition methods.

Analysis of the fatty acid composition of roots and  
 15 seeds of Arabidopsis mutants deficient in microsomal delta-12 desaturation shows that they have reduced levels of 18:2 as well as reduced levels of 16:0 (as much as 40% reduced level in mutant seeds as compared to wild type seeds) [Miquel and Browse (1990) in Plant  
 20 Lipid Biochemistry, Structure, and Utilization, pages 456-458, Ed. Quinn, P. J. and Harwood, J. L., Portland Press. Reduction in the level of 16:0 is also observed in ultra high oleate mutants of B. napus. Thus, one can expect that ultra high level of 18:1 in  
 25 transgenic plants due to antisense inhibition or co-suppression using the claimed sequences will also reduce the level of 16:0.

One particularly useful application of the claimed invention is to combine the high oleate trait of the  
 30 transformed seeds with mutations for altered fatty acid compositions to obtain novel fatty acid compositions and/or improved agronomy. Fatty acid compositions, such as ultra-high oleate levels (EPO 323753 and see below), resulting from mutagenesis alone, such as by combining  
 35 mutations, may be affected in agronomy or seed

09597379.102600



germination. Transgene and mutant genes may be combined either by making a genetic cross between the two lines or by transforming the chimeric genes of the inventions into the mutant lines as known to one skilled in the art.

As examples, the high oleic acid trait of the invention can be combined with high stearate mutations, such as soybean A6 [Hammond, E. G. and Fehr, W. R. (1983)] or low linolenic acid mutations and soybean mutants A5, A23, A16 and C1640 [Fehr, W. R. et al. (1992) in Crop Science 32:903-906]. Oils produced from such combinations will provide improved feedstocks for production of margarines, shortenings, spray coating and frying oils and will eliminate or reduce the need for hydrogenation. Furthermore, these oils would provide a health benefit for consumers, for example by reducing or eliminating trans fatty acids which have recently been found to be associated with high risk to cardiovascular diseases.

It can also be combined with other high oleic acid mutants that will result in even higher oleic acid content. Examples of high oleate mutants include soybean lines A5 and N782245 [Martin, B. A. and Rinne, R. W. (1985) Crop Science 25:1055-1058] and rapeseed lines containing about 69% to 77% oleic acid. A mutant (IMC 129) was derived by chemical mutagenesis of Westar and is presently in commercial production in the U.S. This variety has yields comparable to Westar, a leading Canadian variety. Its high oleate trait can be attributed to a mutation found in the coding region (Glu<sub>106</sub> to Lys<sub>106</sub>) one of the two delta-12 desaturase structural genes found in Westar. Re-mutagenesis of this line resulted in a second mutation in the other delta-12 desaturase structural gene and even higher seed oleic acid content in the double mutant. However the

00697379, 102600

double mutant has significantly higher levels of oleic acid in its leaves and roots than Westar, i.e., the mutant phenotype is not seed specific. The double mutant has extremely poor agronomy characteristics, which appear to be related to the high oleate in its leaves and/or roots. Thus, one can combine the seed-specific high oleic acid content resulting from seed-specific antisense inhibition or cosuppression of the microsomal delta-12 desaturase in transgenic seeds with that in the single mutant with 69-77% oleic acid to obtain levels of seed oleic acid that are as high or higher than that in the double mutant without affecting agronomy. The increased levels of oleic acid in the double homozygotes are associated with reduced levels of saturates. The reduction in saturates is also observed with seed-specific inhibition of delta-12 desaturase in soybean seeds.

#### Selection of Hosts, Promoters and Enhancers

A preferred class of heterologous hosts for the expression of the nucleic acid fragments of the invention are eukaryotic hosts, particularly the cells of higher plants. Particularly preferred among the higher plants are the oil-producing species, such as soybean (Glycine max), rapeseed (including Brassica napus, B. campestris), sunflower (Helianthus annuus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), and peanut (Arachis hypogaea).

Expression in plants will use regulatory sequences functional in such plants. The expression of foreign genes in plants is well-established (De Blaere et al., Meth. Enzymol. (1987) 153:277-291). The source of the promoter chosen to drive the expression of the fragments

of the invention is not critical provided it has sufficient transcriptional activity to accomplish the invention by increasing or decreasing, respectively, the level of translatable mRNA for the fatty acid

5 desaturases in the desired host tissue. Preferred promoters include (a) strong constitutive plant promoters, such as those directing the 19S and 35S transcripts in cauliflower mosaic virus (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987)

10 86:482-493), (b) tissue- or developmentally-specific promoters, and (c) other transcriptional promoter systems engineered in plants, such as those using bacteriophage T7 RNA polymerase promoter sequences to express foreign genes. Examples of tissue-specific

15 promoters are the light-inducible promoter of the small subunit of ribulose 1,5-bis-phosphate carboxylase (if expression is desired in photosynthetic tissues), the maize zein protein promoter (Matzke et al., EMBO J. (1984) 3:1525-1532), and the chlorophyll a/b binding

20 protein promoter (Lampa et al., Nature (1986) 316:750-752).

Particularly preferred promoters are those that allow seed-specific expression. This may be especially useful since seeds are the primary source of vegetable

25 oils and also since seed-specific expression will avoid any potential deleterious effect in non-seed tissues. Examples of seed-specific promoters include, but are not limited to, the promoters of seed storage proteins, which can represent up to 90% of total seed protein in

30 many plants. The seed storage proteins are strictly regulated, being expressed almost exclusively in seeds in a highly tissue-specific and stage-specific manner (Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221; Goldberg et al., Cell (1989) 56:149-160).

Moreover, different seed storage proteins may be expressed at different stages of seed development.

- Expression of seed-specific genes has been studied in great detail (See reviews by Goldberg et al., Cell (1989) 56:149-160 and Higgins et al., Ann. Rev. Plant Physiol. (1984) 35:191-221). There are currently numerous examples of seed-specific expression of seed storage protein genes in transgenic dicotyledonous plants. These include genes from dicotyledonous plants for bean b-phaseolin (Sengupta-Gopalan et al., Proc. Natl. Acad. Sci. USA (1985) 82:3320-3324; Hoffman et al., Plant Mol. Biol. (1988) 11:717-729), bean lectin (Voelker et al., EMBO J. (1987) 6:3571-3577), soybean lectin (Okamuro et al., Proc. Natl. Acad. Sci. USA (1986) 83:8240-8244), soybean Kunitz trypsin inhibitor (Perez-Grau et al., Plant Cell (1989) 1:095-1109), soybean b-conglycinin (Beachy et al., EMBO J. (1985) 4:3047-3053; pea vicilin (Higgins et al., Plant Mol. Biol. (1988) 11:683-695), pea convicilin (Newbigin et al., Planta (1990) 180:461-470), pea legumin (Shirsat et al., Mol. Gen. Genetics (1989) 215:326-331); rapeseed napin (Radke et al., Theor. Appl. Genet. (1988) 75:685-694) as well as genes from monocotyledonous plants such as for maize 15 kD zein (Hoffman et al., EMBO J. (1987) 6:3213-3221), maize 18 kD oleosin (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 888:6181-6185), barley b-hordein (Marris et al., Plant Mol. Biol. (1988) 10:359-366) and wheat glutenin (Colot et al., EMBO J. (1987) 6:3559-3564). Moreover, promoters of seed-specific genes operably linked to heterologous coding sequences in chimeric gene constructs also maintain their temporal and spatial expression pattern in transgenic plants. Such examples include use of *Arabidopsis thaliana* 2S seed storage protein gene promoter to express enkephalin peptides in *Arabidopsis*

and B. napus seeds (Vandekerckhove et al.,  
 Bio/Technology (1989) 7:929-932), bean lectin and bean  
 b-phaseolin promoters to express luciferase (Riggs et  
 al., Plant Sci. (1989) 63:47-57), and wheat glutenin  
 5 promoters to express chloramphenicol acetyl transferase  
 (Colot et al., EMBO J. (1987) 6:3559-3564).

Of particular use in the expression of the nucleic  
 acid fragment of the invention will be the heterologous  
 promoters from several soybean seed storage protein  
 10 genes such as those for the Kunitz trypsin inhibitor  
 (Jofuku et al., Plant Cell (1989) 1:1079-1093; glycinin  
 (Nielson et al., Plant Cell (1989) 1:313-328), and  
 b-conglycinin (Harada et al., Plant Cell (1989)  
 1:415-425). Promoters of genes for a- and b-subunits of  
 15 soybean b-conglycinin storage protein will be  
 particularly useful in expressing the mRNA or the  
 antisense RNA in the cotyledons at mid- to late-stages  
 of seed development (Beachy et al., EMBO J. (1985)  
 4:3047-3053) in transgenic plants. This is because  
 20 there is very little position effect on their expression  
 in transgenic seeds, and the two promoters show  
 different temporal regulation. The promoter for the  
 a-subunit gene is expressed a few days before that for  
 the b-subunit gene. This is important for transforming  
 25 rapeseed where oil biosynthesis begins about a week  
 before seed storage protein synthesis (Murphy et al.,  
 J. Plant Physiol. (1989) 135:63-69).

Also of particular use will be promoters of genes  
 expressed during early embryogenesis and oil bio-  
 30 synthesis. The native regulatory sequences, including  
 the native promoters, of the fatty acid desaturase genes  
 expressing the nucleic acid fragments of the invention  
 can be used following their isolation by those skilled  
 in the art. Heterologous promoters from other genes  
 35 involved in seed oil biosynthesis, such as those for

09697379.102600

- B. napus* isocitrate lyase and malate synthase (Comai et al., Plant Cell (1989) 1:293-300), delta-9 desaturase from safflower (Thompson et al. Proc. Natl. Acad. Sci. USA (1991) 88:2578-2582) and castor (Shanklin et al., Proc. Natl. Acad. Sci. USA (1991) 88:2510-2514), acyl carrier protein (ACP) from *Arabidopsis* (Post-Beittenmiller et al., Nucl. Acids Res. (1989) 17:1777), *B. napus* (Safford et al., Eur. J. Biochem. (1988) 174:287-295), and *B. campestris* (Rose et al., Nucl. Acids Res. (1987) 15:7197), b-ketoacyl-ACP synthetase from barley (Siggaard-Andersen et al., Proc. Natl. Acad. Sci. USA (1991) 88:4114-4118), and oleosin from *Zea mays* (Lee et al., Proc. Natl. Acad. Sci. USA (1991) 88:6181-6185), soybean (Genbank Accession No: X60773) and *B. napus* (Lee et al., Plant Physiol. (1991) 96:1395-1397) will be of use. If the sequence of the corresponding genes is not disclosed or their promoter region is not identified, one skilled in the art can use the published sequence to isolate the corresponding gene and a fragment thereof containing the promoter. The partial protein sequences for the relatively-abundant enoyl-ACP reductase and acetyl-CoA carboxylase are also published (Slabas et al., Biochim. Biophys. Acta (1987) 877:271-280; Cottingham et al., Biochim. Biophys. Acta (1988) 954:201-207) and one skilled in the art can use these sequences to isolate the corresponding seed genes with their promoters. Similarly, the fragments of the present invention encoding fatty acid desaturases can be used to obtain promoter regions of the corresponding genes for use in expressing chimeric genes.

Attaining the proper level of expression of the nucleic acid fragments of the invention may require the use of different chimeric genes utilizing different promoters. Such chimeric genes can be transferred into

host plants either together in a single expression vector or sequentially using more than one vector.

It is envisioned that the introduction of enhancers or enhancer-like elements into the promoter regions of either the native or chimeric nucleic acid fragments of the invention will result in increased expression to accomplish the invention. This would include viral enhancers such as that found in the 35S promoter (Odell et al., Plant Mol. Biol. (1988) 10:263-272), enhancers from the opine genes (Fromm et al., Plant Cell (1989) 1:977-984), or enhancers from any other source that result in increased transcription when placed into a promoter operably linked to the nucleic acid fragment of the invention.

Of particular importance is the DNA sequence element isolated from the gene for the a-subunit of b-conglycinin that can confer 40-fold seed-specific enhancement to a constitutive promoter (Chen et al., Dev. Genet. (1989) 10:112-122). One skilled in the art can readily isolate this element and insert it within the promoter region of any gene in order to obtain seed-specific enhanced expression with the promoter in transgenic plants. Insertion of such an element in any seed-specific gene that is expressed at different times than the b-conglycinin gene will result in expression in transgenic plants for a longer period during seed development.

The invention can also be accomplished by a variety of other methods to obtain the desired end. In one form, the invention is based on modifying plants to produce increased levels of fatty acid desaturases by virtue of introducing more than one copy of the foreign gene containing the nucleic acid fragments of the invention. In some cases, the desired level of polyunsaturated fatty acids may require introduction of

09697379.102600

foreign genes for more than one kind of fatty acid desaturase.

- Any 3' non-coding region capable of providing a polyadenylation signal and other regulatory sequences that may be required for the proper expression of the nucleic acid fragments of the invention can be used to accomplish the invention. This would include 3' ends of the native fatty acid desaturase(s), viral genes such as from the 35S or the 19S cauliflower mosaic virus transcripts, from the opine synthesis genes, ribulose 1,5-bisphosphate carboxylase, or chlorophyll a/b binding protein. There are numerous examples in the art that teach the usefulness of different 3' non-coding regions.

#### Transformation Methods

- Various methods of transforming cells of higher plants according to the present invention are available to those skilled in the art (see EPO Pub. 0 295 959 A2 and 0 318 341 A1). Such methods include those based on transformation vectors utilizing the Ti and Ri plasmids of Agrobacterium spp. It is particularly preferred to use the binary type of these vectors. Ti-derived vectors transform a wide variety of higher plants, including monocotyledonous and dicotyledonous plants (Sukhapinda et al., Plant Mol. Biol. (1987) 8:209-216; Potrykus, Mol. Gen. Genet. (1985) 199:183). Other transformation methods are available to those skilled in the art, such as direct uptake of foreign DNA constructs (see EPO Pub. 0 295 959 A2), techniques of electroporation (Fromm et al., Nature (1986) (London) 319:791) or high-velocity ballistic bombardment with metal particles coated with the nucleic acid constructs (Kline et al., Nature (1987) (London) 327:70). Once transformed, the cells can be regenerated by those skilled in the art.



Of particular relevance are the recently described methods to transform foreign genes into commercially important crops, such as rapeseed (De Block et al., Plant Physiol. (1989) 91:694-701), sunflower (Everett et al., Bio/Technology (1987) 5:1201), and soybean (Christou et al., Proc. Natl. Acad. Sci USA (1989) 86:7500-7504).

#### Application to Molecular Breeding

The 1.6 kb insert obtained from the plasmid pSF2-169K was used as a radiolabelled probe on a Southern blot containing genomic DNA from soybean (*Glycine max* (cultivar Bonus) and *Glycine soja* (PI81762)) digested with one of several restriction enzymes. Different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Hind III and Eco RI. These polymorphisms were used to map two pSF2-169 loci relative to other loci on the soybean genome essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). One mapped to linkage group 11 between 4404.00 and 1503.00 loci (4.5 cM and 7.1 cM from 4404.00 and 1503.00, respectively) and the other to linkage group 19 between 4010.00 and 5302.00 loci (1.9 cM and 2.7 cM from 4010.00 and 5302.00, respectively) [Rafalski, A and Tingey, S. (1993) in Genetic Maps, Ed. O' Brien, S. J.]. The use of restriction fragment length polymorphism (RFLP) markers in plant breeding has been well-documented in the art (Tanksley et al., Bio/Technology (1989) 7:257-264). Thus, the nucleic acid fragments of the invention can be used as RFLP markers for traits linked to expression of fatty acid desaturases. These traits will include altered levels of unsaturated fatty acids. The nucleic acid fragment of the invention can also be used to isolate the fatty acid desaturase gene from variant

09697379-102600

(including mutant) plants with altered levels of unsaturated fatty acids. Sequencing of these genes will reveal nucleotide differences from the normal gene that cause the variation. Short oligonucleotides designed around these differences may also be used in molecular breeding either as hybridization probes or in DNA-based diagnostics to follow the variation in fatty acids. Oligonucleotides based on differences that are linked to the variation may be used as molecular markers in breeding these variant oil traits.

#### EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. All publications, including patents and non-patent literature, referred to in this specification are expressly incorporated by reference herein.

#### EXAMPLE 1

##### ISOLATION OF GENOMIC DNA FLANKING THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA MUTANT LINE 658

##### 30 Identification of an Arabidopsis thaliana T-DNA Mutant with High Oleic Acid Content

A population of Arabidopsis thaliana (geographic race Wassilewskija) transformants containing the modified T-DNA of Agrobacterium tumefaciens was generated by seed transformation as described by

Feldmann et al., (Mol. Gen. Genetics (1987) 208:1-9). In this population the transformants contain DNA sequences encoding the pBR322 bacterial vector, nopaline synthase, neomycin phosphotransferase (NPTII, confers  
 5 kanamycin resistance), and b-lactamase (confers ampicillin resistance) within the T-DNA border sequences. The integration of the T-DNA into different areas of the chromosomes of individual transformants may cause a disruption of plant gene function at or near the  
 10 site of insertion, and phenotypes associated with this loss of gene function can be analyzed by screening the population for the phenotype.

T3 seed was generated from the wild type seed treated with Agrobacterium tumefaciens by two rounds of  
 15 self-fertilization as described by Feldmann et al., (Science (1989) 243:1351-1354). These progeny were segregating for the T-DNA insertion, and thus for any mutation resulting from the insertion. Approximately 10-12 leaves of each of 1700 lines were combined and the  
 20 fatty acid content of each of the 1700 pooled samples was determined by gas chromatography of the fatty acyl methyl esters essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent.  
 25 A line designated "658" produced a sample that gave an altered fatty acid profile compared to those of lines 657 and 659 sampled at the same time (Table 8).

TABLE 8

Fatty Acid Methyl Ester	657 Leaf Pool	659 Leaf Pool	658 Leaf Pool
16:0	14.4	14.1	13.6
16:1	4.4	4.6	4.5
16:2	2.9	2.2	2.7
16:3	13.9	13.3	13.9

009697379-102600

18:0	1.0	1.1	0.9
18:1	2.6	2.5	4.9
18:2	14.0	13.6	12.8
18:3	42.9	46.1	44.4

- Analysis of the fatty acid composition of 12 individual T3 seeds of line 658 indicated that the 658 pool was composed of seeds segregating in three classes: "high", "mid-range" and "low" classes with
- 5 approximately, 37% (12 seeds), 21% (7 seeds), and 14% (3 seeds) oleic acid, respectively (Table 9).

TABLE 9

	<u>"High"</u> <u>Class</u>	<u>"Mid-range"</u> <u>Class</u>	<u>"Low"</u> <u>Class</u>
16:0	8.9	8.7	9.3
16:1c	2.0	1.6	2.6
18:0	4.5	4.3	4.4
18:1	37.0	20.7	14.4
18:2	8.0	24.9	27.7
18:3	10.6	14.3	13.6
20:1	25.5	21.6	20.4

- Thus, the high oleic acid mutant phenotype segregates in an approximately Mendelian ratio. To determine the number of independently segregating T-DNA
- 10 inserts in line 658, 200 T3 seeds were tested for their ability to germinate and grow in the presence of kanamycin [Feldman et al. (1989) Science 243:1351-1354]. In this experiment, only 4 kanamycin-sensitive
- 15 individual plants were identified. The segregation ratio (approximately 50:1) indicated that line 658 harbored three T-DNA inserts. In this and two other experiments a total of 56 kanamycin-sensitive plants were identified; 53 of these were analyzed for fatty

acid composition and at least seven of these displayed oleic acid levels that were higher than would be expected for wild type seedlings grown under these conditions.

- 5        In order to more rigorously test whether the mutation resulting in high oleic acid is the result of T-DNA insertion, Applicants identified a derivative line that was segregating for the mutant fatty acid phenotype as well as a single kanamycin resistance locus. For
- 10 this, approximately 100 T3 plants were individually grown to maturity and seeds collected. One sample of seed from each T3 plant was tested for the ability to germinate and grow in the presence of kanamycin. In addition, the fatty acid compositions of ten additional
- 15 individual seeds from each line were determined. A T3 plant, designated 658-75, was identified whose progeny seeds segregated 28 kanamycin-sensitive to 60 kanamycin-resistant and 7 with either low or intermediate oleic acid to 2 high oleic acid.
- 20        A total of approximately 400 T4 progeny seeds of the derivative line 658-75 were grown and the leaf fatty acid composition analyzed. A total of 91 plants were identified as being homozygous for the high oleic acid trait (18:2/18:1 less than 0.5). The remaining plants
- 25 (18:2/18:1 more than 1.2) could not be definitively assigned to wild type and heterozygous classes on the basis of leaf fatty acid composition and thus could not be used to test linkage between the kanamycin marker and the fatty acid mutation. Eighty three of the 91
- 30 apparently homozygous high oleic acid mutant were tested for the presence of nopaline, another T-DNA marker, in leaf extracts (Errampalli et al. The Plant Cell (1991)3:149-157 and all 83 plants were positive for the presence of nopaline. This tight linkage of the mutant
- 35 fatty acid phenotype and a T-DNA marker provides

to, the microsomal delta-12 and delta-15 desaturases that act on phosphatidylcholine lipid substrates; the chloroplastic or plastid delta-12 and delta-15 desaturases that act on phosphatidyl glycerol and galactolipids; and other desaturases that act on such fatty acid substrates such as phospholipids, galactolipids, and sulfolipids. "Microsomal desaturase" refers to the cytoplasmic location of the enzyme, while "chloroplast desaturase" and "plastid desaturase" refer to the plastid location of the enzyme. These fatty acid desaturases may be found in a variety of organisms including, but not limited to, higher plants, diatoms, and various eukaryotic and prokaryotic microorganisms such as fungi and photosynthetic bacteria and algae.

The term "homologous fatty acid desaturases" refers to fatty acid desaturases that catalyze the same desaturation on the same lipid substrate. Thus, microsomal delta-15 desaturases, even from different plant species, are homologous fatty acid desaturases.

The term "heterologous fatty acid desaturases" refers to fatty acid desaturases that catalyze desaturations at different positions and/or on different lipid substrates. Thus, for example, microsomal delta-12 and delta-15 desaturases, which act on phosphatidylcholine lipids, are heterologous fatty acid desaturases, even when from the same plant. Similarly, microsomal delta-15 desaturase, which acts on phosphatidylcholine lipids, and chloroplast delta-15 desaturase, which acts on galactolipids, are heterologous fatty acid desaturases, even when from the same plant. It should be noted that these fatty acid desaturases have never been isolated and characterized as proteins. Accordingly, the terms such as "delta-12 desaturase" and "delta-15 desaturase" are used as a convenience to describe the proteins encoded by nucleic acid fragments

09697379.102600

that have been isolated based on the phenotypic effects caused by their disruption. They do not imply any catalytic mechanism. For example, delta-12 desaturase refers to the enzyme that catalyzes the formation of a double bond between carbons 12 and 13 of an 18 carbon fatty acid irrespective of whether it "counts" the carbons from the methyl, carboxyl end, or the first double bond. The term "fatty acid desaturase-related enzyme" refers to enzymes whose catalytic product may not be a carbon-carbon double bond but whose mechanism of action is similar to that of a fatty acid desaturase (that is, catalysis of the displacement of a carbon-hydrogen bond of a fatty acid chain to form a fatty-hydroxyacyl intermediate or end-product). Examples include delta-12 hydroxylase which means a delta-12 fatty acid hydroxylase or the oleate hydroxylase responsible for the synthesis of ricinoleic acid from oleic acid.

The term "nucleic acid" refers to a large molecule which can be single-stranded or double-stranded, composed of monomers (nucleotides) containing a sugar, a phosphate and either a purine or pyrimidine. A "nucleic acid fragment" is a fraction of a given nucleic acid molecule. In higher plants, deoxyribonucleic acid (DNA) is the genetic material while ribonucleic acid (RNA) is involved in the transfer of the information in DNA into proteins. A "genome" is the entire body of genetic material contained in each cell of an organism. The term "nucleotide sequence" refers to the sequence of DNA or RNA polymers, which can be single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases capable of incorporation into DNA or RNA polymers. The term "oligomer" refers to short nucleotide sequences, usually up to 100 bases long. As used herein, the term "homologous to" refers

09697379.10600

to the relatedness between the nucleotide sequence of two nucleic acid molecules or between the amino acid sequences of two protein molecules. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.); or by the comparison of sequence similarity between two nucleic acids or proteins, such as by the method of Needleman et al. (*J. Mol. Biol.* (1970) 48:443-453). As used herein, "substantially homologous" refers to nucleotide sequences that have more than 90% overall identity at the nucleotide level with the coding region of the claimed sequence, such as genes and pseudo-genes corresponding to the coding regions. The nucleic acid fragments described herein include molecules which comprise possible variations, both man-made and natural, such as but not limited to (a) those that involve base changes that do not cause a change in an encoded amino acid, or (b) which involve base changes that alter an amino acid but do not affect the functional properties of the protein encoded by the DNA sequence, (c) those derived from deletions, rearrangements, amplifications, random or controlled mutagenesis of the nucleic acid fragment, and (d) even occasional nucleotide sequencing errors.

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding) and following (3' non-coding) the coding region. "Fatty acid desaturase gene" refers to a nucleic acid fragment that expresses a protein with fatty acid desaturase activity. "Native" gene refers to an isolated gene with its own regulatory sequences as found in nature. "Chimeric gene" refers to a gene that comprises heterogeneous regulatory and

09697379.102600



coding sequences not found in nature. "Endogenous" gene refers to the native gene normally found in its natural location in the genome and is not isolated. A "foreign" gene refers to a gene not normally found in the host organism but that is introduced by gene transfer. "Pseudo-gene" refers to a genomic nucleotide sequence that does not encode a functional enzyme.

5 "Coding sequence" refers to a DNA sequence that codes for a specific protein and excludes the non-coding sequences. It may constitute an "uninterrupted coding sequence", i.e., lacking an intron or it may include one or more introns bounded by appropriate splice junctions. An "intron" is a nucleotide sequence that is transcribed in the primary transcript but that is removed through  
10 cleavage and re-ligation of the RNA within the cell to create the mature mRNA that can be translated into a protein.  
15

"Initiation codon" and "termination codon" refer to a unit of three adjacent nucleotides in a coding sequence that specifies initiation and chain termination, respectively, of protein synthesis (mRNA translation). "Open reading frame" refers to the coding sequence uninterrupted by introns between initiation and termination codons that encodes an amino acid sequence.  
20

25 "RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cdNA" refers to a double-stranded DNA that is  
30 complementary to and derived from mRNA. "Sense" RNA  
35

refers to RNA transcript that includes the mRNA. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene by interfering with the processing, transport and/or translation of its primary transcript or mRNA. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. In addition, as used herein, antisense RNA may contain regions of ribozyme sequences that increase the efficacy of antisense RNA to block gene expression. "Ribozyme" refers to a catalytic RNA and includes sequence-specific endoribonucleases.

- As used herein, "suitable regulatory sequences" refer to nucleotide sequences in native or chimeric genes that are located upstream (5'), within, and/or downstream (3') to the nucleic acid fragments of the invention, which control the expression of the nucleic acid fragments of the invention. The term "expression", as used herein, refers to the transcription and stable accumulation of the sense (mRNA) or the antisense RNA derived from the nucleic acid fragment(s) of the invention that, in conjunction with the protein apparatus of the cell, results in altered levels of the fatty acid desaturase(s). Expression or overexpression of the gene involves transcription of the gene and translation of the mRNA into precursor or mature fatty acid desaturase proteins. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of preventing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Cosuppression" refers to the expression of a foreign

gene which has substantial homology to an endogenous gene resulting in the suppression of expression of both the foreign and the endogenous gene. "Altered levels" refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

"Promoter" refers to a DNA sequence in a gene, usually upstream (5') to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. In artificial DNA constructs promoters can also be used to transcribe antisense RNA. Promoters may also contain DNA sequences that are involved in the binding of protein factors which control the effectiveness of transcription initiation in response to physiological or developmental conditions. It may also contain enhancer elements. An "enhancer" is a DNA sequence which can stimulate promoter activity. It may be an innate element of the promoter or a heterologous element inserted to enhance the level and/or tissue-specificity of a promoter. "Constitutive promoters" refers to those that direct gene expression in all tissues and at all times. "Tissue-specific" or "development-specific" promoters as referred to herein are those that direct gene expression almost exclusively in specific tissues, such as leaves or seeds, or at specific development stages in a tissue, such as in early or late embryogenesis, respectively.

The "3' non-coding sequences" refers to the DNA sequence portion of a gene that contains a polyadenylation signal and any other regulatory signal capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by

09697379.102600

affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

"Transformation" herein refers to the transfer of a foreign gene into the genome of a host organism and its genetically stable inheritance. "Restriction fragment length polymorphism" (RFLP) refers to different sized restriction fragment lengths due to altered nucleotide sequences in or around variant forms of genes. "Molecular breeding" refers to the use of DNA-based diagnostics, such as RFLP, RAPDs, and PCR in breeding. "Fertile" refers to plants that are able to propagate sexually.

"Plants" refer to photosynthetic organisms, both eukaryotic and prokaryotic, whereas the term "Higher plants" refers to eukaryotic plants. "Oil-producing species" herein refers to plant species which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (Glycine max), rapeseed and canola (including Brassica napus, B. campestris), sunflower (Helianthus annuus), cotton (Gossypium hirsutum), corn (Zea mays), cocoa (Theobroma cacao), safflower (Carthamus tinctorius), oil palm (Elaeis guineensis), coconut palm (Cocos nucifera), flax (Linum usitatissimum), castor (Ricinus communis) and peanut (Arachis hypogaea). The group also includes non-agronomic species which are useful in developing appropriate expression vectors such as tobacco, rapid cycling Brassica species, and Arabidopsis thaliana, and wild species which may be a source of unique fatty acids.

"Sequence-dependent protocols" refer to techniques that rely on a nucleotide sequence for their utility. Examples of sequence-dependent protocols include, but are not limited to, the methods of nucleic acid and oligomer hybridization and methods of DNA and RNA

amplification such as are exemplified in various uses of the polymerase chain reaction (PCR).

Various solutions used in the experimental manipulations are referred to by their common names such as "SSC", "SSPE", "Denhardt's solution", etc. The composition of these solutions may be found by reference to Appendix B of Sambrook, et al. (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press).

10        T-DNA Mutagenesis and Identification of an  
               Arabidopsis Mutant Defective in  
               Microsomal Delta-12 Desaturation

In T-DNA mutagenesis (Feldmann, et al., Science (1989) 243:1351-1354), the integration of T-DNA in the genome can interrupt normal expression of the gene at or near the site of the integration. If the resultant mutant phenotype can be detected and shown genetically to be tightly linked to the T-DNA insertion, then the "tagged" mutant locus and its wild type counterpart can be readily isolated by molecular cloning by one skilled in the art.

Arabidopsis thaliana seeds were transformed by Agrobacterium tumefaciens C58Clrif strain harboring the avirulent Ti-plasmid pGV3850::pAK1003 that has the T-DNA region between the left and right T-DNA borders replaced by the origin of replication region and ampicillin resistance gene of plasmid pBR322, a bacterial kanamycin resistance gene, and a plant kanamycin resistance gene (Feldmann, et al., Mol. Gen. Genetics (1987) 208:1-9). Plants from the treated seeds were self-fertilized and the resultant progeny seeds, germinated in the presence of kanamycin, were self-fertilized to give rise to a population, designated T3, that was segregating for T-DNA insertions. T3 seeds from approximately 1700 T2 plants were germinated and grown under controlled

09697379.102600

environment. One leaf from each of ten T3 plants of each line were pooled and analyzed for fatty acid composition. One line, designated 658, showed an increased level of oleic acid (18:1). Analysis of twelve individual T3 seeds of line 658 identified two seeds that contained greater than 36% oleic acid while the remaining seeds contained 12-22% oleic acid. The mutant phenotype of increased level of oleic acid in leaf and seed tissues of line 658 and its segregation in individual T3 seeds suggested that line 658 harbors a mutation that affects desaturation of oleic acid to linoleic acid in both leaf and seed tissues. When approximately 200 T3 seeds of line 658 were tested for their ability to germinate in the presence of kanamycin, four kanamycin-sensitive seeds were identified, suggesting multiple, possibly three, T-DNA inserts in the original T2 line. When progeny seeds of 100 individual T3 plants were analyzed for fatty acid composition and their ability to germinate on kanamycin, one plant, designated 658-75, was identified whose progeny segregated 7 wild type:2 mutant for the increased oleic acid and 28 sensitive:60 resistant for kanamycin resistance. Approximately 400 T4 progeny seeds of derivative line 658-75 were grown and their leaves analyzed for fatty acid composition. Ninety one of these seedlings were identified as homozygous for the mutant (high oleic acid) phenotype. Eighty-three of these homozygous plants were tested for the presence of nopaline, another marker for T-DNA, and all of them were nopaline positive. On the basis of these genetic studies it was concluded that the mutation in microsomal delta-12 desaturation was linked to the T-DNA.

006697379.10600

Isolation of Arabidopsis 658-75 Genomic DNA  
Containing the Disrupted Gene Controlling  
Microsomal Delta-12 Desaturation

In order to isolate the gene controlling microsomal  
5 delta-12 desaturation from wild-type Arabidopsis, a  
T-DNA-plant DNA "junction" fragment containing a T-DNA  
border integrated into the host plant DNA was isolated  
from the homozygous mutant plants of the 658-75 line of  
Arabidopsis. For this, genomic DNA from the mutant  
10 plant was isolated and completely digested by either Bam  
HI or Sal I restriction enzymes. In each case, one of  
the resultant fragments was expected to contain the  
origin of replication and ampicillin-resistance gene of  
pBR322 as well as the left T-DNA-plant DNA junction  
15 fragment. Such fragments were rescued as plasmids by  
ligating the digested genomic DNA fragments at a dilute  
concentration to facilitate self-ligation and then using  
the ligated fragments to transform E. coli cells. While  
no ampicillin-resistant colony was obtained from the  
20 plasmid rescue of Sal I-digested plant genomic DNA, a  
single ampicillin-resistant colony was obtained from the  
plasmid rescue of Bam HI-digested plant genomic DNA.  
The plasmid obtained from this transformant was  
designated p658-1. Restriction analysis of plasmid  
25 p658-1 with Bam HI, Sal I and Eco RI restriction enzymes  
strongly suggested that it contained the expected  
14.2 kb portion of the T-DNA (containing pBR322  
sequences) and a putative plant DNA/left T-DNA border  
fragment in a 1.6 kb Eco RI-Bam HI fragment. The 1.6 kb  
30 Eco RI-Bam HI fragment was subcloned into pBluescript SK  
[Stratagene] by standard cloning procedures described in  
Sambrook et al., (Molecular Cloning, A Laboratory  
Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory  
Press) and the resultant plasmid, designated pS1658.

09697379.102600

Isolation of Microsomal Delta-12 Desaturase cDNA  
and Gene from Wild type Arabidopsis

09697379-102600

The 1.6 kb Eco RI-Bam HI fragment, which contained the putative plant DNA flanking T-DNA, in plasmid p658-1 was isolated and used as a radiolabeled hybridization probe to screen a cDNA library made to polyA<sup>+</sup> mRNA from the above-ground parts of Arabidopsis thaliana plants, which varied in size from those that had just opened their primary leaves to plants which had bolted and were flowering [Elledge et al. (1991) Proc. Natl. Acad. Sci. USA 88:1731-1735]. The cDNA inserts in the library were made into an Xho I site flanked by Eco RI sites in lambda Yes vector [Elledge et al. (1991) Proc. Natl. Acad. Sci. USA 88:1731-1735]. Of the several positively-hybridizing plaques, four were subjected to plaque purification. Plasmids were excised from the purified phages by site-specific recombination using the cre-lox recombination system in E. coli strain BNN132 [Elledge et al. (1991) Proc. Natl. Acad. Sci. USA 88:1731-1735]. The four excised plasmids were digested by Eco RI restriction enzyme and shown to contain cDNA inserts ranging in size between 1 kb and 1.5 kb. Partial nucleotide sequence determination and restriction enzyme mapping of all four cDNAs revealed their common identity.

The partial nucleotide sequences of two cDNAs, designated pSF2b and p92103, containing inserts of ca. 1.2 kb and ca. 1.4 kb, respectively, were determined. The composite sequence derived from these plasmids is shown as SEQ ID NO:1 and is expected to be contained completely in plasmid p92103. SEQ ID NO:1 shows the 5' to 3' nucleotide sequence of 1372 base pairs of the Arabidopsis cDNA which encodes microsomal delta-12 fatty acid desaturase. Nucleotides 93-95 are the putative initiation codon of the open reading frame (nucleotides



93-1244), (identified by comparison of other plant delta-12 desaturases in this application). Nucleotides 1242-1244 are the termination codon. Nucleotides 1 to 92 and 1245-1372 are the 5' and 3' untranslated  
5 nucleotides, respectively. The 383 amino acid protein sequence in SEQ ID NO:2 is that deduced from the open reading frame and has an estimated molecular weight of 44 kD.

The gene corresponding to SEQ ID NO:1 was isolated  
10 by screening an *Arabidopsis* genomic DNA library using radiolabeled pSF2b cDNA insert, purifying the positively-hybridizing plaque, and subcloning a 6 kB Hind III insert fragment from the phage DNA in pBluescript vector. The sequence of 2973 nucleotides of  
15 the gene is shown in SEQ ID NO:15. Comparison of the sequences of the gene (SEQ ID NO:15) and the cDNA (SEQ ID NO:1) revealed the presence of a single intron of 1134 bp at a position between nucleotide positions 88 and 89 of the cDNA, which is 4 nucleotides 5' to the  
20 initiation codon.

The 1.6 kB Eco RI-Bam HI genomic border fragment insert in pS1658 was also partially sequenced from the Bam HI and Eco RI ends. Comparison of the nucleotide sequences of the gene (SEQ ID NO:15), the cDNA (SEQ ID  
25 NO:1), the border fragment, and the published sequence of the left end of T-DNA (Yadav et al., (1982) Proc. Natl. Acad. Sci. 79:6322-6326) revealed that a) the sequence of the first 451 nucleotides of the border fragment from the Bam HI end is collinear with that of  
30 nucleotides 539 (Bam HI site) to 89 of the cDNA, b) from the Eco RI end, the border fragment is collinear from nucleotides 1 to 61 with that of the left end of T-DNA (except for a deletion of 9 contiguous nucleotides at position 42 in the border fragment), and is collinear  
35 from nucleotides 57 to 104 with that of nucleotides

41-88 of the cDNA, and c) the sequence divergences between the border fragment and the cDNA are due to the presence of the intron in the border fragment. These results show that the T-DNA disrupted the microsomal  
 5 delta-12 desaturase gene in the transcribed region between the promoter and the coding region and 5' to the intron in the untranslated sequence.

A phage DNA containing Arabidopsis microsomal delta-12 desaturase gene was used as a RFLP marker on a  
 10 Southern blot containing genomic DNA from several progeny of Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with Hind III. This mapped the microsomal delta-12 desaturase gene 13.6 cM proximal to locus  
 15 c3838, 9.2 cM distal to locus 1At228, and 4.9 cM proximal to Fad D locus on chromosome 3 [Koorneef, M. et al., (1993) in Genetic Maps, Ed. O'Brien, S. J.; Yadav et al. (1993) Plant Physiology 103:467-476]. This position corresponds closely to previously suggested  
 20 locus for microsomal delta-12 desaturation (Fad 2) [Hugly, S. et al., (1991) Heredity 82:4321].

The open reading frames in SEQ ID NO:1 and in sequences encoding Arabidopsis microsomal delta-15 desaturase [WO 9311245], Arabidopsis plastid delta-15  
 25 desaturase [WO 9311245], and cyanobacterial desaturase, des A, [Wada, et al., Nature (1990) 347:200-203; Genbank ID:CSDESA; GenBank Accession No:X53508] as well as their deduced amino acid sequences were compared by the method of Needleman et al. [J. Mol. Biol. (1970) 48:443-453]  
 30 using gap weight and gap length weight values of 5.0 and 0.3, respectively, for the nucleotide sequences and 3.0 and 0.1, respectively, for protein sequences. The overall identities are summarized in Table 2.

TABLE 2

Percent Identity Between Different Fatty Acid  
Desaturases at the Nucleotide and Amino Acid Levels

		a3	ad	des A
a2	nucleotide	48(8 gaps)	46(6 gaps)	43(10 gaps)
	amino acid	39(9 gaps)	34(8 gaps)	24(10 gaps)
a3	nucleotide	-	65(1 gap)	43(9 gaps)
	amino acid	-	65(2 gaps)	26(11 gaps)
ad	nucleotide	-	-	43(9 gaps)
	amino acid	-	-	26(11 gaps)

- a2, a3, ad, and des A refer, respectively, to SEQ ID NO:1/2, Arabidopsis microsomal delta-15 desaturase, Arabidopsis plastid delta-15 desaturase, and cyanobacterial desaturase, des A. The percent identities in each comparison are shown at both the nucleotide and amino acid levels; the number of gaps imposed by the comparisons are shown in brackets following the percent identities. As expected on the basis of unsuccessful attempts in using delta-15 fatty acid nucleotide sequences as hybridization probes to isolate nucleotide sequences encoding microsomal delta-12 fatty acid desaturase, the overall homology at the nucleotide level between microsomal delta-12 fatty acid desaturase (SEQ ID NO:1) and the nucleotide sequences encoding the other three desaturases is poor (ranging between 43% and 48%). At the amino acid level too, the microsomal delta-12 fatty acid desaturase (SEQ ID NO:2) is poorly related to cyanobacterial des A (less than 24% identity) and the plant delta-15 desaturases (less than 39% identity).

While the overall relatedness between the deduced amino acid sequence of the said invention and the published fatty acid desaturases is limited, more significant identities are observed in shorter stretches

of amino acid sequences in the above comparisons. These results confirmed that the T-DNA in line 658-75 had interrupted the normal expression of a fatty acid desaturase gene. Based on the fatty acid phenotype of homozygous mutant line 658-75, Applicants concluded that SEQ ID NO:1 encoded the delta-12 desaturase. Further, Applicants concluded that it was the microsomal delta-12 desaturase, and not the chloroplastic delta-12 desaturase, since: a) the mutant phenotype was expressed strongly in the seed but expressed poorly, if at all, in the leaf of line 658-75, and b) the delta-12 desaturase polypeptide, by comparison to the microsomal and plastid delta-15 desaturase polypeptides [WO 9311245], did not have an N-terminal extension of a transit peptide expected for a nuclear-encoded plastid desaturase.

Plasmid p92103 was deposited on October 16, 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69095.

Expression Of Microsomal Delta-12 Fatty Acid Desaturase In Arabidopsis Fad2-1 Mutant To Complement Its Mutation In Delta-12 Fatty Acid Desaturation

To confirm the identity of SEQ ID NO:1 (Arabidopsis microsomal delta-12 fatty acid desaturase cDNA) a chimeric gene comprising of SEQ ID NO:1 was transformed into an Arabidopsis mutant affected in microsomal delta-12 fatty acid desaturation. For this, the ca. 1.4 kb Eco RI fragment containing the cDNA (SEQ ID NO:1) was isolated from plasmid p92103 and sub-cloned in pGA748 vector [An et. al.(1988) Binary Vectors. In: Plant Molecular Biology Manual. Eds Gelvin, S. B. et al. Kluwer Academic Press], which was previously linearized with Eco RI restriction enzyme. In one of the resultant binary plasmid, designated pGA-Fad2, the cDNA was placed

00697375-102600

in the sense orientation behind the CaMV 35S promotor of the vector to provide constitutive expression.

- Binary vector pGA-Fad2 was transformed by the freeze/thaw method [Holsters et al. (1978) Mol. Gen. Genet. 163:181-187] into Agrobacterium tumefaciens strain R1000, carrying the Ri plasmid pRiA4b from Agrobacterium rhizogenes [Moore et al., (1979) Plasmid 2:617-626] to result in transformants R1000/pGA-Fad2.
- Agrobacterium strains R1000 and R1000/pGA-Fad2 were used to transform Arabidopsis mutant fad2-1 [Miquel, M. & Browse, J. (1992) Journal of Biological Chemistry 267:1502-1509] and strain R1000 was used to transform wild type Arabidopsis. Young bolts of plants were sterilized and cut so that a single node was present in each explant. Explants were inoculated by Agrobacteria and incubated at 25°C in the dark on drug-free MS minimal organics medium with 30 g/L sucrose (Gibco). After four days, the explants were transferred to fresh MS medium containing 500 mg/L cefotaxime and 250 mg/ml carbenicillin for the counterselection of Agrobacterium. After 5 days, hairy roots derived from R1000/pGA-Fad2 transformation were excised and transferred to the same medium containing 50 mg/ml kanamycin. Fatty acid methyl esters were prepared from 5-10 mm of the roots essentially as described by Browse et al., (Anal. Biochem. (1986) 152:141-145) except that 2.5% H<sub>2</sub>SO<sub>4</sub> in methanol was used as the methylation reagent and samples were heated for 1.5 h at 80°C to effect the methanolysis of the seed triglycerides. The results are shown in Table 3. Root samples 41 to 46, 48 to 51, 58, and 59 are derived from transformation of fad2-1 plants with R1000/pGA Fad2; root samples 52, 53, and 57 were derived from transformation of fad2-1 plants with R1000 and serve as controls; root sample 60 is derived from

transformation of wild type Arabidopsis with R1000 and also serves as a control.

**TABLE 3**  
Fatty acid Composition in Transgenic  
Arabidopsis fad2-1 Hairy Roots Transformed  
with Agrobacterium R1000/pGA-fad2

Sample	16:0	16:1	18:0	18:1	18:2	18:3
41	24.4	1.8	1.7	5.0	29.4	33.8
42	25.6	3.7	1.3	20.0	22.0	27.5
43	23.6	-	1.6	7.2	27.6	36.1
44	24.4	1.3	4.6	16.0	18.1	33.6
45	20.7	-	8.1	44.7	11.8	14.8
46	20.1	-	1.8	7.5	33.7	36.0
48	26.1	2.9	2.1	9.5	17.6	33.4
49	30.8	1.0	2.4	8.7	18.7	31.1
50	19.8	1.9	3.3	27.7	21.8	24.4
51	20.9	1.1	5.0	13.7	25.0	32.1
58	23.5	0.3	1.4	3.6	22.1	45.9
59	22.6	0.6	1.4	2.8	29.9	40.4
52, cont.	12.3	-	2.6	64.2	4.6	16.4
53, cont.	20.3	9.1	2.2	55.2	1.7	9.2
57, cont.	10.4	2.4	0.7	65.9	3.8	12.7
60, WT	23.0	1.7	0.8	6.0	35.0	31.8

- 5            These results show that expression of Arabidopsis microsomal delta-12 desaturase in a mutant Arabidopsis lacking delta-12 desaturation can result in partial to complete complementation of the mutant. The decrease in oleic acid levels in transgenic roots is accompanied by
- 10 increases in the levels of both 18:2 and 18:3. Thus, overexpression of this gene in other oil crops, especially canola, which is a close relative of Arabidopsis and which naturally has high levels of 18:1 in seeds, is also expected to result in higher levels of 18:2, which in conjunction with

the overexpression of the microsomal delta-15 fatty acid desaturase will result in very high levels of 18:3.

Using Arabidopsis Microsomal Delta-12 Desaturase  
cDNA as a Hybridization Probe to Isolate Microsomal  
Delta-12 Desaturase cDNAs from Other Plant Species

Evidence for conservation of the delta-12 desaturase sequences amongst species was provided by using the Arabidopsis cDNA insert from pSF2b as a hybridization probe to clone related sequences from Brassica napus, and soybean. Furthermore, corn and castor bean microsomal delta-12 fatty acid desaturase were isolated by PCR using primers made to conserved regions of microsomal delta-12 desaturases.

Cloning of a Brassica napus Seed  
cDNA Encoding Seed Microsomal Delta-12  
Fatty Acid Desaturase

For the purpose of cloning the Brassica napus seed cDNA encoding a delta-12 fatty acid desaturase, the cDNA insert from pSF2b was isolated by digestion of pSF2b with EcoR I followed by purification of the 1.2 kb insert by gel electrophoresis. The 1.2 kb fragment was radiolabeled and used as a hybridization probe to screen a lambda phage cDNA library made with poly A<sup>+</sup> mRNA from developing Brassica napus seeds 20-21 days after pollination. Approximately 600,000 plaques were screened under low stringency hybridization conditions (50 mM Tris pH 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 ug denatured calf thymus DNA and 50°C) and washes (two washes with 2X SSC, 0.5% SDS at room temperature for 15 min each, then twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min each, and then twice with 0.2X SSC, 0.5% SDS at 50°C for 15 min each). Ten strongly-hybridizing phage were plaque-purified and the size of the cDNA inserts was determined by PCR amplification of the insert using phage as template and

T3/T7 oligomers for primers. Two of these phages, 165D and 165F, had PCR amplified inserts of 1.6 kb and 1.2 kb respectively and these phages were also used to excise the phagemids as described above. The phagemid derived from phage 165D, designated pCF2-165D, contained a 1.5 kb insert which was sequenced completely on one strand. SEQ ID NO:3 shows the 5' to 3' nucleotide sequence of 1394 base pairs of the Brassica napus cDNA which encodes delta-12 desaturase in plasmid pCF2-165D. Nucleotides 99 to 101 and nucleotides 1248 to 1250 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 99 to 1250). Nucleotides 1 to 98 and 1251 to 1394 are, respectively, the 5' and 3' untranslated nucleotides. The 383 amino acid protein sequence deduced from the open reading frame in SEQ ID NO:3 is shown in SEQ ID NO:4. While the other strand can easily be sequenced for confirmation, comparisons of SEQ ID NOS:1 and 3 and of SEQ ID NOS:2 and 4, even admitting of possible sequencing errors, showed an overall homology of approximately 84% at both the nucleotide and amino acid levels, which confirmed that pCF2-165D is a Brassica napus seed cDNA that encoded delta-12 desaturase. Plasmid pCF2-165D has been deposited on October 16, 1992 with the American Type Culture Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69094.

Cloning of a Soybean (Glycine max)  
cDNA Encoding Seed Microsomal Delta-12  
Fatty Acid Desaturase

A cDNA library was made to poly A<sup>+</sup> mRNA isolated from developing soybean seeds, and screened as described above.. Radiolabelled probe prepared from pSF2b as described above was added, and allowed to hybridize for 18 h at 50°C. The probes were washed as described

09697379.102600



above. Autoradiography of the filters indicated that there were 14 strongly hybridizing plaques, and numerous weakly hybridizing plaques. Six of the 14 strongly hybridizing plaques were plaque purified as described

5 above and the cDNA insert size was determined by PCR amplification of the insert using phage as template and T3/T7 oligomers for primers. One of these phages, 169K, had an insert sizes of 1.5 kb and this phage was also used to excise the phagemid as described above. The

10 phagemid derived from phage 169K, designated pSF2-169K, contained a 1.5 kb insert which was sequenced completely on both strands. SEQ ID NO:5 shows the 5' to 3' nucleotide sequence of 1473 base pairs of soybean (*Glycine max*) cDNA which encodes delta-12 desaturase in

15 plasmid pSF2-169K. Nucleotides 108 to 110 and nucleotides 1245 to 1247 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 108 to 1247). Nucleotides 1 to 107 and 1248 to 1462 are, respectively, the 5' and 3' untranslated nucleotides. The 380 amino acid protein sequence deduced from the open reading frame in SEQ ID NO:5 is shown in SEQ ID NO:6. Comparisons of SEQ ID NOS:1 and 5 and of SEQ ID NOS:2 and 6, even admitting of possible sequencing errors, showed an overall homology

20 of approximately 65% at the nucleotide level and approximately 70% at the amino acid level, which confirmed that pSF2-169K is a soybean seed cDNA that encoded delta-12 desaturase. A further description of this clone can be obtained by comparison of the SEQ ID

30 NO:1, SEQ ID NO:3, and SEQ ID NO:5 and by analyzing the phenotype of transgenic plants produced by using chimeric genes incorporating the insert of pSF2-169K, in sense or antisense orientation, with suitable regulatory sequences. Plasmid pSF2-169K was deposited on

35 October 16, 1992 with the American Type Culture

09697379.102600

Collection of Rockville, Maryland under the provisions of the Budapest Treaty and bears accession number ATCC 69092.

Cloning of a Corn (*Zea mays*)

5        cDNA Encoding Seed Microsomal Delta-12  
          Fatty Acid Desaturase

Corn microsomal delta-12 desaturase cDNA was isolated using a PCR approach. For this, a cDNA library was made to poly A<sup>+</sup> RNA from developing corn embryos in  
 10 Lambda Zap II vector. This library was used as template for PCR using sets of degenerate oligomers NS3 (SEQ ID NO:13) and RB5A/B (SEQ ID NOS:16 and 17) as sense and antisense primers, respectively. NS3 and RB5A/B correspond to stretches of amino acids 101-109 and  
 15 318-326, respectively, of SEQ ID NO:2, which are conserved in most microsomal delta-12 desaturases (for example, SEQ ID NOS:2, 4, 6, 8). PCR was carried out using a PCR kit (Perkin-Elmer) by 40 cycles of 94°C 1', 45°C, 1 min, and 55°C, 2 min. Analyses of the PCR  
 20 products on an agarose gel showed the presence of a product of the expected size (720 bp), which was absent in control reactions containing either the sense or antisense primers alone. The fragment was gel purified and then used as a probe for screening the corn cDNA  
 25 library at 60°C as described above. One positively-hybridizing plaque was purified and partial sequence determination of its cDNA showed it to be a nucleotide sequence encoding microsomal delta-12 desaturase but truncated at the 3' end. The cDNA insert encoding the  
 30 partial desaturase was gel isolated and used to probe the corn cDNA library again. Several positive plaques were recovered and characterized. DNA sequence analysis revealed that all of these clones seem to represent the same sequence with the different length of 5' or 3'  
 35 ends. The clone containing the longest insert,

06697379.102600

designated pFad2#1, was sequenced completely. The total length of the cDNA is 1790 bp (SEQ ID NO:7) comprising of an open reading frame from nucleotide 165 to 1328 bp that encoded a polypeptide of 388 amino acids. The deduced amino acid sequence of the polypeptide (SEQ ID NO:8) shared overall identities of 71%, 40%, and 38% to Arabidopsis microsomal delta-12 desaturase, Arabidopsis microsomal delta-15 desaturase, and Arabidopsis plastid delta-15 desaturase, respectively. Furthermore, it lacked an N-terminal amino acid extension that would indicate it is a plastid enzyme. Based on these considerations, it is concluded that it encodes a microsomal delta-12 desaturase.

#### Isolation of cDNAs Encoding

##### Delta-12 Microsomal Fatty Acid Desaturases and Desaturase-Related Enzymes from Castor Bean Seed

Polysomal mRNA was isolated from castor beans of stages I-II (5-10 DAP) and also from castor beans of stages IV-V (20-25 DAP). Ten ng of each mRNA was used for separate RT-PCR reactions, using the Perkin-Elmer RT-PCR kit. The reverse transcriptase reaction was primed with random hexamers and the PCR reaction with degenerate delta-12 desaturase primers NS3 and NS9 (SEQ ID NOS:13 and 14). The annealing-extension temperature of the PCR reaction was 50°C. A DNA fragment of approx. 700 bp was amplified from both stage I-II and stage IV-V mRNA. The amplified DNA fragment from one of the reactions was gel purified and cloned into a pGEM-T vector using the Promega pGEM-T PCR cloning kit to create the plasmid pRF2-1C. The 700 bp insert in pRF2-1C was sequenced, as described above, and the resulting DNA sequence is shown in SEQ ID NO:9. The DNA sequence in SEQ ID NO:9 contains an open-reading frame encoding 219 amino acids (SEQ ID NO:10) which has 81% identity (90% similarity) with amino acids 135 to 353 of

the *Arabidopsis* microsomal delta-12 desaturase described in SEQ ID NO:2. The cDNA insert in pRF2-1C is therefore a 676 bp fragment of a full-length cDNA encoding a castor bean seed microsomal delta-12 desaturase. The

5 full length castor bean seed microsomal delta-12 desaturase cDNA may isolated by screening a castor seed cDNA library, at 60°C, with the labeled insert of pRF2-1C as described in the example above. The insert in pRF2-1C may also be used to screen castor bean

10 libraries at lower temperatures to isolate delta-12 desaturase-related sequences, such as the delta-12 hydroxylase.

A cDNA library made to poly A<sup>+</sup> mRNA isolated from developing castor beans (stages IV-V, 20-25 DAP) was

15 screened as described above. Radiolabeled probe prepared from pSF2b or pRF2-1C, as described above, were added, and allowed to hybridize for 18 h at 50°C. The filters were washed as described above. Autoradiography of the filters indicated that there were numerous

20 hybridizing plaques, which appeared either strongly-hybridising or weakly-hybridising. Three of the strongly hybridising plaques (190A-41, 190A-42 and 190A-44) and three of the weakly hybridising plaques, (190B-41, 190b-43 and 197c-42), were plaque purified

25 using the methods described above. The cDNA insert size of the purified phages were determined by PCR amplification of the insert using phage as template and lambda-gt11 oligomers (Clontech lambda-gt11 Amplimers) for primers. The PCR-amplified inserts of the amplified

30 phages were subcloned into pBluescript (Pharmacia) which had been cut with Eco RI and filled in with Klenow (Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory Press). The resulting plasmids were called pRF190a-41,

35 pRF190a-42, pRF190a-44, pRF190b-41, pRF190b-43 and

09697379.102600

09697379-102600

pRF197c-42. All of the inserts were about 1.1 kb with the exception of pRF197c-42 which was approx. 1.5 kb. The inserts in the plasmids were sequenced as described above. The insert in pRF190b-43 did not contain any open reading frame and was not identified. The inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 were identical. The insert in pRF197c-42 contained all of the nucleotides of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 plus an additional approx. 400 bp. It was deduced therefore that the insert in pRF197c-42 was a longer version of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 and all were derived from the same full-length mRNA. The complete cDNA sequence of the insert in plasmid pRF197c-42 is shown in SEQ ID NO:11. The deduced amino acid sequence of SEQ ID NO:11, shown in SEQ ID NO:12, is 78.5% identical (90% similarity) to the castor microsomal delta-12 desaturase described above (SEQ ID NO:10) and 66% identical (80% similarity) to the Arabidopsis delta-12 desaturase amino acid sequence in SEQ ID NO:2. These similarities confirm that pRF197c-42 is a castor bean seed cDNA that encodes a microsomal delta-12 desaturase or a microsomal delta-12 desaturase-related enzyme, such as a delta-12 hydroxylase. Specific PCR primers for pRF2-1C and pRF197c-42 were made. For pRF2-1C the upstream primer was bases 180 to 197 of the cDNA sequence in SEQ ID NO:9. For pRF197c-42 the upstream primer was bases 717 to 743 of the cDNA sequence in SEQ ID NO:11. A common downstream primer was made corresponding to the exact complement of the nucleotides 463 to 478 of the sequence described in SEQ ID NO:9. Using RT-PCR with random hexamers and the above primers it was observed that the cDNA contained in pRF2-1C is expressed in both Stage I-II and Stage IV-V castor bean seeds whereas the cDNA contained in plasmid

pRF197c-42 is expressed only in Stage IV-V castor bean seeds, i.e., it is only expressed in tissue actively synthesizing ricinoleic acid. Thus, it is possible that this cDNA encodes a delta-12 hydroxylase.

- 5        There is enough deduced amino acid sequence from the two castor sequences described in SEQ ID NOS:10 and 12 to compare the highly conserved region corresponding to amino acids 311 to 353 of SEQ ID NO:2. When SEQ ID NOS:2, 4, 6, 8, and 10 are aligned by the Hein method
- 10       described above the consensus sequence corresponds exactly to the amino acids 311 to 353 of SEQ ID NO:2. All of the seed microsomal delta-12 desaturases described above have a high degree of identity with the consensus over this region, namely Arabidopsis (100%
- 15       identity), soybean (90% identity), corn (95% identity), canola (93% identity) and one (pRF2-1c) of the castor bean sequences (100% identity). The other castor bean seed delta-12 desaturase or desaturase-related sequence (pRF197c-42) however has less identity with the
- 20       consensus, namely 81% for the deduced amino acid sequence of the insert in pRF197c-42 (described in SEQ ID NO:12). Thus while it remains possible that the insert in pRF197c-42 encodes a microsomal delta-12 desaturase, this observation supports the hypothesis
- 25       that it encodes a desaturase-related sequence, namely the delta-12 hydroxylase.

- An additional approach to cloning a castor bean seed delta-12 hydroxylase is the screening of a differential population of cDNAs. A lambda-Zap
- 30       (Stratagene) cDNA library made to polysomal mRNA isolated from developing castor bean endosperm (stages IV-V, 20-25 DAP) was screened with <sup>32</sup>P-labeled cDNA made from polysomal mRNA isolated from developing castor bean endosperm (stage I-II, 5-10 DAP) and with <sup>32</sup>P-labeled
- 35       cDNA made from polysomal mRNA isolated from developing

castor bean endosperm (stages IV-V, 20-25 DAP). The library was screened at a density of 2000 plaques per 137 mm plate so that individual plaques were isolated. About 60,000 plaques were screened and plaques which  
5 hybridised with late (stage IV/V) cDNA but not early (stage I/II) cDNA, which corresponded to about 1 in every 200 plaques, were pooled.

The library of differentially expressed cDNAs may be screened with the castor delta-12 desaturase cDNA  
10 described above and/or with degenerate oligonucleotides based on sequences of amino conserved among delta-12 desaturases to isolate related castor cDNAs, including the cDNA encoding the delta-12 oleate hydroxylase enzyme. These regions of amino acid conservation may  
15 include, but are not limited to amino acids 101 to 109, 137 to 145, and 318 to 327 of the amino acid sequence described in SEQ ID NO:2 or any of the sequences described in Table 7 below. Examples of such oligomers are SEQ ID NOS:13, 14, 16, and 17. The insert in  
20 plasmid pCF2-197c may be cut with Eco RI to remove vector sequences, purified by gel electrophoresis and labeled as described above. Degenerate oligomers based on the above conserved amino acid sequences may be labeled with <sup>32</sup>P as described above. The cDNAs cloned  
25 from the developing endosperm difference library which do not hybridize with early mRNA probe but do hybridize with late mRNA probe and hybridize with either castor delta-12 desaturase cDNA or with an oligomer based on delta-12 desaturase sequences are likely to be the  
30 castor delta-12 hydroxylase. The pBluescript vector containing the putative hydroxylase cDNA can be excised and the inserts directly sequenced, as described above.

Clones such as pRF2-1C and pRF197c-42, and other clones from the differential screening, which, based on  
35 their DNA sequence, are less related to castor bean seed

09697379.102600

microsomal delta-12 desaturases and are not any of the known fatty-acid desaturases described above or in WO 9311245, may be expressed, for example, in soybean embryos or another suitable plant tissue, or in a microorganism, such as yeast, which does not normally contain ricinoleic acid, using suitable expression vectors and transformation protocols. The presence of novel ricinoleic acid in the transformed tissue(s) expressing the castor cDNA would confirm the identity of the castor cDNA as DNA encoding for an oleate hydroxylase.

#### Sequence Comparisons Among Seed Microsomal Delta-12 Desaturases

The percent overall identities between coding regions of the full-length nucleotide sequences encoding microsomal delta-12 desaturases was determined by their alignment by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 5.0 and 0.3 (Table 4). Here, a2, c2, s2, z2 and des A refer, respectively, to the nucleotide sequences encoding microsomal delta-12 fatty acid desaturases from *Arabidopsis* (SEQ ID NO:1), *Brassica napus* (SEQ ID NO:3), soybean (SEQ ID NO:5), corn (SEQ ID NO:7), and cyanobacterial des A, whereas r2 refers to the microsomal delta-12 desaturase or desaturase-related enzyme from castor bean (SEQ ID NO:12).

TABLE 4

Percent Identity Between the Coding Regions of Nucleotide Sequences Encoding Different Microsomal Delta-12 Fatty Acid Desaturases

	<u>c2</u>	<u>s2</u>	<u>z2</u>	<u>des A</u>
a2	84	66	64	43
c2	-	65	62	42
s2	-	-	62	42



The overall relatedness between the deduced amino acid sequences of microsomal delta-12 desaturases or desaturase-related enzymes of the invention (i.e., SEQ ID NOS:2, 4, 6, 8, and 12) determined by their alignment by the method of Needleman et al. (J. Mol. Biol. (1970) 48:443-453) using gap weight and gap length weight values of 3.0 and 0.1, respectively, is shown in Table 5. Here a2, c2, s2, z2 and des A refer, respectively, to microsomal delta-12 fatty acid desaturases from Arabidopsis (SEQ ID NO:2), Brassica napus (SEQ ID NO:4), soybean (SEQ ID NO:6), corn (SEQ ID NO:8), and cyanobacterial des A, whereas r2 refers to the microsomal desaturase or desaturase-related enzyme from castor bean (SEQ ID NO:12). The relatedness between the enzymes is shown as percent overall identity/percent overall similarity.

TABLE 5  
Relatedness Between Different Microsomal  
Delta-12 Fatty Acid Desaturases

	<u>c2</u>	<u>s2</u>	<u>r2</u>	<u>z2</u>	<u>des A</u>
a2	84/89	70/85	66/80	71/83	24/50
c2	-	67/80	63/76	69/79	24/51
s2	-	-	67/83	66/82	23/49
r2	-	-	-	61/78	24/51
z2	-	-	-	-	25/49

The high degree of overall identity (60% or greater) at the amino acid levels between the Brassica napus, soybean, castor and corn enzymes with that of Arabidopsis microsomal delta-12 desaturase and their lack of an N-terminal extension of a transit peptide expected for a nuclear-encoded chloroplast desaturase leads Applicants to conclude that SEQ ID NOS:4, 6, 8, 10, and 12 encode the microsomal delta-12

desaturases or desaturase-related enzymes. Further confirmation of this identity will come from biological function, that is, by analyzing the phenotype of transgenic plants or other organisms produced by using

5 chimeric genes incorporating the above-mentioned sequences in sense or antisense orientation, with-suitable regulatory sequences. Thus, one can isolate cDNAs and genes for homologous fatty acid desaturases from the same or different higher plant species,

10 especially from the oil-producing species. Furthermore, based on these comparisons, the Applicants expect all higher plant microsomal delta-12 desaturases from all higher plants to show an overall identity of 60% or more and to be able to readily isolate homologous fatty acid

15 desaturase sequences using SEQ ID NOS:1, 3, 5, 7, 9, and 11 by sequence-dependent protocols.

The overall percent identity at the amino acid level, using the above alignment method, between selected plant desaturases is illustrated in Table 6.

**TABLE 6**  
Percent Identity Between Selected Plant Fatty Acid  
Desaturases at the Amino Acid Level

	<u>a3</u>	<u>aD</u>	<u>c3</u>	<u>cD</u>	<u>S3</u>
a2	38	33	38	35	34
a3	-	65	93	66	67
aD	-	-	66	87	65
c3	-	-	-	67	67
cD	-	-	-	-	65

In Table 6, a2, a3, aD, c3, cD, and S3 refer, respectively, to SEQ ID NO:2, Arabidopsis microsomal delta-15 desaturase, Arabidopsis plastid delta-15

20 desaturase, canola microsomal delta-15 desaturase, canola plastid delta-15 desaturase, and soybean microsomal delta-15 desaturase. Based on these

comparisons, the delta-15 desaturases, of both microsomal and plastid types, have overall identities of 65% or more at the amino acid level, even when from the same plant species. Based on the above the Applicants expect microsomal delta-12 desaturases from all higher plants to show similar levels of identity (that is, 60% or more identity at the amino acid level) and that SEQ ID NOS:1, 3, 5, 7, and 9 can also be used as hybridization probe to isolate homologous delta-12 desaturase sequences, and possibly for sequences for fatty acid desaturase-related enzymes, such as oleate hydroxylase, that have an overall amino acid homology of 50% or more.

Similar alignments of protein sequences of plant microsomal fatty acid delta-12 desaturases [SEQ ID NOS:2, 4, 6, and 8] and plant delta-15 desaturases [microsomal and plastid delta-15 desaturases from *Arabidopsis* and *Brassica napus*, WO 9311245] allows identification of amino acid sequences conserved between the different desaturases (Table 7).

**TABLE 7**  
Amino Acid Sequences Conserved Between  
Plant Microsomal Delta-12 Desaturases and Microsomal and  
Plastid Delta-15 Desaturases

Region	Conserved AA Positions in SEQ ID NO:2	Consensus Conserved AA Sequence in $\Delta^{12}$ Desaturases	Consensus Conserved AA Sequence in $\Delta^{15}$ Desaturases	Consensus AA Sequence
A	39-44	<u>AIPPHC</u>	<u>AIPKHC</u>	AIP(P/K)HC
B	86-90	<u>WPL(I)YW</u>	<u>WPLYW</u>	WP(L/I)YW
C	104-109	<u>AHECGH</u>	<u>GHDCGH</u>	(A/G)H(D/E)CGH
D	130-134	<u>LLVPY</u>	<u>ILVPY</u>	(L/I)LVPY
E	137-142	<u>WKYSHR</u>	<u>WRISHR</u>	W(K/R)(Y/I)SHR
F	140-145	<u>SHRRHH</u>	<u>SHRTHH</u>	SHR(R/T)HH
G	269-274	<u>ITYLQ</u>	<u>VTYLH</u>	(I/V)TYL(Q/H)
H	279-282	<u>LPHY</u>	<u>LPWY</u>	LP(H/W)Y

I	289-294	<u>WL(R/K)GAL</u>	<u>YL</u> RGGL	(W/Y)L(R/K)G(A/G)L
J	296-302	<u>TVDRDYG</u>	<u>TLDRDYG</u>	T(V/L)DRDYG
K	314-321	<u>THVAHHLF</u>	<u>THVIHHLF</u>	THV(A/I)HHLF
L	318-327	<u>HHLESTMPHY</u>	<u>HHLEFQIPHY</u>	HHFL(S/P) (T/Q)(I/M)PHY

Table 7 shows twelve regions of conserved amino acid sequences, designated A-L (column 1), whose positions in SEQ ID NO:2 are shown in column 2. The consensus sequences for these regions in plant delta-12 fatty acid desaturases and plant delta-15 fatty acid desaturases are shown in columns 3 and 4, respectively; amino acids are shown by standard abbreviations, the underlined amino acids are conserved between the delta-12 and the delta-15 desaturases, and amino acids in brackets represent substitutions found at that position. The consensus sequence of these regions are shown in column 5. These short conserved amino acids and their relative positions further confirm that the isolated isolated cDNAs encode a fatty acid desaturase.

Isolation of Nucleotide Sequences Encoding  
Homologous and Heterologous Fatty acid Desaturases  
and Desaturase-like Enzymes

Fragments of the instant invention may be used to isolate cDNAs and genes of homologous and heterologous fatty acid desaturases from the same species as the fragments of the invention or from different species. Isolation of homologous genes using sequence-dependent protocols is well-known in the art and Applicants have demonstrated that Arabidopsis microsomal delta-12 desaturase cDNA sequence can be used to isolate cDNA for related fatty acid desaturases from Brassica napus, soybean, corn and castor bean.

More importantly, one can use the fragments containing SEQ ID NOS:1, 3, 5, 7, and 9 or their

00697379-102600

smaller, more conserved regions to isolate novel fatty acid desaturases and fatty acid desaturase-related enzymes.

In a particular embodiment of the present invention, regions of the nucleic acid fragments of the invention that are conserved between different desaturases may be used by one skilled in the art to design a mixture of degenerate oligomers for use in sequence-dependent protocols aimed at isolating nucleic acid fragments encoding homologous or heterologous fatty acid desaturase cDNA's or genes. For example, in the polymerase chain reaction (Innis, et al., Eds, (1990) PCR Protocols: A Guide to Methods and Applications, Academic Press, San Diego), two short pieces of the present fragment of the invention can be used to amplify a longer fatty acid desaturase DNA fragment from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleotide sequences with one primer based on the fragment of the invention and the other on either the poly A<sup>+</sup> tail or a vector sequence. These oligomers may be unique sequences or degenerate sequences derived from the nucleic acid fragments of the invention. The longer piece of homologous fatty acid desaturase DNA generated by this method could then be used as a probe for isolating related fatty acid desaturase genes or cDNAs from *Arabidopsis* or other species and subsequently identified by differential screening with known desaturase sequences and by nucleotide sequence determination. The design of oligomers, including long oligomers using deoxyinosine, and "guessmers" for hybridization or for the polymerase chain reaction are known to one skilled in the art and discussed in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). Short stretches of

009697379.102600

amino acid sequences that are conserved between cyanobacterial delta-12 desaturase (Wada et al., Nature (1990) 347:200-203) and plant delta-15 desaturases [WO 9311245] were previously used to make oligo-nucleotides that were degenerate and/or used deoxyinosine/s. One set of these oligomers made to a stretch of 12 amino acids conserved between cyanobacterial delta-12 desaturase and higher plant delta-15 desaturases was successful in cloning the plastid delta-12 desaturase cDNAs; these plant desaturases have more than 50% identity to the cyanobacterial delta-12 desaturase. Some of these oligonucleotides were also used as primers to make polymerase chain reaction (PCR) products using poly A<sup>+</sup> RNA from plants. However, none of the oligonucleotides and the PCR products were successful as radiolabeled hybridization probes in isolating nucleotide sequences encoding microsomal delta-12 fatty acid desaturases. Thus, as expected, none of the stretches of four or more amino acids conserved between Arabidopsis delta-12 and Arabidopsis delta-15 desaturases are identical in the cyanobacterial desaturase, just like none of the stretches of four or more amino acids conserved between Arabidopsis delta-15 and the cyanobacterial desaturase are identical in SEQ ID NO:2. Stretches of conserved amino acids between the present invention and delta-15 desaturases now allow for the design of oligomers to be used to isolate sequences encoding other desaturases and desaturase-related enzymes. For example, conserved stretches of amino acids between delta-12 desaturase and delta-15 desaturase, shown in Table 7, are useful in designing long oligomers for hybridization as well as shorter ones for use as primers in the polymerase chain reaction. In this regard, sequences conserved between delta-12 and delta-15 desaturases (shown in Table 7)

09697379.102600

evidence that the high oleic acid trait in mutant 658 is the result of T-DNA insertion.

#### Plasmid Rescue and Analysis

One-half and one microgram of genomic DNA from the  
5 homozygous mutant plants of the 658-75 line, prepared from leaf tissue as described [Rogers, S. O. and A. J. Bendich (1985) Plant Molecular Biology 5:69-76], was digested with 20 units of either Bam HI or Sal I restriction enzyme (Bethesda Research Laboratory) in a  
10 50  $\mu$ L reaction volume according to the manufacturer's specifications. After digestion the DNA was extracted with buffer-saturated phenol (Bethesda Research Laboratory) followed by precipitation in ethanol. One-half to one microgram of Bam HI or Sal I digested  
15 genomic DNA was resuspended in 200  $\mu$ L or 400  $\mu$ L of ligation buffer containing 50 mM Tris-Cl, pH 8.0, 10 mM  $MgCl_2$ , 10 mM dithiothreitol, 1 mM ATP, and 4 units of T4 DNA ligase (Bethesda Research Laboratory). The dilute DNA concentration of approximate 2.5  $\mu$ g/mL in the  
20 ligation reaction was chosen to facilitate circularization, as opposed to intermolecular joining. The reaction was incubated for 16 h at 16°C. Competent DH10B cells (Bethesda Research Laboratory) were transfected with 10 ng of ligated DNA per 100  $\mu$ L of  
25 competent cells according to the manufacturer's specifications. Transformants from Sal I or Bam HI digests were selected on LB plates (10 g Bacto-tryptone, 5 g Bacto-yeast extract, 5 g NaCl, 15 g agar per liter, pH 7.4) containing 100  $\mu$ g/mL ampicillin. After  
30 overnight incubation at 37°C the plates were scored for ampicillin-resistant colonies.

A single ampicillin-resistant transformant derived from Bam HI-digested plant DNA was used to start a culture in 35 mL LB medium (10 g Bacto-tryptone, 5 g  
35 yeast-extract, 5 g NaCl per liter) containing 25 mg/L

09697379.102600

ampicillin. The culture was incubated with shaking overnight at 37°C and the cells were then collected by centrifugation at 1000xg for 10 min. Plasmid DNA, designated p658-1, was isolated from the cells by the alkaline lysis method of Birnboim et al. [Nucleic Acid Research (1979) 7:1513-1523], as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed (1989) Cold Spring Harbor Laboratory Press). Plasmid p658-1 DNA was digested by restriction enzymes Bam HI, Eco RI and Sal I (Bethesda Research Laboratory) and electrophoresed through a 1% agarose gel in 1xTBE buffer (0.089M tris-borate, 0.002M EDTA). The restriction pattern indicated the presence in this plasmid of the expected 14.2 kB T-DNA fragment and a 1.6 kB putative plant DNA/T-DNA border fragment.

#### EXAMPLE 2

#### CLONING OF ARABIDOPSIS THALIANA MICROSOMAL DELTA-12 DESATURASE cDNA USING GENOMIC DNA FLANKING THE T-DNA SITE OF INSERTION IN ARABIDOPSIS THALIANA' MUTANT LINE 658-75 AS A HYBRIDIZATION PROBE

Two hundred nanograms of the 1.6 kB Eco RI-Bam HI fragment from plasmid p658-1, following digestion of the plasmid with Eco RI and Bam HI and purification by electrophoresis in agarose, was radiolabelled with alpha[32P]-dCTP using a Random Priming Labeling Kit (Bethesda Research Laboratory) under conditions recommended by the manufacturer.

The radiolabeled DNA was used as a probe to screen an Arabidopsis cDNA library made from RNA isolated from above ground portions of various growth stages (Elledge et al., (1991) Proc. Nat. Acad. Sci., 88:1731-1735) essentially as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor Laboratory Press). For this, approximately 17,000 plaque-forming units were plated on

09697379.102600



- seven 90mm petri plates containing a lawn of LE392 E. coli cells on NZY agar media (5 g NaCl, 2 g MgSO<sub>4</sub>-7 H<sub>2</sub>O, 5 g yeast extract, 10 g casein acid hydrolysate, 13 g agar per liter). Replica filters of the phage
- 5 plaques were prepared by adsorbing the plaques onto nitrocellulose filters (BA85, Schleicher and Schuell) then soaking successively for five min each in 0.5 M NaOH/1 M NaCl, 0.5 M Tris(pH 7.4)/1.5 M NaCl and 2xSSPE (0.36 M NaCl, 20 mM NaH<sub>2</sub>PO<sub>4</sub>(pH 7.4), 20 mM EDTA
- 10 (pH 7.4)). The filters were then air dried and baked for 2 h at 80°C. After baking the filters were wetted in 2X SSPE, and then incubated at 42°C in prehybridization buffer (50% Formamide, 5X SSPE, 1% SDS, 5X Denhardt's Reagent, and 100 ug/mL denatured salmon
- 15 sperm DNA) for 2 h. The filters were removed from the prehybridization buffer, and then transferred to hybridization buffer (50% Formamide, 5X SSPE, 1% SDS, 1X Denhardt's Reagent, and 100 ug/mL denatured salmon sperm DNA) containing the denatured radiolabeled probe (see
- 20 above) and incubated for 40 h at 42°C. The filters were washed three times in 2X SSPE/0.2% SDS at 42°C (15 min each) and twice in 0.2X SSPE/0.2% SDS at 55°C (30 min each), followed by autoradiography on Kodak XAR-5 film with an intensifying screen at -80°C, overnight.
- 25 Fifteen plaques were identified as positively-hybridizing on replica filters. Five of these were subjected to plaque purification essentially as described in Sambrook et al., (Molecular Cloning, A Laboratory Manual, 2nd ed. (1989), Cold Spring Harbor
- 30 Laboratory Press). The lambda YES-R cDNA clones were converted to plasmid by propagating the phage in the E. coli BNN-132 cells, which expresses Cre protein that excises the cDNA insert as a double-stranded plasmid by cre-mediated in vivo site-specific recombination at a
- 35 'lox' sites present in the phage. Ampicillin-resistant

09697379-102600

plasmid clones containing cDNA inserts were grown in liquid culture, and plasmid DNA was prepared using the alkaline lysis method as previously described. The sizes of the resulting plasmids were analyzed by

5 electrophoresis in agarose gels. The agarose gels were treated with 0.5 M NaOH/1 M NaCl, and 0.5 M Tris(pH 7.4), 1.5 M NaCl for 15 min each, and the gel was then dried completely on a gel drier at 65°C. The gel was hydrated in 2X SSPE and incubated overnight, at

10 42°C, in hybridization buffer containing the denatured radiolabeled probe, followed by washing as described above. After autoradiography, the inserts of four of the purified cDNA clones were found to have hybridized to the probe. Plasmid DNA from the hybridizing clones

15 was purified by equilibration in a CsCl/ethidium bromide gradient (see above). The four cDNA clones were sequenced using Sequenase T7 DNA polymerase (US Biochemical Corp.) and the manufacturer's instructions, beginning with primers homologous to vector sequences

20 that flank the cDNA insert. After comparing the partial sequences of the inserts obtained from the four clones, it was apparent that they each contained sequences in common. One cDNA clone, p92103, containing ca. 1.4 kB cDNA insert, was sequenced. The longest three clones

25 were subcloned into the plasmid vector pBluescript (Stratagene). One of these clones, designated pSF2b, containing ca 1.2 kB cDNA insert was also sequenced serially with primers designed from the newly acquired sequences as the sequencing experiment progressed. The

30 composite sequence derived from pSF2b and p92103 is shown in SEQ ID NO:1.

09697379.102600

EXAMPLE 3CLONING OF PLANT FATTY ACIDDESATURASE cDNAs USING THE ARABIDOPSIS THALIANA  
MICROSOMAL DELTA-12 DESATURASE cDNA CLONE AS A  
HYBRIDIZATION PROBE

5 An approximately 1.2 kb fragment containing the  
Arabidopsis delta-12 desaturase coding sequence of SEQ  
 ID NO:1 was obtained from plasmid pSF2b. This plasmid  
 was digested with EcoR I and the 1.2 kb delta-12  
 10 desaturase cDNA fragment was purified from the vector  
 sequence by agarose gel electrophoresis. The fragment  
 was radiolabelled with  $^{32}\text{P}$  as previously described.

Cloning of a Brassica napus Seed  
cDNA Encoding Microsomal Delta-12 Fatty AcidDesaturase

15 The radiolabelled probe was used to screen a  
Brassica napus seed cDNA library. In order to construct  
 the library, Brassica napus seeds were harvested 20-21  
 days after pollination, placed in liquid nitrogen, and  
 20 polysomal RNA was isolated following the procedure of  
 Kamalay et al., (Cell (1980) 19:935-946). The  
 polyadenylated mRNA fraction was obtained by affinity  
 chromatography on oligo-dT cellulose (Aviv et al., Proc.  
 Natl. Acad. Sci. USA (1972) 69:1408-1411). Four  
 25 micrograms of this mRNA were used to construct a seed  
 cDNA library in lambda phage (Uni-ZAP™ XR vector) using  
 the protocol described in the ZAP-cDNA™ Synthesis Kit  
 (1991 Stratagene Catalog, Item #200400). Approximately  
 600,000 clones were screened for positively hybridizing  
 30 plaques using the radiolabelled EcoR I fragment from  
 pSF2b as a probe essentially as described in Sambrook et  
 al., (Molecular Cloning: A Laboratory Manual, 2nd ed.  
 (1989) Cold Spring Harbor Laboratory Press) except that  
 low stringency hybridization conditions (50 mM Tris, pH  
 35 7.6, 6X SSC, 5X Denhardt's, 0.5% SDS, 100 µg denatured

009697379.102600

calf thymus DNA and 50°C) were used and post-hybridization washes were performed twice with 2X SSC, 0.5% SDS at room temperature for 15 min, then twice with 0.2X SSC, 0.5% SDS at room temperature for 15 min, and then twice with 0.2X SSC, 0.5% SDS at 50°C for 15 min. Ten positive plaques showing strong hybridization were picked, plated out, and the screening procedure was repeated. From the secondary screen nine pure phage plaques were isolated. Plasmid clones containing the cDNA inserts were obtained through the use of a helper phage according to the *in vivo* excision protocol provided by Stratagene. Double-stranded DNA was prepared using the alkaline lysis method as previously described, and the resulting plasmids were size-analyzed by electrophoresis in agarose gels. The largest one of the nine clones, designated pCF2-165D, contained an approximately 1.5 kb insert which was sequenced as described above. The sequence of 1394 bases of the cDNA insert of pCF2-165D is shown in SEQ ID NO:3. Contained in the insert but not shown in SEQ ID NO:3 are approximately 40 bases of the extreme 5' end of the 5' non-translated region and a poly A tail of about 38 bases at the extreme 3' end of the insert.

#### Cloning of a Soybean Seed

#### cDNA Encoding Microsomal Delta-12 Fatty Acid Desaturase

A cDNA library was made as follows: Soybean embryos (ca. 50 mg fresh weight each) were removed from the pods and frozen in liquid nitrogen. The frozen embryos were ground to a fine powder in the presence of liquid nitrogen and then extracted by Polytron homogenization and fractionated to enrich for total RNA by the method of Chirgwin et al. (Biochemistry (1979) 18:5294-5299). The nucleic acid fraction was enriched for poly A<sup>+</sup>RNA by passing total RNA through an oligo-dT

00697379.102600

cellulose column and eluting the poly A<sup>+</sup>RNA with salt as described by Goodman et al. (Meth. Enzymol. (1979) 68:75-90). cDNA was synthesized from the purified poly A<sup>+</sup>RNA using cDNA Synthesis System (Bethesda Research Laboratory) and the manufacturer's instructions. The resultant double-stranded DNA was methylated by Eco RI DNA methylase (Promega) prior to filling-in its ends with T4 DNA polymerase (Bethesda Research Laboratory) and blunt-end ligation to phosphorylated Eco RI linkers using T4 DNA ligase (Pharmacia). The double-stranded DNA was digested with Eco RI enzyme, separated from excess linkers by passage through a gel filtration column (Sephadex CL-4B), and ligated to lambda ZAP vector (Stratagene) according to manufacturer's instructions. Ligated DNA was packaged into phage using the Gigapack packaging extract (Stratagene) according to manufacturer's instructions. The resultant cDNA library was amplified as per Stratagene's instructions and stored at -80°C.

Following the instructions in the Lambda ZAP Cloning Kit Manual (Stratagene), the cDNA phage library was used to infect *E. coli* BB4 cells and approximately 600,000 plaque forming units were plated onto 150 mm diameter petri plates. Duplicate lifts of the plates were made onto nitrocellulose filters (Schleicher & Schuell). The filters were prehybridized in 25 mL of hybridization buffer consisting of 6X SSPE, 5X Denhardt's solution, 0.5% SDS, 5% dextran sulfate and 0.1 mg/mL denatured salmon sperm DNA (Sigma Chemical Co.) at 50°C for 2 h. Radiolabelled probe prepared from pSF2b as described above was added, and allowed to hybridize for 18 h at 50°C. The filters were washed exactly as described above. Autoradiography of the filters indicated that there were 14 strongly hybridizing plaques. The 14 plaques were subjected to a

09697379.102600

second round of screening as before. Numerous, strongly hybridizing plaques were observed on 6 of the 14 filters, and one, well-isolated from other phage, was picked from each of the six plates for further analysis.

5       Following the Lambda ZAP Cloning Kit Instruction Manual (Stratagene), sequences of the pBluescript vector, including the cDNA inserts, from the purified phages were excised in the presence of a helper phage and the resultant phagemids were used to infect *E. coli*  
 10   XL-1 Blue cells. DNA from the plasmids was made by the Promega "Magic Miniprep" according to the manufacturers instructions. Restriction analysis indicated that the plasmids contained inserts ranging in size from 1 kb to 2.5 kb. The alkali-denatured double-stranded DNA from  
 15   one of these, designated pSF2-169K contained an insert of 1.6 kb, was sequenced as described above. The nucleotide sequence of the cDNA insert in plasmid pSF2-169K shown in SEQ ID NO:5.

Cloning of a Corn (*Zea mays*)

20       cDNA Encoding Seed Microsomal Delta-12  
           Fatty Acid Desaturase

      Corn microsomal delta-12 desaturase cDNA was isolated using a PCR approach. For this, a cDNA library was made to poly A<sup>+</sup> RNA from developing corn embryos in  
 25   Lambda ZAP II vector (Stratagene). 5-10 ul of this library was used as a template for PCR using 100 pmol each of two sets of degenerate oligomers NS3 (SEQ ID NO:13) and equimolar amounts of RB5a/b (that is, equimolar amounts of SEQ ID NOS:16/17) as sense and  
 30   antisense primers, respectively. NS3 and RB5a/b correspond to stretches of amino acids 101-109 and 318-326, respectively, of SEQ ID NO:2, which are conserved in most microsomal delta-12 desaturases (SEQ ID NOS:2, 4, 6, 8). PCR was carried out using the PCR  
 35   kit (Perkin-Elmer) using 40 cycles of 94°C 1 min, 45°C,

1 min, and 55°C, 2 min. Analyses of the PCR products on an agarose gel showed the presence of a product of the expected size (720 bp), which was absent in control reactions containing either the sense or antisense primers alone. The PCR product fragment was gel purified and then used as a probe for screening the same corn cDNA library at 60°C as described above. One positively-hybridizing plaque was purified and partial sequence determination of its cDNA showed it to be a nucleotide sequence encoding microsomal delta-12 desaturase but truncated at the 3' end. The cDNA insert encoding the partial desaturase was gel isolated and used to probe the corn cDNA library again. Several positive plaques were recovered and characterized. DNA sequence analysis revealed that all of these clones seem to represent the same sequence with the different length of 5' or 3' ends. The clone containing the longest insert, designated pFad2#1, was sequenced completely. SEQ ID NO:7 shows the 5' to 3' nucleotide sequence of 1790 base pairs of corn (*Zea mays*) cDNA which encodes microsomal delta-12 desaturase in plasmid pFad2#1. Nucleotides 165 to 167 and nucleotides 1326 to 1328 are, respectively, the putative initiation codon and the termination codon of the open reading frame (nucleotides 164 to 1328). SEQ ID NO:8 is the 387 amino acid protein sequence deduced from the open reading frame (nucleotides 164 to 1328) in SEQ ID NO:7. The deduced amino acid sequence of the polypeptide shared overall identities of 71%, 40%, and 38% to *Arabidopsis* microsomal delta-12 desaturase, *Arabidopsis* microsomal delta-15 desaturase, and *Arabidopsis* plastid delta-15 desaturase, respectively. Furthermore, it lacked an N-terminal amino acid extension that would indicate it is a plastid enzyme. Based on these considerations, it

09697379.102600

is concluded that it encodes a microsomal delta-12 desaturase.

Cloning of a cDNA Encoding A Microsomal Delta-12 Desaturase and of cDNAs Encoding Microsomal Delta-12 Desaturase-Related Enzymes from Castor Bean Seed

- 5 Castor microsomal delta-12 desaturase cDNA was isolated using a RT-PCR approach. Polysomal mRNA was isolated from castor beans of stages I-II (5-10 DAP) and also from castor beans of stages IV-V (20-25 DAP).
- 10 Ten ng of each mRNA was used for separate RT-PCR reactions, using the Perkin-Elmer RT-PCR kit with the reagent concentration as recommended by the kit protocol. The reverse transcriptase reaction was primed with random hexamers and the PCR reaction with 100 pmol
- 15 each of the degenerate delta-12 desaturase primers NS3 and NS9 (SEQ ID NOS:13 and 14, respectively). The reverse transcriptase reaction was incubated at 25°C for 10 min, 42°C for 15 min, 99°C for 5 min and 5°C for 5 min. The PCR reaction was incubated at 95°C for 2 min
- 20 followed by 35 cycles of 95°C for 1 min/50°C for 1 min. A final incubation at 60°C for 7 min completed the reaction. A DNA fragment of 720 bp was amplified from both stage I-II and stage IV-V mRNA. The amplified DNA fragment from one of the reactions was gel purified and
- 25 cloned into a pGEM-T vector using the Promega pGEM-T PCR cloning kit to create the plasmid pRF2-1C. The 720 bp insert in pRF2-1C was sequenced, as described above, and the resulting DNA sequence is shown in SEQ ID NO:9. The DNA sequence in SEQ ID NO:9 contains an open-reading
- 30 frame encoding 219 amino acids (SEQ ID NO:10), which has 81% identity (90% similarity) with amino acids 135 to 353 of the *Arabidopsis* microsomal delta-12 desaturase described in SEQ ID NO:2. The cDNA insert in pRF2-1C is therefore a 673 bp fragment of a full-length cDNA
- 35 encoding a castor bean seed microsomal delta-12

09697379.102600



desaturase. The full length castor bean seed microsomal delta-12 desaturase cDNA may isolated by screening a castor seed cDNA library, at 60°C, with the labeled insert of pRF2-1C as described in the example above.

- 5 The insert in pRF2-1C may also be used to screen castor bean libraries at lower temperatures to isolate delta-12 desaturase related sequences, such as the delta-12 hydroxylase.

- A cDNA library made to poly A<sup>+</sup> mRNA isolated from  
10 developing castor beans (stages IV-V, 20-25 DAP) was screened as described above. Radiolabeled probe prepared from pSF2b or pRF2-1C, as described above, were added, and allowed to hybridize for 18 h at 50°C. The filters were washed as described above. Autoradiography  
15 of the filters indicated that there were numerous hybridizing plaques, which appeared either strongly hybridising or weakly hybridising. Three of the strongly hybridising plaques (190A-41, 190A-42 and 190A-44) and three of the weakly hybridising plaques,  
20 (190B-41, 190b-43 and 197c-42), were plaque purified using the methods described above. The cDNA insert size of the purified phages were determined by PCR amplification of the insert using phage as template and lambda-gt11 oligomers (Clontech lambda-gt11 Amplimers)  
25 for primers. The PCR-amplified inserts of the amplified phages were subcloned into pBluescript (Pharmacia) which had been cut with Eco RI and filled in with Klenow (Sambrook et al. (Molecular Cloning, A Laboratory Approach, 2nd. ed. (1989) Cold Spring Harbor Laboratory  
30 Press). The resulting plasmids were called pRF190a-41, pRF190a-42, pRF190a-44, pRF190b-41, pRF190b-43 and pRF197c-42. All of the inserts were about 1.1 kb with the exception of pRF197c-42 which was approx. 1.5 kb. The inserts in the plasmids were sequenced as described  
35 above. The insert in pRF190b-43 did not contain any

09697379.102600

open reading frame and was not identified. The inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 were identical. The insert in pRF197c-42 contained all of the nucleotides of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 plus an additional approx. 400 bp. It was deduced therefore that the insert in pRF197c-42 was a longer version of the inserts in pRF190a-41, pRF190a-42, pRF190a-44 and pRF190b-41 and all were derived from the same full-length mRNA. The complete cDNA sequence of the insert in plasmid pRF197c-42 is shown in SEQ ID NO:11. The deduced amino acid sequence of SEQ ID NO:11, shown in SEQ ID NO:12, is 78.5% identical (90% similarity) to the castor microsomal delta-12 desaturase described above (SEQ ID NO:10) and 66% identical (80% similarity) to the Arabidopsis delta-12 desaturase amino acid sequence in SEQ ID NO:2. These similarities confirm that pRF197c-42 is a castor bean seed cDNA that encodes a microsomal delta-12 desaturase or a microsomal delta-12 desaturase-related enzyme, such as a delta-12 hydroxylase. Specific PCR primers for pRF2-1C and pRF197c-42 were made. For pRF2-1c the upstream primer was bases 180 to 197 of the cDNA sequence in SEQ ID NO:9. For pRF197c-42 the upstream primer was bases 717 to 743 of the cDNA sequence in SEQ ID NO:11. A common downstream primer was made corresponding to the exact complement of the nucleotides 463 to 478 of the sequence described in SEQ ID NO:9. Using RT-PCR with random hexamers and the above primers, and the incubation temperatures described above, it was observed that mRNA which gave rise to the cDNA contained in pRF2-1C is present in both Stage I-II and Stage IV-V castor bean seeds whereas mRNA which gave rise to the cDNA contained in plasmid pRF197c-42 is present only in Stage IV-V castor bean seeds, i.e., it is only expressed in tissue actively synthesizing

09697379.102600

ricinoleic acid. Thus it is possible that this cDNA encodes a delta-12 hydroxylase.

Clones such as pRF2-1C and pRF197c-42, and other clones from the differential screening, which, based on their DNA sequence, are less related to castor bean seed microsomal delta-12 desaturases and are not any of the known fatty-acid desaturases described above or in WO 9311245, may be expressed, for example, in soybean embryos or another suitable plant tissue, or in a microorganism, such as yeast, which does not normally contain ricinoleic acid, using suitable expression vectors and transformation protocols. The presence of novel ricinoleic acid in the transformed tissue(s) expressing the castor cDNA would confirm the identity of the castor cDNA as DNA encoding for an oleate hydroxylase.

#### EXAMPLE 4

#### USE OF THE ARABIDOPSIS THALIANA DELTA-12 DESATURASE GENOMIC CLONE AS A RESTRICTION FRAGMENT LENGTH POLYMORPHISM (RFLP) MARKER TO MAP THE DELTA-12 DESATURASE LOCUS IN ARABIDOPSIS

The gene encoding *Arabidopsis* microsomal delta-12 desaturase was used to map the genetic locus encoding the microsomal delta-12 desaturase of *Arabidopsis thaliana*. pSF2b cDNA insert encoding *Arabidopsis* microsomal delta-12 desaturase DNA was radiolabeled and used to screen an *Arabidopsis* genomic DNA library. DNA from several pure strongly-hybridizing phages was isolated. Southern blot analysis of the DNA from different phages using radiolabeled pSF2b cDNA insert as the probe identified a 6 kb Hind III insert fragment to contain the coding region of the gene. This fragment was subcloned in pBluescript vector to result in plasmid pAGF2-6 and used for partial sequence determination. This sequence (SEQ ID NO:15) confirmed that it is the

microsomal delta-12 desaturase gene. DNA from two phages was isolated and labelled with  $^{32}\text{P}$  using a random priming kit from Pharmacia under conditions recommended by the manufacturer. The radioactive DNA was used to

5 probe a Southern blot containing genomic DNA from Arabidopsis thaliana (ecotype Wassileskija and marker line W100 ecotype Landesberg background) digested with one of several restriction endonucleases. Following hybridization and washes under standard conditions

10 (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. (1989) Cold Spring Harbor Laboratory Press), autoradiograms were obtained. A different pattern of hybridization (polymorphism) was identified in Hind III-digested genomic DNAs using one of the phage

15 DNAs. This polymorphism was located to a 7 kB Hind III fragment in the phage DNA that revealed the polymorphism. The 7 kb fragment was subcloned in pBluescript vector to result in plasmid pAGF2-7. Plasmid pAGF2-7 was restricted with Hind III enzyme and

20 used as a radiolabelled probe to map the polymorphism essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The radiolabelled DNA fragment was applied as described above to Southern blots of Hind III-digested genomic DNA isolated from 117

25 recombinant inbred progeny (derived from single-seed descent lines to the  $F_6$  generation) resulting from a cross between Arabidopsis thaliana marker line W100 and ecotype Wassileskija (Burr et al., Genetics (1988) 118:519-526). The bands on the autoradiograms were

30 interpreted as resulting from inheritance of either paternal (ecotype Wassileskija) or maternal (marker line W100) DNA or both (a heterozygote). The resulting segregation data were subjected to genetic analysis using the computer program Mapmaker (Lander et al.,

35 Genomics (1987) 1:174-181). In conjunction with

09697379.102600

previously obtained segregation data for 63 anonymous RFLP markers and 9 morphological markers in Arabidopsis thaliana (Chang et al., Proc. Natl. Acad. Sci. USA (1988) 85:6856-6860; Nam et al., Plant Cell (1989) 1:699-705), a single genetic locus was positioned corresponding to the microsomal delta-12 desaturase gene. The location of the microsomal delta-12 desaturase gene was thus determined to be 13.6 cM proximal to locus c3838, 9.2 cM distal to locus 1At228, and 4.9 cM proximal to FadD locus on chromosome 3 [Koorneef, M. et. al. (1993) in Genetic Maps, Ed. O'Brien, S. J.; Yadav et al. (1993) Plant Physiology 103:467-476.]

#### EXAMPLE 5

#### 15     USE OF SOYBEAN MICROSOMAL DELTA-12 DESATURASE cDNA           SEQUENCE AS A RESTRICTION FRAGMENT           LENGTH POLYMORPHISM (RFLP) MARKER

The 1.6 kb insert obtained from the plasmid pSF2-169K as previously described was radiolabelled with <sup>32</sup>P using a Random Priming Kit from Bethesda Research Laboratories under conditions recommended by the manufacturer. The resulting radioactive probe was used to probe a Southern blot (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed. (1989) Cold Spring Harbor Laboratory Press) containing genomic DNA from soybean (Glycine max (cultivar Bonus) and Glycine soja (PI81762)) digested with one of several restriction enzymes. After hybridization and washes under low stringency conditions (50 mM Tris, pH 7.5, 6X SSPE, 10% dextran sulfate, 1% SDS at 56°C for the hybridization and initial washes, changing to 2X SSPE and 0.1% SDS for the final wash), autoradiograms were obtained, and different patterns of hybridization (polymorphisms) were identified in digests performed with restriction enzymes Hind III and Eco RI. These polymorphisms were used to

map two pSF2-169k loci relative to other loci on the soybean genome essentially as described by Helentjaris et al., (Theor. Appl. Genet. (1986) 72:761-769). The map positions of the polymorphisms were determined to be in linkage group 11 between 4404.00 and 1503.00 loci (4.5 cM and 7.1 cM from 4404.00 and 1503.00, respectively) and linkage group 19 between 4010.00 and 5302.00 loci (1.9 cM and 2.7 cM from 4010.00 and 5302.00, respectively) [Rafalski, A. and Tingey, S. (1993) in Genetic Maps, Ed. O' Brien, S. J.].

#### EXAMPLE 6

#### EXPRESSION OF MICROSOMAL DELTA-12 DESATURASE IN SOYBEANS

##### Construction of Vectors for Transformation of

##### Glycine max for Reduced Expression of

##### Microsomal Delta-12 Desaturases in Developing Soybean Seeds

Plasmids containing the antisense G. max microsomal delta-12 desaturase cDNA sequence under control of the soybean Kunitz Trypsin Inhibitor 3 (KTI3) promoter (Jofuku and Goldberg, Plant Cell (1989) 1:1079-1093), the Phaseolus vulgaris 7S seed storage protein (phaseolin) promoter (Sengupta-Gopalan et al., Proc. Natl. Acad. Sci. USA (1985) 82:3320-3324; Hoffman et al., Plant Mol. Biol. (1988) 11:717-729) and soybean beta-conglycinin promoter (Beachy et al., EMBO J. (1985) 4:3047-3053), were constructed. The construction of vectors expressing the soybean delta-12 desaturase antisense cDNA under the control of these promoters was facilitated by the use of the following plasmids: pML70, pCW108 and pCW109A.

The pML70 vector contains the KTI3 promoter and the KTI3 3' untranslated region and was derived from the commercially available vector pTZ18R (Pharmacia) via the intermediate plasmids pML51, pML55, pML64 and pML65. A 2.4 kb Bst BI/Eco RI fragment of the complete soybean

KTi3 gene (Jofuku and Goldberg (1989) Plant Cell 1:1079-1093), which contains all 2039 nucleotides of the 5' untranslated region and 390 bases of the coding sequence of the KTi3 gene ending at the Eco RI site corresponding to bases 755 to 761 of the sequence described in Jofuku et al (1989) Plant Cell 1:427-435, was ligated into the Acc I/Eco RI sites of pTZ18R to create the plasmid pML51. The plasmid pML51 was cut with Nco I, filled in using Klenow, and religated, to destroy an Nco I site in the middle of the 5' untranslated region of the KTi3 insert, resulting in the plasmid pML55. The plasmid pML55 was partially digested with Xmn I/Eco RI to release a 0.42 kb fragment, corresponding to bases 732 to 755 of the above cited sequence, which was discarded. A synthetic Xmn I/Eco RI linker containing an Nco I site, was constructed by making a dimer of complementary synthetic oligo-nucleotides consisting of the coding sequence for an Xmn I site (5'-TCTTCC-3') and an Nco I site (5'-CCATGGG-3') followed directly by part of an Eco RI site (5'-GAAGG-3'). The Xmn I and Nco I/Eco RI sites were linked by a short intervening sequence (5'-ATAGCCCCCAA-3'). This synthetic linker was ligated into the Xmn I/Eco RI sites of the 4.94 kb fragment to create the plasmid pML64. The 3' untranslated region of the KTi3 gene was amplified from the sequence described in Jofuku et al (Ibid.) by standard PCR protocols (Perkin Elmer Cetus, GeneAmp PCR kit) using the primers ML51 and ML52. Primer ML51 contained the 20 nucleotides corresponding to bases 1072 to 1091 of the above cited sequence with the addition of nucleotides corresponding to Eco RV (5'-GATATC-3'), Nco I (5'-CCATGG-3'), Xba I (5'-TCTAGA-3'), Sma I (5'-CCCGGG-3') and Kpn I (5'-GGTACC-3') sites at the 5' end of the primer. Primer ML52 contained to the exact compliment of the

09697379.102600

09697379.102600

nucleotides corresponding to bases 1242 to 1259 of the above cited sequence with the addition of nucleotides corresponding to Sma I (5'-CCCGGG-3'), Eco RI (5'-GAATTC-3'), Bam HI (5'-GGATCC-3') and Sal I (5'-GTCGAC-3') sites at the 5' end of the primer. The PCR-amplified 3' end of the KTi3 gene was ligated into the Nco I/Eco RI sites of pML64 to create the plasmid pML65. A synthetic multiple cloning site linker was constructed by making a dimer of complementary synthetic oligonucleotides consisting of the coding sequence for Pst I (5'-CTGCA-3'), Sal I (5'-GTCGAC-3'), Bam HI (5'-GGATCC-3') and Pst I (5'-CTGCA-3') sites. The linker was ligated into the Pst I site (directly 5' to the KTi3 promoter region) of pML65 to create the plasmid pML70.

The 1.46 kb Sma I/Kpn I fragment from pSF2-169K (soybean delta-12 desaturase cDNA described above) was ligated into the corresponding sites in pML70 resulting in the plasmid pBS10. The desaturase cDNA fragment was in the reverse (antisense) orientation with respect to the KTi3 promoter in pBS10. The plasmid pBS10 was digested with Bam HI and a 3.47 kb fragment, representing the KTi3 promoter/antisense desaturase cDNA/KTi3-3' end transcriptional unit was isolated by agarose gel electrophoresis. The vector pML18 consists of the non-tissue specific and constitutive cauliflower mosaic virus (35S) promoter (Odell et al., Nature (1985) 313:810-812; Hull et al., Virology (1987) 86:482-493), driving expression of the neomycin phosphotransferase gene described in (Beck et al. (1982) Gene 19:327-336) followed by the 3' end of the nopaline synthase gene including nucleotides 848 to 1550 described by (Depicker et al., (1982) J. Appl. Genet. 1:561-574). This transcriptional unit was inserted into the commercial cloning vector pGEM9Z (Gibco-BRL) and is flanked at the



5' end of the 35S promoter by the restriction sites Sal I, Xba I, Bam HI and Sma I in that order. An additional Sal I site is present at the 3' end of the NOS 3' sequence and the Xba I, Bam HI and Sal I sites are unique. The 3.47 kb transcriptional unit released from pBS10 was ligated into the Bam HI site of the vector pML18. When the resulting plasmids were double digested with Sma I and Kpn I, plasmids containing inserts in the desired orientation yielded 3 fragments of 5.74, 2.69 and 1.46 kb. A plasmid with the transcriptional unit in the correct orientation was selected and was designated pBS13.

The pCW108 vector contains the bean phaseolin promoter and 3' untranslated region and was derived from the commercially available pUC18 plasmid (Gibco-BRL) via plasmids AS3 and pCW104. Plasmid AS3 contains 495 base pairs of the bean (*Phaseolus vulgaris*) phaseolin (7S seed storage protein) promoter starting with 5'-TGGTCTTTTGGT-3' followed by the entire 1175 base pairs of the 3' untranslated region of the same gene (see sequence descriptions in Doyle et al., (1986) J. Biol. Chem. 261:9228-9238 and Slightom et al., (1983) Proc. Natl. Acad. Sci. USA, 80:1897-1901. Further sequence description may be found in WO 9113993) cloned into the Hind III site of pUC18. The additional cloning sites of the pUC18 multiple cloning region (Eco RI, Sph I, Pst I and Sal I) were removed by digesting with Eco RI and Sal I, filling in the ends with Klenow and religating to yield the plasmid pCW104. A new multiple cloning site was created between the 495bp of the 5' phaseolin and the 1175bp of the 3' phaseolin by inserting a dimer of complementary synthetic oligo-nucleotides consisting of the coding sequence for a Nco I site (5'-CCATGG-3') followed by three filler bases (5'-TAG-3'), the coding sequence for a Sma I site

(5'-CCCGGG-3'), the last three bases of a Kpn I site (5'-TAC-3'), a cytosine and the coding sequence for an Xba I site (5'-TCTAGA-3') to create the plasmid pCW108. This plasmid contains unique Nco I, Sma I, Kpn I and Xba I sites directly behind the phaseolin promoter. The 1.4 kb Eco RV/Sma I fragment from pSF2-169K was ligated into the Sma I site of the commercially available phagemid pBC SK+ (Stratagene). A phagemid with the cDNA in the desired orientation was selected by digesting with Pfl MI/Xho I to yield fragments of approx. 1 kb and 4 kb and designated pM1-SF2. The 1.4 kb Xmn I/Xba I fragment from pM1-SF2 was inserted into the Sma I/Xba I sites of pCW108 to yield the plasmid pBS11, which has the soybean delta-12 desaturase cDNA in the reverse (3'-5') orientation behind the phaseolin promoter. The plasmid pBS11 was digested with Bam HI and a 3.07 kb fragment, representing the phaseolin promoter/antisense desaturase cDNA/phaseolin 3' end transcriptional unit was isolated by agarose gel electrophoresis and ligated into the Hind III site of pML18 (described above). When the resulting plasmids were digested with Xba I, plasmids containing inserts in the desired orientation yielded 2 fragments of 8.01 and 1.18 kb. A plasmid with the transcriptional unit in the correct orientation was selected and was designated pBS14.

The vector pCW109A contains the soybean b-conglycinin promoter sequence and the phaseolin 3' untranslated region and is a modified version of vector pCW109 which was derived from the commercially available plasmid pUC18 (Gibco-BRL). The vector pCW109 was made by inserting into the Hind III site of the cloning vector pUC18 a 555 bp 5' non-coding region (containing the promoter region) of the b-conglycinin gene followed by the multiple cloning sequence containing the restriction endonuclease sites for Nco I, Sma I, Kpn I

and Xba I, as described for pCW108 above, then 1174 bp of the common bean phaseolin 3' untranslated region into the Hind III site (described above). The b-conglycinin promoter region used is an allele of the published b-conglycinin gene (Doyle et al., J. Biol. Chem. (1986) 261:9228-9238) due to differences at 27 nucleotide positions. Further sequence description of this gene may be found in Slightom (W0 9113993). To facilitate use in antisense constructions, the Nco I site and potential translation start site in the plasmid pCW109 was destroyed by digestion with Nco I, mung bean exonuclease digestion and re-ligation of the blunt site to give the modified plasmid pCW109A. The plasmid pCW109A was digested with Hind III and the resulting 1.84 kb fragment, which contained the b-conglycinin/antisense delta-12 desaturase cDNA/phaseolin 3' untranslated region, was gel isolated. The plasmid pML18 (described above) was digested with Xba I, filled in using Klenow and religated, in order to remove the Xba I site. The resulting plasmid was designated pBS16. The 1.84 kb fragment of plasmid pCW109A (described above) was ligated into the Hind III site of pBS16. A plasmid containing the insert in the desired orientation yielded a 3.53 kb and 4.41 kb fragment when digested with Kpn I and this plasmid was designated pCST2. The Xmn I/Xba I fragment of pML1-SF2 (described above) was ligated into the Sma I/Xba I sites of pCST2 to yield the vector pST11.

#### Transformation Of Somatic Soybean Embryo Cultures and Regeneration Of Soybean Plants

Soybean embryogenic suspension cultures were maintained in 35 mL liquid media (SB55 or SBP6) on a rotary shaker, 150 rpm, at 28°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. Cultures were subcultured every four weeks by

09697379-102600

inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures were transformed with pCS3FdSTlR by the method of particle gun bombardment (see Kline et al. (1987) Nature (London) 327:70). A DuPont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

To 50 mL of a 60 mg/mL 1 mm gold particle suspension was added (in order); 5 uL DNA(1 ug/uL), 20 uL spermidine (0.1 M), and 50 uL  $\text{CaCl}_2$  (2.5 M). The particle preparation was agitated for 3 min, spun in a microfuge for 10 sec and the supernatant removed. The DNA-coated particles were then washed once in 400 uL 70% ethanol and re suspended in 40 uL of anhydrous ethanol. The DNA/particle suspension was sonicated three times for 1 sec each. Five uL of the DNA-coated gold particles were then loaded on each macro carrier disk.

Approximately 300-400 mg of a four week old suspension culture was placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue were normally bombarded. Membrane rupture pressure was set at 1000 psi and the chamber was evacuated to a vacuum of 28 inches of mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was placed back into liquid and cultured as described above.

Eleven days post bombardment, the liquid media was exchanged with fresh SB55 containing 50 mg/mL hygromycin. The selective media was refreshed weekly. Seven weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed

and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Thus each new line was treated as independent transformation event. These suspensions can then be maintained as suspensions of embryos clustered in an immature developmental stage through subculture or regenerated into whole plants by maturation and germination of individual somatic embryos.

Transformed embryogenic clusters were removed from liquid culture and placed on a solid agar media (SB103) containing no hormones or antibiotics. Embryos were cultured for eight weeks at 26°C with mixed florescent and incandescent lights on a 16:8 h day/night schedule. During this period, individual embryos were removed from the clusters and analyzed at various stages of embryo development. After eight weeks somatic embryos become suitable for germination. For germination, eight week old embryos were removed from the maturation medium and dried in empty petri dishes for 1 to 5 days. The dried embryos were then planted in SB71-1 medium where they were allowed to germinate under the same lighting and germination conditions described above. Germinated embryos were transferred to sterile soil and grown to maturity for seed collection.

TABLE 10

Media:		B5 Vitamin Stock
SB55 and SBP6 Stock		10 g m-inositol
Solutions		100 mg nicotinic acid
(g/L):		100 mg pyridoxine HCl
MS Sulfate 100X Stock		1 g thiamine
MgSO <sub>4</sub> 7H <sub>2</sub> O	37.0	SB55 (per Liter)
MnSO <sub>4</sub> H <sub>2</sub> O	1.69	10 mL each MS stocks
ZnSO <sub>4</sub> 7H <sub>2</sub> O	0.86	1 mL B5 Vitamin stock
CuSO <sub>4</sub> 5H <sub>2</sub> O	0.0025	0.8 g NH <sub>4</sub> NO <sub>3</sub>
MS Halides 100X Stock		3.033 g KNO <sub>3</sub>
CaCl <sub>2</sub> 2H <sub>2</sub> O	44.0	1 mL 2,4-D (10mg/mL stock)
KI	0.083	60 g sucrose
CoCl <sub>2</sub> 6H <sub>2</sub> O	0.00125	0.667 g asparagine
KH <sub>2</sub> PO <sub>4</sub>	17.0	pH 5.7
H <sub>3</sub> BO <sub>3</sub>	0.62	For SBP6- substitute 0.5 mL
Na <sub>2</sub> MoO <sub>4</sub> 2H <sub>2</sub> O	0.025	2,4-D
MS FeEDTA 100X Stock		SB103 (per Liter)
Na <sub>2</sub> EDTA	3.724	MS Salts
FeSO <sub>4</sub> 7H <sub>2</sub> O	2.784	6% maltose
		750 mg MgCl <sub>2</sub>
		0.2% Gelrite
		pH 5.7
		SB71-1 (per liter)
		B5 salts
		1ml B5 vitamin stock
		3% sucrose
		750mg MgCl <sub>2</sub>
		0.2% gelrite
		pH 5.7

09697379-102600

Analysis Of Transgenic Glycine Max Embryos and  
Seeds Containing An Antisense Delta-15 Desaturase:  
Demonstration That The Phenotype Of Transgenic Soybean  
Somatic Embryos Is Predictive Of The Phenotype Of Seeds  
Derived From Plants Regenerated From Those Embryos

- While in the globular embryo state in liquid culture as described above, somatic soybean embryos contain very low amounts of triacylglycerol or storage proteins typical of maturing, zygotic soybean embryos.
- At this developmental stage, the ratio of total triacylglyceride to total polar lipid (phospholipids and glycolipid) is about 1:4, as is typical of zygotic soybean embryos at the developmental stage from which the somatic embryo culture was initiated. At the globular stage as well, the mRNAs for the prominent seed proteins (alpha' subunit of beta-conglycinin, Kunitz Trypsin Inhibitor 3 and Soybean Seed Lectin) are essentially absent. Upon transfer to hormone free media to allow differentiation to the maturing somatic embryo state as described above, triacylglycerol becomes the most abundant lipid class. As well, mRNAs for alpha'-subunit of beta-conglycinin, Kunitz Trypsin Inhibitor 3 and Soybean Seed Lectin become very abundant messages in the total mRNA population. In these respects the somatic soybean embryo system behaves very similarly to maturing zygotic soybean embryos *in vivo*, and is therefore a good and rapid model system for analyzing the phenotypic effects of modifying the expression of genes in the fatty acid biosynthesis pathway.
- Furthermore, the model system is predictive of the fatty acid composition of seeds from plants derived from transgenic embryos. Liquid culture globular embryos transformed with a vector containing a soybean microsomal delta-15 desaturase, in a reverse orientation and under the control of soybean conglycinin promoter

09697379.102600

- (pCS3FdST 1R), gave rise to mature embryos with a reduced 18:3 content (WO 9311245). A number of embryos from line A2872 (control tissue transformed with pCST) and from lines 299/1/3, 299/15/1, 303/7/1, 306/3/1, 306/4/3, 306/4/5 (line 2872 transformed with plasmid pCS3FdST1R) were analyzed for fatty acid content. Fatty acid analysis was performed as described in WO 9311245 using single embryos as the tissue source. Mature, somatic embryos from each of these lines were also regenerated into soybean plants by transfer to regeneration medium as described above. A number of seeds taken from plants regenerated from these embryo lines were analyzed for fatty acid content. The relative fatty-acid composition of embryos taken from tissue transformed with pCS3FdST1R was compared with relative fatty-acid composition of seeds taken from plants derived from embryos transformed with pCS3FdST1R. Also, relative fatty acid compositions of embryos and seeds transformed with pCS3FdST1R were compared with control tissue, transformed with pCST. In all cases where a reduced 18:3 content was seen in a transgenic embryo line, compared with the control, a reduced 18:3 content was also observed in segregating seeds of plants derived from that line, when compared with the control seed (Table 11).

TABLE 11

Antisense Delta-15 Desaturase:  
Relative 18:3 Content Of Embryos And Seeds Of Control  
(A2172) And Transgenic (299-, 303-, 306-) Soybean Lines

Soybean Line	Embryo	Embryo	Seed	Seed
	av. %18:3	lowest %18:3	av. %18:3*	lowest %18:3
A2872 (control)	12.1 (2.6)	8.5	8.9 (0.8)	8.0
299/1/3	5.6 (1.2)	4.5	4.3 (1.6)	2.5
299/15/1	8.9 (2.2)	5.2	2.5 (1.8)	1.4



303/7/1	7.3 (1.1)	5.9	4.9 (1.9)	2.8
306/3/1	7.0 (1.9)	5.3	2.4 (1.7)	1.3
306/4/3	8.5 (1.9)	6.4	4.5 (2.2)	2.7
306/4/5	7.6 (1.6)	5.6	4.6 (1.6)	2.7

\*Seeds which were segregating with wild-type phenotype and without a copy of the transgene are not included in these averages. The number in brackets is S.D., n=10.

Thus the Applicants conclude that an altered polyunsaturated fatty acid phenotype observed in a transgenic, mature somatic embryo line is predictive of an altered fatty acid composition of seeds of plants derived from that line.

Analysis Of Transgenic Glycine Max Embryos Containing An Antisense Microsomal Delta-12 Desaturase Construct

The vectors pBS13, pBS14 and pST11 contain the soybean microsomal delta-12 desaturase cDNA, in the antisense orientation, under the control of the soybean Kunitz Trypsin Inhibitor 3 (KTI3), Phaseolus phaseolin, and soybean beta-conglycinin promoters as described above. Liquid culture globular embryos transformed with vectors pBS13, pBS14 and pST11, gave rise to mature embryo lines as described above. Fatty acid analysis was performed as described in WO 9311245 using single, mature embryos as the tissue source. A number of embryos from line A2872 (control tissue transformed with pCST) and from line A2872 transformed with vectors pBS13, pBS14 and pST11 were analyzed for fatty acid content. About 30% of the transformed lines showed an increased 18:1 content when compared with control lines transformed with pCST described above, demonstrating that the delta-12 desaturase had been inhibited in these lines. The remaining transformed lines showed relative fatty acid compositions similar to those of the control line. The relative 18:1 content of the lines showing an increased 18:1 content was as high as 50% compared with

09697379.102600

a maximum of 12.5% in the control embryo lines. The average 18:1 content of embryo lines which showed an increased 18:1 content was about 35% (Table 12). In all the lines showing an increased 18:1 content there was a proportional decrease in the relative 18:2 content (Table 13).

TABLE 12

Summary Of Experiment In Which Soybean Embryos Were Transformed With Plasmids Containing A Soybean Antisense Microsomal Delta-12 Desaturase cDNA

	# of <u>Vector Lines</u>	# of lines		
		with high 18:1	highest 18:1	av. (%) 18:1
pCST (control)	---	---	12.5	10.5
pBS13	11	4	53.5	35.9
pBS14	11	2	48.7	32.6
pST11	11	3	50.1	35.9

In Table 12 the average 18:1 of transgenics is the average of all embryos transformed with a particular vector whose relative 18:1 content is greater than two standard deviations from the highest control value (12.5). The control average is the average of ten A2872 embryos (standard deviation = 1.2). The data in Table 12 are derived from Table 13 below.

TABLE 13

Relative Fatty Acid Contents Of Embryo Lines  
Transformed With Plasmids Containing A  
Soybean Antisense Delta-12 Desaturase cDNA

Embryo Line	Relative % Fatty-Acid Content				
A2872 (control)					
#	16:0	18:0	18:1	18:2	18:3
1	11.7	3.2	11.7	52.7	16.1
2	16.4	4.0	10.8	47.1	19.3
3	17.1	3.4	8.3	48.3	20.6
4	15.3	2.7	9.4	51.1	19.0
5	15.2	3.6	10.8	51.0	17.5
6	18.6	3.9	10.9	45.8	18.1
7	14.6	3.4	12.5	52.3	16.4
8	14.2	3.5	11.2	53.9	16.7
9	15.2	3.2	9.8	49.5	16.1
10	19.0	3.8	9.6	47.4	19.0
G335/4/197 (pBS13)					
#	16:0	18:0	18:1	18:2	18:3
1	12.2	3.3	42.0	23.0	17.4
2	12.4	2.7	22.4	39.0	21.9
3	12.0	3.2	42.0	23.2	18.4
G335/4/221 (pBS13)					
#	16:0	18:0	18:1	18:2	18:3
1	12.2	2.7	30.4	36.0	17.9
2	11.5	2.4	14.3	53.4	17.6
3	13.0	2.6	15.2	47.4	19.9
4	12.0	2.6	27.4	37.9	19.1
5	11.7	2.7	25.1	42.3	15.6
6	11.7	3.4	21.6	44.3	17.8
7	12.0	2.5	11.3	53.6	20.0
8	12.0	2.5	20.8	44.1	19.5
9	11.7	2.6	25.3	39.6	18.3

00697379.10500

## G335/8/174 (pBS13)

#	16:0	18:0	18:1	18:2	18:3
1	14.1	2.1	30.3	32.1	20.3
2	14.7	2.5	5.9	40.6	34.8
3	14.3	2.4	7.3	45.2	29.8

## G335/8/202 (pBS13)

#	16:0	18:0	18:1	18:2	18:3
1	11.7	1.5	30.1	32.4	23.3
2	11.4	2.3	48.5	20.6	16.1
3	12.9	2.3	46.6	17.1	19.5
4	12.7	2.6	32.0	31.1	20.5
5	12.9	1.9	41.7	23.5	18.9
6	12.3	2.6	40.1	25.6	17.9
7	11.3	2.4	53.5	16.6	14.5
8	11.4	2.5	15.5	21.7	17.8
9	10.2	2.0	45.4	23.2	18.5
10	12.8	2.2	43.2	23.5	16.9

## G335/6/42 (pBS14)

#	16:0	18:0	18:1	18:2	18:3
1	13.7	2.4	38.6	28.2	15.6
2	12.6	2.3	37.6	28.8	17.2
3	11.7	3.0	48.7	21.1	14.6

## G335/6/104 (pBS14)

#	16:0	18:0	18:1	18:2	18:3
1	13.8	2.5	30.5	35.4	16.0
2	12.3	2.3	14.6	53.2	16.4
3	12.7	2.6	27.1	36.6	20.0
4	12.6	2.2	32.1	34.9	17.4
5	12.7	2.6	23.2	41.2	19.3
6	12.6	2.2	11.7	52.5	20.1
7	13.3	2.1	23.3	41.2	18.4

## G335/1/25 (pST11)

#	16:0	18:0	18:1	18:2	18:3
1	13.7	2.8	50.7	17.5	12.1
2	14.5	3.0	41.8	23.5	15.0

009201" 64E6960

	3	13.9	2.9	49.1	16.8	13.6
	4	12.3	2.8	47.5	19.3	14.8
G335/2/7/1 (pST11)						
	#	16:0	18:0	18:1	18:2	18:3
	1	15.5	4.3	21.8	38.0	17.5
	2	17.8	4.1	22.0	39.5	14.0
	3	15.2	3.0	20.5	42.2	16.5
G335/2/118 (pST11)						
	#	16:0	18:0	18:1	18:2	18:3
	1	14.1	2.7	44.7	22.6	14.0
	2	15.8	2.8	37.7	26.9	14.8
	3	17.3	3.4	23.3	37.9	16.0

N.B. All other transformed embryos (24 lines) had fatty acid profiles similar to those of the control.

- One of these embryo lines, G335/1/25, had an average 18:2 content of less than 20% and an average 18:1 content greater than 45% (and as high as 53.5%). The Applicants expect, based on the data in Table 13, that seeds derived from plants regenerated from such lines will have an equivalent or greater increase in 18:1 content and an equivalent or greater increase decrease in 18:2 content.

Analysis Of Transgenic Glycine Max Seeds Containing  
An Antisense Delta-12 Desaturase Construct

- The vector pST11 contains the soybean microsomal delta-12 desaturase cDNA, in the antisense orientation, under the control of the soybean beta-conglycinin promoter as described above. Liquid culture globular embryos transformed with vector pST11 gave rise to mature embryo lines as described above. These embryo were germinated, as described above, without analysis. Plants regenerated from these embryos were self-fertilized and between 15 and 30 seeds were taken from each plant. These seeds were analyzed for fatty acid

composition as described in WO 9311245. The relative fatty acid content, expressed as a percentage of the five major fatty acids are shown below in Table 13-A:

**TABLE 13-A:**  
Relative Fatty Acids Contents Of Seeds Containing A  
Soybean Antisense Delta-12 Desaturase cDNA

LINE #	SEED #	16 0	18 0	18 1	18 2	18 3
336/3/4/1	F1	11.4	2.6	19.7	55.5	10.3
	F2	14.9	3.5	14.1	54.9	12.0
	F3	10.7	2.6	22.4	53.0	10.7
	F4	12.0	3.1	21.7	53.8	8.9
	F5	10.7	2.4	18.8	55.6	12.2
	F6	10.9	2.8	22.3	53.9	9.6
	F7	11.6	2.6	19.1	55.9	10.3
	F8	12.2	2.6	21.8	52.5	10.5
	F9	10.6	2.7	23.8	51.1	11.5
	F10	12.3	2.5	18.3	54.2	12.5
	F11	12.9	3.6	12.6	58.4	12.3
	F12	12.4	3.4	16.5	54.9	12.6
	F13	13.4	3.4	10.5	55.9	16.8
	F14	11.4	3.2	26.3	51.0	7.7
	F15	11.8	2.9	20.3	53.7	11.0
336/7/1/1	F16	13.2	2.6	25.8	43.1	14.8
	F17	9.6	3.2	65.4	11.9	9.5
	F18	10.1	2.6	41.8	37.0	8.0
	F19	8.6	2.4	60.3	18.7	9.4
	F20	11.4	3.2	41.5	30.6	12.9
	F21	8.7	2.9	63.2	16.9	8.2
	F22	10.8	3.0	39.4	35.2	11.6
	F23	7.8	2.5	79.9	3.4	6.8
	F24	9.0	2.4	63.0	17.5	8.1
	F25	9.5	2.6	60.4	19.6	8.0
	F26	8.8	3.5	67.2	11.5	8.7
	F27	10.4	3.1	53.7	22.5	10.2

009201-626666

	F28	11.1	3.0	47.0	27.8	11.1
	F29	10.6	2.8	47.3	30.6	8.4
	F30	9.7	2.7	59.0	18.2	10.4
	F136	11.6	3.2	27.4	49.3	8.4
	F137	10.7	2.8	24.2	53.6	8.3
	F138	9.8	2.6	45.6	32.2	9.4
	F139	12.1	3.2	47.0	27.2	10.3
	F140	9.7	2.6	50.8	28.2	8.1
	F141	8.4	3.5	66.3	12.0	9.4
	F142	10.3	2.5	30.9	48.5	7.3
	F143	9.1	3.6	72.1	7.0	7.5
	F144	9.8	3.2	58.6	18.2	9.7
	F145	8.3	2.8	71.8	9.5	7.0
	F146	11.1	2.8	25.9	51.5	8.6
	F147	9.5	3.0	43.1	34.5	9.4
	F148	10.6	2.5	26.7	50.6	9.1
	F149	11.7	3.3	20.2	55.4	9.0
336/7/3/3	F31	11.5	2.7	20.0	55.0	10.7
	F32	12.6	3.0	24.1	52.0	8.1
	F33	11.6	2.9	20.5	56.1	8.5
	F34	13.0	3.0	20.2	53.9	10.0
	F35	11.8	2.8	20.8	53.2	11.2
	F36	10.9	3.0	29.5	48.2	8.2
	F37	11.2	3.0	26.4	49.2	10.0
	F38	12.7	3.4	15.3	56.9	11.6
	F39	11.5	3.2	19.0	54.6	11.7
	F40	10.8	3.1	23.0	52.9	10.2
	F41	11.0	3.3	32.1	45.0	8.4
	F42	13.6	3.8	13.2	55.7	13.7
	F43	10.6	2.7	25.7	50.9	10.1
	F44	9.8	2.5	25.8	49.8	12.1
	F45	12.4	3.0	19.7	53.8	11.1
336/7/4/11	F46	11.4	2.9	36.4	39.4	9.8
	F47	12.8	3.2	23.4	50.6	10.1
	F48	11.0	2.8	26.0	49.6	10.4

09697379.102600

	F49	11.7	2.9	22.9	52.9	9.2
	F50	10.8	2.9	32.2	46.6	7.2
	F51	11.4	3.4	20.0	54.0	11.1
	F52	11.6	3.0	26.6	49.6	9.0
	F53	12.6	2.7	31.3	43.4	9.6
	F54	10.8	3.4	23.0	51.3	11.0
	F55	11.5	2.8	22.4	54.1	8.8
	F56	11.1	2.7	25.8	50.6	9.5
	F57	11.8	3.4	20.1	54.3	10.3
	F58	13.0	3.3	21.1	51.2	11.4
	F59	12.4	3.2	19.5	53.7	10.8
	F60	9.7	2.9	41.7	37.4	8.1
336/7/5/7	F61	11.0	3.5	43.8	31.2	10.1
	F62	12.1	3.2	18.4	56.8	9.5
	F63	10.9	3.9	48.6	26.0	10.4
	F64	12.6	3.7	14.3	58.8	10.4
	F65	9.1	2.9	61.7	16.6	9.1
	F66	9.9	3.3	59.4	17.6	9.5
	F67	9.5	2.9	59.3	16.5	11.2
	F68	10.3	4.0	46.0	28.2	11.2
	F69	11.6	2.5	30.6	46.4	8.6
	F70	10.7	3.6	54.6	22.1	8.6
	F71	12.5	4.0	16.0	56.4	10.9
	F72	11.1	2.8	42.2	34.9	8.6
	F73	11.6	3.4	23.0	53.4	8.2
	F74	12.1	3.5	15.8	58.0	10.4
	F75	10.8	3.4	51.4	24.2	9.7
336/7/7/19	F76	11.4	2.7	41.7	32.8	11.5
	F77	8.6	2.5	76.6	5.1	6.8
	F78	9.5	3.2	63.2	14.6	9.1
	F79	8.2	2.8	77.7	3.8	6.8
	F80	9.6	3.4	69.8	7.9	9.2
	F81	8.3	2.8	72.5	8.6	7.0
	F82	11.1	2.8	41.5	31.9	12.3
	F83	10.2	3.0	65.7	11.2	9.4

009697379.10600



F84	8.8	2.7	69.7	10.5	7.6
F85	8.5	2.9	75.2	5.5	7.2
F86	9.5	2.6	71.3	7.8	8.3
F87	10.1	2.9	60.5	17.1	9.0
F88	8.5	2.5	76.6	4.9	6.8
F89	9.6	2.4	68.7	10.5	8.1
F90	10.7	3.5	55.1	20.2	10.4
F150	9.1	2.9	66.9	12.5	8.0
F151	7.8	2.6	77.1	6.1	6.1
F152	9.5	2.9	29.7	48.8	8.6
F153	7.6	2.8	74.9	6.5	7.6
F154	12.4	3.1	18.6	54.7	10.7
F155	8.4	2.9	79.0	3.0	5.8
F156	10.2	3.3	51.7	25.4	9.1
F157	10.5	2.5	62.5	15.8	8.3
F158	9.6	3.3	71.1	8.5	6.9
F159	7.3	2.9	77.4	4.5	7.4
F160	12.4	3.3	20.4	54.4	9.2
F161	8.9	2.9	75.4	5.8	6.4
F162	12.0	3.7	38.4	35.2	10.4
F163	11.1	3.6	37.9	36.6	10.7
F164	8.3	3.7	77.5	3.5	6.7
F165	11.0	2.8	53.6	20.8	11.1
F91	12.8	3.7	17.3	55.3	10.6
F92	9.1	2.8	66.3	12.8	8.5
F93	8.9	2.2	70.5	10.4	7.4
F94	10.5	2.5	28.1	50.9	7.6
F95	10.9	3.2	42.8	31.9	10.8
F96	11.4	3.4	44.9	30.5	9.5
F97	11.7	2.7	35.8	39.5	9.8
F98	9.7	2.5	49.0	27.7	10.9
F99	12.3	3.5	15.9	57.1	10.8
F100	9.4	3.0	56.7	21.2	9.3
F101	9.8	2.6	64.4	13.4	9.6
F102	12.3	2.8	19.6	54.4	10.7

336/7/5/5

009201\* 62526960

336/7/7/1	F103	10.0	2.9	57.3	19.0	10.8
	F104	8.9	2.6	68.7	12.2	7.3
	F105	10.2	2.1	56.3	23.3	7.7
	F106	12.3	3.2	15.7	57.9	10.6
	F107	11.2	2.7	18.8	57.4	9.7
	F108	10.9	3.3	24.3	50.1	11.2
	F109	12.2	2.5	21.0	52.5	11.5
	F110	10.8	2.6	44.5	30.4	11.5
	F111	9.8	2.9	33.4	43.9	9.8
	F112	12.5	2.7	18.0	55.7	10.7
	F113	12.4	2.8	18.4	55.2	11.2
	F114	11.0	2.4	29.3	49.4	7.5
	F115	13.0	2.9	15.2	54.8	13.9
	F116	10.6	2.6	41.1	35.4	9.8
	F117	11.4	3.1	21.4	52.7	11.1
	F118	11.2	3.2	24.7	51.3	9.4
	F119	12.0	2.5	24.3	51.4	9.3
	F120	12.2	2.8	20.5	51.8	12.3
336/7/7/3	F121	11.3	3.0	23.9	53.0	8.4
	F122	6.7	2.5	77.7	4.4	8.0
	F123	7.8	3.2	70.1	10.2	8.4
	F124	8.4	3.4	65.1	12.9	9.7
	F125	11.5	3.2	16.3	57.6	11.0
	F126	11.5	2.7	20.1	51.5	14.2
	F127	11.6	2.6	24.8	50.8	9.7
	F128	9.2	2.6	41.5	38.1	8.7
	F129	7.7	2.6	78.4	3.6	7.0
	F130	10.4	2.6	52.6	24.9	9.4
336/3/3/1	F131	12.2	2.9	20.0	54.8	9.8
	F132	8.0	3.0	77.0	4.6	7.0
	F133	10.2	3.3	22.0	52.9	11.6
	F134	11.8	3.0	16.3	56.0	12.9
	F135	8.7	2.6	53.3	25.2	9.7
	F166	11.8	3.3	25.9	49.4	9.6
	F167	13.1	3.5	15.5	54.8	13.0

009201" 62526960

	F168	12.9	2.7	22.8	51.6	9.9
	F169	12.8	3.0	17.7	55.6	10.8
	F170	13.7	3.2	15.4	55.9	11.8
	F171	12.6	3.1	17.6	54.2	12.4
	F172	12.7	3.3	17.6	54.2	12.2
	F173	11.7	3.2	15.6	56.8	12.5
	F174	12.5	3.5	17.2	56.7	9.9
	F175	12.6	3.1	22.2	51.2	10.8
	F176	12.7	3.2	17.6	54.9	11.5
	F177	12.5	3.0	22.0	52.1	10.3
	F178	12.3	2.6	32.2	43.9	9.0
	F179	12.0	3.0	17.8	56.0	11.2
	F180	12.6	3.4	18.8	55.0	10.1
	F181	11.5	2.5	27.6	48.1	10.3
	F182	14.9	3.4	14.7	49.8	17.1
	F183	10.7	2.5	24.7	53.9	8.2
	F184	14.1	3.2	13.2	54.6	14.8
	F185	13.3	3.6	16.3	54.8	11.9
336/3/4/7	F186	11.9	4.0	12.7	58.7	12.8
	F187	11.2	2.8	21.2	56.1	8.6
	F188	12.3	3.5	12.3	58.6	13.2
	F189	12.5	3.8	12.7	58.2	12.7
	F190	12.4	3.1	13.5	55.2	15.8
	F191	13.5	3.7	12.2	55.9	14.6
	F192	12.1	2.9	16.4	56.7	11.8
	F193	11.8	3.5	14.9	56.8	13.0
	F194	13.0	3.9	12.5	56.5	14.0
	F195	11.8	2.8	19.4	53.9	12.0
	F196	12.7	3.8	11.1	58.1	14.2
	F197	12.4	3.5	15.1	56.1	12.8
	F198	12.2	3.6	16.2	57.2	10.8
	F199	13.1	3.1	13.3	57.4	13.0
	F200	11.8	4.0	13.5	58.8	11.9
	F201	11.9	4.1	12.4	59.6	11.9
	F202	13.0	4.0	12.1	56.8	14.1

09697379.102500

336/5/7/3	F203	13.0	3.6	15.0	55.6	12.8
	F204	11.8	4.0	17.9	55.3	11.0
	F205	13.5	3.9	10.8	57.0	14.7
	F206	10.2	2.5	21.3	54.3	11.6
	F207	11.6	3.5	14.4	56.6	14.0
	F208	12.4	3.4	16.6	55.1	12.4
	F209	12.0	3.0	23.9	51.4	9.6
	F210	12.8	3.0	18.0	55.3	10.8
	F211	11.2	2.7	25.6	50.3	10.0
	F212	12.8	3.2	12.1	57.6	14.3
336/7/3/9	F213	11.2	2.8	18.2	55.4	12.3
	F214	13.8	2.9	18.6	51.2	13.5
	F215	11.8	2.2	21.0	46.4	18.6
	F216	10.9	2.5	34.2	43.7	8.6
	F217	10.0	2.6	36.9	42.8	7.7
	F218	12.0	2.9	18.1	54.5	12.4
	F219	11.8	3.2	19.8	54.2	11.0
	F220	10.3	2.5	26.7	49.9	10.6
	F221	12.8	3.3	17.7	53.2	13.0
	F222	11.7	2.7	25.6	49.6	10.4
	F223	12.2	3.2	15.1	55.6	13.8
	F224	12.2	3.3	18.8	55.5	10.2
	F225	11.4	2.4	27.4	49.0	9.7
	F226	11.7	3.2	21.1	52.0	11.9
	F227	12.0	3.2	14.3	55.6	14.8
	F228	12.0	3.0	21.9	52.9	10.2
	F229	11.6	2.9	18.0	54.8	12.6
	F230	14.0	3.1	15.6	55.6	11.6
	F231	11.1	2.5	23.9	51.8	10.7
	F232	13.0	2.8	14.0	57.6	12.6
	F233	12.6	3.1	17.5	54.8	12.0
	F234	13.6	3.6	17.8	51.5	13.4
	F235	13.0	3.7	16.7	54.2	12.4
	F236	12.8	3.4	19.0	53.2	11.6
	F237	10.5	2.5	30.5	46.6	9.9

00697379.106600

	F238	11.5	2.8	21.5	54.5	9.5
	F239	13.8	3.0	17.1	54.6	11.6
	F240	14.3	3.9	15.3	53.8	12.7
	F241	14.2	3.4	9.9	57.9	14.5
	F242	12.5	2.9	22.8	52.1	9.7
	F243	11.9	3.3	22.4	52.4	10.0
	F244	14.5	2.5	18.4	45.0	19.6
	F245	11.3	3.2	26.5	48.4	10.6
336/7/3/13	F246	13.0	2.8	25.8	46.3	12.1
	F247	12.5	3.3	16.7	56.5	10.9
	F248	11.6	2.7	31.0	44.1	10.5
	F249	13.6	4.0	14.2	53.4	14.8
	F250	10.8	3.0	36.8	41.2	8.2
	F251	13.5	2.6	16.4	50.1	17.4
	F252	14.2	2.9	18.1	53.4	11.3
	F253	13.7	3.7	18.9	49.2	14.5
	F254	11.4	2.7	28.8	47.8	9.2
	F255	12.9	3.6	18.4	52.9	12.2
	F256	11.2	3.0	23.9	50.2	11.6
	F257	13.2	3.2	18.7	53.2	11.8
	F258	10.8	2.9	31.0	45.9	9.3
	F259	13.0	3.8	17.5	53.7	12.0
	F260	13.2	3.6	18.5	50.7	14.0
	F261	12.6	3.8	24.2	49.7	9.6
	F262	14.7	3.2	14.1	54.0	14.0
	F263	14.7	4.5	11.3	55.3	14.1
	F264	12.8	3.2	19.4	54.4	10.1
	F265	12.2	3.7	14.6	56.2	13.3
336/7/4/2	F266	12.5	3.9	46.6	28.2	8.8
	F267	9.6	2.5	61.8	18.5	7.1
	F268	10.1	2.9	62.6	13.8	9.8
	F269	12.4	3.2	25.5	49.5	9.1
	F270	8.2	3.0	72.1	9.2	6.8
	F271	14.3	3.8	14.8	57.2	9.8
	F272	12.2	3.3	17.6	49.0	17.9

009697379.102600

	F273	11.7	3.2	41.7	34.3	8.8
	F274	11.6	3.5	44.1	30.1	10.5
	F275	13.5	3.5	22.4	48.7	11.8
	F276	12.5	3.4	23.1	48.6	12.4
	F277	13.0	3.8	22.6	49.0	11.6
	F278	12.3	3.7	27.1	46.5	10.4
	F279	10.6	3.0	58.2	19.5	8.6
	F280	12.8	4.0	13.0	59.7	10.4
	F281	12.8	3.4	17.4	56.6	9.8
	F282	10.6	4.4	46.3	29.5	9.1
	F283	13.4	4.4	23.7	47.2	11.3
	F284	11.2	3.0	29.2	46.7	9.4
	F285	11.9	3.5	36.8	38.5	8.8
336/7/4/9	F286	9.9	2.6	39.6	37.8	10.1
	F287	8.9	2.7	57.3	23.6	7.5
	F288	10.4	2.6	45.2	33.2	8.0
	F289	10.0	2.9	30.3	46.3	10.5
	F290	10.7	2.5	34.9	42.4	9.3
	F291	9.9	2.9	52.8	25.2	8.7
	F292	10.5	2.4	39.0	38.1	9.6
	F293	9.2	2.7	33.8	43.9	10.2
	F294	11.6	2.7	29.0	46.6	9.9
	F295	12.2	2.8	18.0	56.6	10.2
	F296	10.7	2.7	17.4	56.6	12.5
	F297	10.4	2.4	27.1	41.9	7.7
	F298	10.4	2.8	19.8	58.1	8.7
	F299	10.9	2.6	18.0	57.9	10.3
	F300	9.4	2.4	40.4	39.4	7.8
	F301	9.5	2.7	45.8	33.5	8.0
	F302	10.0	2.5	40.1	38.0	9.4
	F303	9.4	2.6	45.1	33.2	9.1
	F304	10.6	2.8	22.9	51.9	11.8
	F305	10.1	2.3	34.5	45.1	7.8
336/7/6/10	F306	8.8	3.4	71.4	9.1	7.3
	F307	13.0	3.0	20.5	56.0	7.4

09697379.102600

336/7/7/9

F308	7.8	3.0	76.3	5.6	6.8
F309	11.1	3.1	24.4	54.2	7.1
F310	9.3	2.5	63.5	11.2	13.5
F311	9.3	3.4	70.0	9.1	8.1
F312	8.0	3.8	61.1	16.4	10.5
F313	8.6	3.3	64.1	14.5	9.5
F314	8.7	3.1	72.5	9.7	5.7
F315	9.0	3.6	58.0	20.8	8.5
F316	12.3	3.8	14.5	59.8	9.4
F317	10.1	3.9	13.6	63.6	8.8
F318	10.9	3.1	16.6	57.3	12.1
F319	8.7	2.7	74.1	7.5	6.8
F320	10.4	3.1	63.2	13.2	9.8
F321	9.6	3.2	50.4	28.2	8.6
F322	7.6	3.4	57.4	23.4	8.0
F323	9.1	2.8	66.0	14.6	7.4
F324	10.3	3.3	60.0	15.6	10.7
F325	10.1	3.5	53.7	21.8	10.7
F326	12.3	3.9	16.5	57.8	9.4
F327	8.4	3.1	72.8	7.6	7.4
F328	11.2	2.9	34.6	43.4	7.8
F329	9.5	3.3	60.8	16.6	9.8
F330	11.7	3.2	31.5	44.7	8.7
F331	12.2	2.9	22.8	53.3	8.8
F332	9.7	2.6	52.9	26.6	8.2
F333	12.2	3.3	28.7	47.6	8.2
F334	10.4	3.6	54.3	21.8	9.8
F335	10.0	3.3	64.9	11.8	10.0
F336	9.0	3.0	73.2	6.0	8.7
F337	9.2	2.7	68.8	10.1	9.2
F338	9.9	3.0	62.7	14.7	9.6
F339	8.5	3.2	72.2	6.6	9.4
F340	12.8	3.7	15.2	55.2	13.0
F341	13.2	2.8	22.5	49.8	11.7
F342	11.2	4.2	42.1	30.8	11.6

009201.6267980

	F343	12.6	3.9	26.8	44.9	11.7
	F344	9.1	3.3	66.6	10.8	9.9
	F345	13.8	4.3	17.9	43.0	21.0
336/7/7/17	F346	9.9	3.2	58.2	17.5	11.2
	F347	9.1	2.8	75.0	4.8	7.8
	F348	8.8	2.8	64.1	14.0	10.2
	F349	9.7	2.9	56.4	20.0	11.0
	F350	13.9	3.9	12.8	51.1	18.2
	F351	9.2	3.3	70.0	7.8	9.6
	F352	11.3	2.5	44.4	32.9	8.9
	F353	12.2	3.0	20.6	53.8	10.4
	F354	12.0	2.8	22.1	52.8	10.4
	F355	14.1	3.5	15.2	54.2	13.1
	F356	12.7	3.0	19.7	54.2	10.4
	F357	8.0	2.4	71.4	11.2	7.0
	F358	13.0	3.4	14.3	56.1	13.1
	F359	8.3	3.0	69.4	10.3	8.8
	F360	8.7	2.6	75.8	4.7	7.8
	F361	13.1	3.3	25.4	44.8	13.3
	F362	10.6	3.6	56.0	17.2	12.6
	F363	13.9	4.0	12.6	55.5	14.0
	F364	12.9	3.8	20.7	51.0	11.4
	F365	12.7	2.6	29.0	42.7	12.8

The 18:1 content of 95 seeds taken from non-transformed A2872 soybean seeds ranged from 9.9% to 32.2% with a mean 18:1 content of 18.4%. Segregating seeds of plants derived from embryos transformed with pST11 ranged from 12.6% to 79.9% with a mean of 44.6% (Table 13-A). Seeds which had an elevated 18:1 content relative to untransformed plants also had a reduced total saturate content. For example, the total saturate content of untransformed A2872 seeds is around 15% (11% 16:0, 4% 18:0). The total saturate content of seed F122 (line 336/7/7/3, 18:1 content 77.7%) was 9.2% (6.7% 16:0, 2.5% 18:0).



EXAMPLE 7EXPRESSION OF MICROSOMAL DELTA-12 DESATURASE IN CANOLAConstruction Of Vectors For Transformation ofBrassica Napus For Reduced Expression of5 Microsomal Delta-12 Desaturasesin Developing Canola Seeds

An extended poly A tail was removed from the canola delta-12 desaturase sequence contained in plasmid pCF2-165D and additional restriction sites for cloning were introduced as follows. A PCR primer was synthesized corresponding to bases 354 through 371 of SEQ ID NO:3. The second PCR primer was synthesized as the complement to bases 1253 through 1231 with 15 additional bases (GCAGATATCGCGGCC) added to the 5' end. The additional bases encode both an EcoRV site and a NotI site. pCF2-165D was used as the template for PCR amplification using these primers. The 914 base pair product of PCR amplification was digested with EcoRV and PflMI to give an 812 base pair product corresponding to bases 450 through 1253 of pCF2-165D with the added NotI site.

pCF2-165D was digested with PstI, the PstI overhang was blunted with Klenow fragment and then digested with PflMI. The 3.5 kB fragment corresponding to pBluescript along with the 5' 450 bases of the canola Fad2 cDNA was gel purified and ligated to the above described 812 base pair fragment. The ligation product was amplified by transformation of E. coli and plasmid DNA isolation. The EcoRI site remaining at the cloning junction between pBluescript and the canola Fad2 cDNA was destroyed by digestion, blunting and religation. The recovered plasmid was called pM2CFd2.

pM2CFd2 was digested with EcoRV and SmaI to remove the Fad2 insert as a blunt ended fragment. The fragment was gel purified and cloned into the SmaI site of pBC

09697379.102600

(Stratagene, La Jolla, CA). A plasmid with the NotI site introduced by PCR oriented away from the existing NotI site in pBC was identified by NotI digestion and gel fractionation of the digests. The resulting construct then had NotI sites at both ends of the canola Fad2 cDNA fragment and was called pM3CFd2.

- Vectors for transformation of the antisense cytoplasmic delta-12 desaturase constructions under control of the  $\beta$ -conglycinin, Kunitz trypsin inhibitor III, napin and phaseolin promoters into plants using Agrobacterium tumefaciens were produced by constructing a binary Ti plasmid vector system (Bevan, (1984) Nucl. Acids Res. 12:8711-8720). One starting vector for the system, (pZS199) is based on a vector which contains:
- (1) the chimeric gene nopaline synthase/neomycin phosphotransferase as a selectable marker for transformed plant cells (Brevan et al. (1984) Nature 304: 184-186),
  - (2) the left and right borders of the T-DNA of the Ti plasmid (Brevan et al. (1984) Nucl. Acids Res. 12:8711-8720),
  - (3) the E. coli lacZ a-complementing segment (Vieria and Messing (1982) Gene 19:259-267) with unique restriction endonuclease sites for Eco RI, Kpn I, Bam HI, and Sal I,
  - (4) the bacterial replication origin from the Pseudomonas plasmid pVS1 (Itoh et al. (1984) Plasmid 11:206-220), and
  - (5) the bacterial neomycin phosphotransferase gene from Tn5 (Berg et al. (1975) Proc. Natnl. Acad. Sci. U.S.A. 72:3628-3632) as a selectable marker for transformed A. tumefaciens. The nopaline synthase promoter in the plant selectable marker was replaced by the 35S promoter (Odell et al. (1985) Nature, 313:810-813) by a standard restriction endonuclease digestion and ligation strategy. The 35S promoter is required for efficient Brassica napus transformation as described below. A second vector (pZS212) was constructed by reversing the

order of restriction sites in the unique site cloning region of pZS199

Canola napin promoter expression cassettes were constructed as follows: Ten oligonucleotide primers were synthesized based upon the nucleotide sequence of napin lambda clone CGN1-2 published in European Patent Application EP 255378). The oligonucleotide sequences were:

- BR42 and BR43 corresponding to bases 1132 to 1156 (BR42) and the complement of bases 2248 to 2271 (BR43) of the sequence listed in Figure 2 of EP 255378.
- BR45 and BR46 corresponding to bases 1150 to 1170 (BR46) and the complement of bases 2120 to 2155 (BR45) of the sequence listed in Figure 2 of EP 255378. In addition BR46 had bases corresponding to a Sal I site (5'-GTCGAC-3') and a few additional bases (5'-TCAGGCCT-3') at its 5' end and BR45 had bases corresponding to a Bgl II site (5'-AGATCT-3') and two (5'-CT-3') additional bases at the 5' end of the primer,
- BR47 and BR48 corresponding to bases 2705 to 2723 (BR47) and bases 2643 to 2666 (BR48) of the sequence listed in Figure 2 of EP 255378. In addition BR47 had two (5'-CT-3') additional bases at the 5' end of the primer followed by bases corresponding to a Bgl II site (5'-AGATCT-3') followed by a few additional bases (5'-TCAGGCCT-3'),
- BR49 and BR50 corresponding to the complement of bases 3877 to 3897 (BR49) and the complement of bases 3985 to 3919 (BR50) of the sequence listed in Figure 2 of EP 255378. In addition BR49 had bases corresponding to a Sal I site (5'-GTCGAC-3') and a few additional bases (5'-TCAGGCCT-3') at its 5' end,
- BR57 and BR58 corresponding to the complement of bases 3875 to 3888 (BR57) and bases 2700 to 2714 (BR58) of

09697379-102600

the sequence listed in Figure 2 of EP 255378. In addition the 5' end of BR57 had some extra bases (5'-CCATGG-3') followed by bases corresponding to a Sac I site (5'-GAGCTC-3') followed by more additional  
 5 bases (5'-GTCGACGAGG-3'). The 5' end of BR58 had additional bases (5'-GAGCTC-3') followed by bases corresponding to a Nco I site (5'-CCATGG-3') followed by additional bases (5'-AGATCTGGTACC-3').

- BR61 and BR62 corresponding to bases 1846 to 1865  
 10 (BR61) and bases 2094 to 2114 (BR62) of the sequence listed in Figure 2 of EP 255378. In addition the 5' end of BR 62 had additional bases (5'-GACA-3') followed by bases corresponding to a Bgl II site (5'-AGATCT-3') followed by a few additional bases  
 15 (5'-GCGGCCGC-3').

Genomic DNA from the canola variety 'Hyola401' (Zeneca Seeds) was used as a template for PCR amplification of the napin promoter and napin terminator regions. The promoter was first amplified using primers  
 20 BR42 and BR43, and reamplified using primers BR45 and BR46. Plasmid pIMC01 was derived by digestion of the 1.0 kb promoter PCR product with SalI/BglII and ligation into SalI/BamHI digested pBluescript SK<sup>+</sup> (Stratagene). The napin terminator region was amplified using primers  
 25 BR48 and BR50, and reamplified using primers BR47 and BR49. Plasmid pIMC06 was derived by digestion of the 1.2 kb terminator PCR product with SalI/BglII and ligation into SalI/BglII digested pSP72 (Promega). Using pIMC06 as a template, the terminator region was  
 30 reamplified by PCR using primer BR57 and primer BR58. Plasmid pIMC101 containing both the napin promoter and terminator was generated by digestion of the PCR product with SacI/NcoI and ligation into SacI/NcoI digested pIMC01. Plasmid pIMC101 contains a 2.2 kb napin  
 35 expression cassette including complete napin 5' and 3'

09697379.102600

non-translated sequences and an introduced NcoI site at the translation start ATG. Primer BR61 and primer BR62 were used to PCR amplify an ~270 bp fragment from the 3' end of the napin promoter. Plasmid pIMC401 was obtained  
5 by digestion of the resultant PCR product with EcoRI/BglII and ligation into EcoRI/BglII digested pIMC101. Plasmid pIMC401 contains a 2.2 kb napin expression cassette lacking the napin 5' non-translated sequence and includes a NotI site at the transcription  
10 start.

To construct the antisense expression vector, pM3CFd2 was digested with NotI as was pIMC401. The delta-12 desaturase containing insert from the digest of pM3CFd2 was gel isolated and ligated into the NotI  
15 digested and phosphatase treated pIMC401. An isolate in which the delta-12 desaturase was oriented antisense to the napin promoter was selected by digestion with XhoI and PflMI to give plasmid pNCFd2R. pNCFd2R was digested with SalI, phosphatase treated and ligated into pZS212  
20 which had been opened by the same treatment. A plasmid with desired orientation of the introduced napin:delta-12 desaturase antisense transcription unit relative to the selectable marker was chosen by digestion with PvuI and the resulting binary vector was  
25 given the name pZNCfD2R.

Plasmid pML70 (described in Example 6 above) was digested with NcoI, blunted then digested with KpnI. Plasmid pM2CFd was digested with KpnI and SmaI and the isolated fragment ligated into the opened pML70 to give  
30 the antisense expression cassette pMKCFd2R. The promoter:delta-12 desaturase:terminator sequence was removed from pMKCFd2R by BamHI digestion and ligated into pZS199 which had been BamHI digested and phosphatase treated. The desired orientation relative  
35 to the selectable marker was determined by digestion

with XhoI and PflMI to give the expression vector pZKCFd2R.

The expression vector containing the  $\beta$ -conglitin promoter was constructed by SmaI and EcoRV digestion of pM2CFd2 and ligation into SmaI cut pML109A. An isolate with the antisense orientation was identified by digestion with XhoI and PflMI, and the transcription unit was isolated by SalI and EcoRI digestion. The isolated SalI-EcoRI fragment was ligated into EcoRI-SalI digested pZS199 to give pCCFd2R.

The expression vector containing the phaseolin promoter was obtained using the same procedure with pCW108 as the starting, promoter containing vector and pZS212 as the binary portion of the vector to give pZPhCFd2R.

#### Agrobacterium-Mediated Transformation Of Brassica Napus

The binary vectors pZNCFd2R, pZCCFd2R, pZPhCFd2R, and pZNCFd2R were transferred by a freeze/thaw method (Holsters et al. (1978) Mol Gen Genet 163:181-187) to the Agrobacterium strain LBA4404/pAL4404 (Hockema et al. (1983), Nature 303:179-180).

Brassica napus cultivar "Westar" was transformed by co-cultivation of seedling pieces with disarmed Agrobacterium tumefaciens strain LBA4404 carrying the appropriate binary vector.

B. napus seeds were sterilized by stirring in 10% Chlorox, 0.1% SDS for thirty min, and then rinsed thoroughly with sterile distilled water. The seeds were germinated on sterile medium containing 30 mM CaCl<sub>2</sub> and 1.5% agar, and grown for six days in the dark at 24°C.

Liquid cultures of Agrobacterium for plant transformation were grown overnight at 28°C in Minimal A medium containing 100 mg/L kanamycin. The bacterial cells were pelleted by centrifugation and resuspended at

a concentration of  $10^8$  cells/mL in liquid Murashige and Skoog Minimal Organic medium containing 100  $\mu$ M acetosyringone.

*B. napus* seedling hypocotyls were cut into 5 mm segments which were immediately placed into the bacterial suspension. After 30 min, the hypocotyl pieces were removed from the bacterial suspension and placed onto BC-35 callus medium containing 100  $\mu$ M acetosyringone. The plant tissue and *Agrobacteria* were co-cultivated for three days at 24°C in dim light.

The co-cultivation was terminated by transferring the hypocotyl pieces to BC-35 callus medium containing 200 mg/L carbenicillin to kill the *Agrobacteria*, and 25 mg/L kanamycin to select for transformed plant cell growth. The seedling pieces were incubated on this medium for three weeks at 28°C under continuous light.

After four weeks, the segments were transferred to BS-48 regeneration medium containing 200 mg/L carbenicillin and 25 mg/L kanamycin. Plant tissue was subcultured every two weeks onto fresh selective regeneration medium, under the same culture conditions described for the callus medium. Putatively transformed calli grew rapidly on regeneration medium; as calli reached a diameter of about 2 mm, they were removed from the hypocotyl pieces and placed on the same medium lacking kanamycin.

Shoots began to appear within several weeks after transfer to BS-48 regeneration medium. As soon as the shoots formed discernable stems, they were excised from the calli, transferred to MSV-1A elongation medium, and moved to a 16:8 h photoperiod at 24°C.

Once shoots had elongated several internodes, they were cut above the agar surface and the cut ends were dipped in Rootone. Treated shoots were planted directly into wet Metro-Mix 350 soilless potting medium. The pots

were covered with plastic bags which were removed when the plants were clearly growing -- after about ten days.

Plants were grown under a 16:8 h photoperiod, with a daytime temperature of 23°C and a nighttime temperature of 17°C. When the primary flowering stem began to elongate, it was covered with a mesh pollen-containment bag to prevent outcrossing. Self-pollination was facilitated by shaking the plants several times each day. Fifty-one plants have thus far been obtained from transformations using both pZCCFd2R and pZPhCfd2R, 40 plants have been obtained from pZKCFd2R and 26 from pZNCfd2R.

#### Minimal A Bacterial Growth Medium

Dissolve in distilled water:

- 15      10.5 grams potassium phosphate, dibasic
- 4.5 grams potassium phosphate, monobasic
- 1.0 gram ammonium sulfate
- 0.5 gram sodium citrate, dihydrate
- Make up to 979 mL with distilled water
- 20      Autoclave
- Add 20 mL filter-sterilized 10% sucrose
- Add 1 mL filter-sterilized 1 M MgSO<sub>4</sub>

#### Brassica Callus Medium BC-35

Per liter:

- 25      Murashige and Skoog Minimal Organic Medium (MS salts, 100 mg/L i-inositol, 0.4 mg/L thiamine; GIBCO #510-3118)
- 30 grams sucrose
- 18 grams mannitol
- 30      0.5 mg/L 2,4-D
- 0.3 mg/L kinetin
- 0.6% agarose
- pH 5.8

#### Brassica Regeneration Medium BS-48

- 35      Murashige and Skoog Minimal Organic Medium

09697379.102600



Gamborg B5 Vitamins (SIGMA #1019)

10 grams glucose

250 mg xylose

600 mg MES

5 0.4% agarose

pH 5.7

Filter-sterilize and add after autoclaving:

2.0 mg/L zeatin

0.1 mg/L IAA

10 Brassica Shoot Elongation Medium MSV-1A

Murashige and Skoog Minimal Organic Medium

Gamborg B5 Vitamins

10 grams sucrose

0.6% agarose

15 pH 5.8

Analysis Of Transgenic Brassica Napus Seeds Containing  
An Antisense Microsomal Delta-12 Desaturase Construct

Fifty-one plants were obtained from transformation with both pZPhCFd2R and pZCCFd2R, 40 were obtained from pZKCFd2R, and 26 from pZNCFd2R. The relative levels of oleate (18:1), linoleate (18:2) and linolate (18:3) change during development so that reliable determination of seed fatty acid phenotype is best obtained from seed which has undergone normal maturation and drydown.

20 Relatively few transformed plants have gone through to maturity, however seeds were sampled from plants which had been transferred to pots for at least 80 days and which had pods that had yellowed and contained seeds with seed coats which had black pigmentation. Plants  
25 were chosen for early analysis based on promoter type, presence and copy number of the inserted delta-12 desaturase antisense gene and fertility of the plant.

30 Fatty acid analysis was done on either individual seeds from transformed and control plants, or on 40 mg  
35 of bulk seed from individual plants as described in

- Example 6. Southern analysis for detection of the presence of canola delta-12 desaturase antisense genes was done on DNA obtained from leaves of transformed plants. DNA was digested either to release the
- 5 promoter:delta-12 desaturase fragment from the transformation vector or to cut outside the coding region of the delta-12 desaturase antisense gene, but within the left and right T-DNA borders of the vector.

TABLE 14

Relative Fatty Acid Profiles of Microsomal Delta-12 Desaturase Antisense Transformed and Control *Brassica Napus* Seeds

PLANT #	PROMOTER	COPY#	AGE*	% of TOTAL FATTY ACIDS				
				16:0	18:0	18:1	18:2	18:3
Westar	control	none	82	4.6	1.2	64.6	20.9	6.6
151-22	phaseolin	>8	82	4.4	1.0	76.6	10.0	6.2
158-8	napin	1	83	3.5	1.5	81.3	6.3	4.6
westar	control	none	106	4.1	1.7	64.4	19.9	7.1
151-22	phaseolin	>8	106	4.2	1.9	74.4	9.9	6.3
151-127	phaseolin	0	106	4.1	2.3	68.4	16.9	5.2
151-268	phaseolin	1	106	4.2	2.7	73.3	12.0	4.2
153-83	conglycinin	2	106	4.1	1.6	68.5	16.7	6.3

\*Seed sampling date in days after the plant was transferred to soil

10

- The expected fatty acid phenotype for antisense suppression of the delta-12 desaturase is decreased relative content of 18:2 with a corresponding increase in 18:1. Plant numbers 151-22 and 158-8 both show a
- 15 substantial decrease in 18:2 content of bulk seed when compared to the westar control at 83 days after planting. Plant 151-22 also shows this difference at maturity in comparison to either the westar control or plant 151-127, which was transformed with the selectable
- 20 marker gene but not the delta-12 desaturase antisense gene.

Since the fatty acid analysis was done on seeds from the primary transformant, individual seed should be segregating for the presence of the transgene copy or copies. The segregating phenotypes serve as an internal control for the effect of the delta-12 desaturase antisense gene. The relative fatty acid phenotypes for 10 individual westar seeds, 10 individual 151-22 seeds and 12 individual 158-8 seeds are given in Table 15 below.

TABLE 15

Relative Fatty Acid Profiles for Individual Seeds of Control and Genetically Segregating Delta-12 Desaturase Transformed Brassica Napus Seeds

<u>westar control</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
4.65	1.05	63.45	21.31	7.29
4.65	1.37	65.41	20.72	6.18
3.86	1.31	62.19	22.50	8.18
4.46	1.41	66.81	19.40	5.63
4.76	1.30	61.90	22.39	7.65
4.59	1.10	64.77	20.62	6.56
4.61	1.16	68.66	18.20	5.07
4.71	1.26	67.28	19.32	5.18
4.67	0.98	61.96	22.93	7.61
4.73	1.33	63.85	21.65	6.23
<u>151-22</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
4.56	1.08	73.40	12.40	7.60
4.25	1.20	77.90	10.00	5.40
4.40	1.00	76.90	10.10	5.90
4.40	0.94	77.40	9.40	6.10
4.50	1.00	73.60	11.30	7.90
4.60	0.98	75.40	10.50	6.50
4.49	0.96	76.70	9.90	6.00
4.20	1.10	77.20	9.70	5.50

0099201-67226960

4.20	1.00	80.00	7.90	4.90
4.50	1.00	78.00	8.80	5.80
<u>158-8</u>				
<u>16:0</u>	<u>18:0</u>	<u>18:1</u>	<u>18:2</u>	<u>18:3</u>
3.62	1.67	84.45	3.60	3.73
3.46	1.64	85.56	3.02	3.36
3.48	1.61	83.64	4.43	4.21
3.53	1.40	83.80	4.41	4.36
3.48	1.39	83.66	4.35	4.44
3.80	1.50	68.17	16.57	7.56
3.41	1.40	83.76	4.38	4.40
3.49	1.29	82.77	5.16	4.60
3.77	1.39	69.47	16.40	6.54
3.44	1.36	83.86	4.49	4.27
3.48	1.38	83.15	4.91	4.53
3.55	1.92	83.69	4.20	3.70

- The westar control shows comparatively little seed to seed variation in content of 18:1 or 18:2. Further the ratio of 18:3/18:2 remains very constant between
- 5 seeds at about 0.35. Plant #158-8 should show a segregation ratio of either 1:2:1 or 1:3 since by Southern analysis it contains a single transgene. The 1:2:1 ratio would indicate a semi-dominant, copy number effect while the 1:3 ratio would indicate complete
- 10 dominance. Two wild type 158-8 segregants are clear in Table 15, while the remaining seeds may either be the same, or the two seeds at greater than 84% 18:1 may represent the homozygous transgeneic. In either case the fatty acid phenotypes of the seeds are as expected
- 15 for effective delta-12 desaturase suppression in this generation. The fatty acid phenotypes of the seeds of plant 151-22 show variation in their 18:1 and 18:2 content, with 18:1 higher than the control average and 18:2 lower. The segregation is apparently quite

complex, as would be expected of a multi-copy transgenic plant.

#### EXAMPLE 8

##### COMBINATION OF HIGH OLEIC TRAITS IN

##### TRANSFORMANT 158-8-1 AND RAPESEED MUTANT IMC 129

*Brassica napus* primary transformant 158-8 that was transformed with a chimeric napin:antisense rapeseed delta-12 desaturase gene (see Example 7 herein) was selfed to give rise to seeds, designated T2 generation, that were segregating for high oleic acid phenotypes. Several high oleic T2 seeds were germinated and leaf DNA of the T2 plants was used in quantitative Southern analysis to identify individuals homozygous for the transformed chimeric gene. One homozygous T2 plant, designated 158-8-1, was selfed and the fatty acid composition of the T3 progeny determined by single seed analysis (Table 16). The data confirmed that 158-8-1 was homozygous for the high oleate trait.

TABLE 16

Fatty Acid Composition of Single Seeds of a Selfed T2 Transgenic Rapeseed Line 158-8-1

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
1	3.91	1.59	82.36	5.75	5.78
2	3.81	1.53	83.34	5.55	5.16
3	3.82	1.55	85.29	4.47	4.27
4	3.67	1.75	83.21	5.05	5.72
5	3.74	1.79	82.64	5.29	5.86
6	3.92	1.63	83.43	5.51	4.91
7	3.82	1.47	84.10	5.10	4.89
8	3.83	1.67	82.63	5.97	5.34
Average	3.82	1.62	83.38	5.34	5.24

T2 plant 158-8-1 was also used as the male parent in a cross with either rapeseed cultivar Westar (control) or IMC 129. Three plants were grown from the F1 hybrid seed from each cross and selfed. Single, segregating F2 seeds from each cross were analyzed for fatty acid composition by imbibing the seeds in water overnight, removing the seed coat and cutting off approximately one-half of the outer cotyledon. Fatty acids in the partial cotyledon were analyzed by direct transesterification in methanol and gas chromatography as described in Example 7.

The combined analysis from 272 F2 seeds from the three 158-8-1 X IMC 129 crosses and of 60 F2 seeds from two 158-8-1 X Westar crosses are shown in Table 17 as % individuals in each class of 18:1 content.

TABLE 17  
Frequency Distribution of F2 seeds of 158-8-1 X IMC 129  
Cross on the Basis of Oleic Acid Content

<u>% 18:1</u> <u>class</u>	<u>% seeds from</u> <u>158-8-1 X Westar cross</u>	<u>% of seeds from</u> <u>158-8-1 X IMC129 cross</u>
61-63	0	0.4
64-66	16.7	1.1
67-69	5	1.1
70-72	3.3	5.1
73-75	0	8.8
76-78	13.3	7.0
79-81	45	7.7
82-84	15	22.1
85-87	1.7	36.8
88-90	0	9.9

The data shows that the average 18:1 content in F2 progeny of the cross involving IMC 129 was significantly higher than that involving Westar.

Thirty F2 half-seeds selected for their very high oleic acid content were germinated and their leaf DNA subjected to molecular analysis (Southern analysis and polymerase chain reaction) with gene specific probes to identify 6 F2 plants that were homozygous for both the mutant delta-12 desaturase gene from IMC 129 and the transgene from 158-8-1. The fatty acid composition of the F2 seeds from which these 6 F2 plants were derived is shown in Table 18 and their average fatty acid compositions is compared to those of 158-8-1 T3 seeds (Table 16) and to those of Westar control seeds (Table 19).

**TABLE 18**

Fatty Acid Composition of Six F2 Seeds Homozygous for the Transgene and High Oleic Acid Mutation in IMC 129

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
1	2.7	1.9	89.0	2.1	3.5
2	3.0	1.4	89.2	2.4	3.6
3	3.0	1.3	88.2	3.0	3.4
4	2.6	1.3	90.9	2.4	2.9
5	3.3	1.1	84.4	4.6	6.3
6	3.2	1.3	88.4	2.65	4.4
Average	2.97	1.38	88.4	2.86	4.0

**TABLE 19**

Seed Fatty Acid Composition of Westar, 158-8-1 and Six Seeds Homozygous for the Transgene from 158-8-1 and High Oleic Acid Mutation from IMC 129.

Progeny	Relative Fatty Acid Content (%)				
	16:0	18:0	18:1	18:2	18:3
Westar	4.20	2.20	63.00	19.90	10.60
158-8-1	3.33	1.63	83.38	5.34	5.24
158-8-1 x IMC129	2.97	1.38	88.40	2.86	4.02

The data in Table 19 shows that the the seed oleic acid content in the double homozygotes resulting from the 158-8-1 X IMC 129 cross are higher than either parent and that this increase is associated with reduced  
5 saturates (16:0+18:0). Furthermore, the double homozygote plants appear normal in their growth and development.

00697379.102600



SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY
- (ii) TITLE OF INVENTION: GENES FOR MICROSOMAL  
FATTY ACID DELTA-12  
DESATURASES AND RELATED  
ENZYMES FROM PLANTS
- (iii) NUMBER OF SEQUENCES: 17
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: E. I. DU PONT DE NEMOURS  
AND COMPANY
  - (B) STREET: 1007 MARKET STREET
  - (C) CITY: WILMINGTON
  - (D) STATE: DELAWARE
  - (E) COUNTRY: U.S.A.
  - (F) ZIP: 19898
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: MacIntosh
  - (C) OPERATING SYSTEM: MacIntosh 6.0
  - (D) SOFTWARE: Microsoft Word 4.0
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: U.S. 07/977,339
  - (B) FILING DATE: 17-NOV-1992
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Siegell, Barbara C.
  - (B) REGISTRATION NUMBER: 30,684
  - (C) REFERENCE/DOCKET NUMBER: BB-1043-B
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (302) 992-4927
  - (B) TELEFAX: (302) 892-7949
  - (C) TELEX: 835420

09697379.102600

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1372 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Arabidopsis thaliana*
- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: p92103
- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 93..1244
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

AGAGAGAGAG ATTCTGCGGA GGAGCTTCTT CTTCGTAGGG TGTTACATCGT TATTAACGTT      60
ATCGCCCCCTA CGTCAGCTCC ATCTCCAGAA AC  ATG GGT GCA GGT GGA AGA ATG      113
                               Met Gly Ala Gly Gly Arg Met
                               1                               5

CCG GTT CCT ACT TCT TCC AAG AAA TCG GAA ACC GAC ACC ACA AAG CGT      161
Pro Val Pro Thr Ser Ser Lys Lys Ser Glu Thr Asp Thr Thr Lys Arg
    10                               15                               20

GTG CCG TGC GAG AAA CCG CCT TTC TCG GTG GGA GAT CTG AAG AAA GCA      209
Val Pro Cys Glu Lys Pro Pro Phe Ser Val Gly Asp Leu Lys Lys Ala
    25                               30                               35

ATC CCG CCG CAT TGT TTC AAA CGC TCA ATC OCT CGC TCT TTC TCC TAC      257
Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr
    40                               45                               50                               55

CTT ATC AGT GAC ATC ATT ATA GCC TCA TGC TTC TAC TAC GTC GCC ACC      305
Leu Ile Ser Asp Ile Ile Ala Ser Cys Phe Tyr Tyr Val Ala Thr
    60                               65                               70

AAT TAC TTC TCT CTC CTC CCT CAG CCT CTC TCT TAC TTG GCT TGG CCA      353
Asn Tyr Phe Ser Leu Leu Pro Gln Pro Leu Ser Tyr Leu Ala Trp Pro
    75                               80                               85

CTC TAT TGG GCC TGT CAA GGC TGT GTC CTA ACT GGT ATC TGG GTC ATA      401
Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Ile Trp Val Ile
    90                               95                               100

```

00697379.102600

GCC CAC GAA TGC GGT CAC CAC GCA TTC AGC GAC TAC CAA TGG CTG GAT	449
Ala His Glu Cys Gly His His Ala Phe Ser Asp Tyr Gln Trp Leu Asp	
105 110 115	
GAC ACA GTT GGT CTT ATC TTC CAT TCC TTC CTC CTC GTC CCT TAC TTC	497
Asp Thr Val Gly Leu Ile Phe His Ser Phe Leu Leu Val Pro Tyr Phe	
120 125 130 135	
TCC TGG AAG TAT AGT CAT CGC CGT CAC CAT TCC AAC ACT GGA TCC CTC	545
Ser Trp Lys Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu	
140 145 150	
GAA AGA GAT GAA GTA TTT GTC CCA AAG CAG AAA TCA GCA ATC AAG TGG	593
Glu Arg Asp Glu Val Phe Val Pro Lys Gln Lys Ser Ala Ile Lys Trp	
155 160 165	
TAC GGG AAA TAC CTC AAC AAC CCT CTT GGA CGC ATC ATG ATG TTA ACC	641
Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ile Met Met Leu Thr	
170 175 180	
GTC CAG TTT GTC CTC GGG TGG CCC TTG TAC TTA GCC TTT AAC GTC TCT	689
Val Gln Phe Val Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser	
185 190 195	
GGC AGA CCG TAT GAC GGG TTC GCT TGC CAT TTC CCC AAC GCT CCC	737
Gly Arg Pro Tyr Asp Gly Phe Ala Cys His Phe Phe Pro Asn Ala Pro	
200 205 210 215	
ATC TAC AAT GAC CGA GAA CGC CTC CAG ATA TAC CTC TCT GAT GCG GGT	785
Ile Tyr Asn Asp Arg Glu Arg Leu Gln Ile Tyr Leu Ser Asp Ala Gly	
220 225 230	
ATT CTA GCC GTC TGT TTT GGT CTT TAC CGT TAC GCT GCT GCA CAA GGG	833
Ile Leu Ala Val Cys Phe Gly Leu Tyr Arg Tyr Ala Ala Ala Gln Gly	
235 240 245	
ATG GCC TCG ATG ATC TGC CTC TAC GGA GTA CCG CTT CTG ATA GTG AAT	881
Met Ala Ser Met Ile Cys Leu Tyr Gly Val Pro Leu Leu Ile Val Asn	
250 255 260	
GCG TTC CTC GTC TTG ATC ACT TAC TTG CAG CAC ACT CAT CCC TCG TTG	929
Ala Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu	
265 270 275	
CCT CAC TAC GAT TCA TCA GAG TGG GAC TGG CTC AGG GGA GCT TTG GCT	977
Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala	
280 285 290 295	
ACC GTA GAC AGA GAC TAC GGA ATC TTG AAC AAG GTG TTC CAC AAC ATT	1025
Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile	
300 305 310	
ACA GAC ACA CAC GTG GCT CAT CAC CTG TTC TCG ACA ATG CCG CAT TAT	1073
Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr	
315 320 325	

09697379.102600

AAC GCA ATG GAA GCT ACA AAG GCG ATA AAG CCA ATT CTG GGA GAC TAT 1121  
 Asn Ala Met Glu Ala Thr Lys Ala Ile Lys Pro Ile Leu Gly Asp Tyr  
 330 335 340  
 TAC CAG TTC GAT GGA ACA CCG TGG TAT GTA GCG ATG TAT AGG GAG GCA 1169  
 Tyr Gln Phe Asp Gly Thr Pro Trp Tyr Val Ala Met Tyr Arg Glu Ala  
 345 350 355  
 AAG GAG TGT ATC TAT GTA GAA CCG GAC AGG GAA GGT GAC AAG AAA GGT 1217  
 Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Glu Gly Asp Lys Lys Gly  
 360 365 370 375  
 GTG TAC TGG TAC AAC AAT AAG TTA TGAGCATGAT GGTGAAGAAA TTGTCGACCT 1271  
 Val Tyr Trp Tyr Asn Asn Lys Leu  
 380  
 TTCTCTTGTC TGTTTGCTT TTGTTAAAGA AGCTATGCTT CGTTTAAATA ATCTTATGT 1331  
 CCATTTTGTT GTGTATGAC ATTTGGCTG CTCATTATGT T 1372

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 383 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Ala Gly Gly Arg Met Pro Val Pro Thr Ser Ser Lys Lys Ser  
 1 5 10 15  
 Glu Thr Asp Thr Thr Lys Arg Val Pro Cys Glu Lys Pro Pro Phe Ser  
 20 25 30  
 Val Gly Asp Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45  
 Ile Pro Arg Ser Phe Ser Tyr Leu Ile Ser Asp Ile Ile Ile Ala Ser  
 50 55 60  
 Cys Phe Tyr Tyr Val Ala Thr Asn Tyr Phe Ser Leu Leu Pro Gln Pro  
 65 70 75 80  
 Leu Ser Tyr Leu Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
 85 90 95  
 Leu Thr Gly Ile Trp Val Ile Ala His Glu Cys Gly His His Ala Phe  
 100 105 110  
 Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
 115 120 125  
 Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
 130 135 140

09697379.10500

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys  
 145 150 155 160  
 Gln Lys Ser Ala Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu  
 165 170 175  
 Gly Arg Ile Met Met Leu Thr Val Gln Phe Val Leu Gly Trp Pro Leu  
 180 185 190  
 Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Phe Ala Cys  
 195 200 205  
 His Phe Phe Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln  
 210 215 220  
 Ile Tyr Leu Ser Asp Ala Gly Ile Leu Ala Val Cys Phe Gly Leu Tyr  
 225 230 235 240  
 Arg Tyr Ala Ala Gln Gly Met Ala Ser Met Ile Cys Leu Tyr Gly  
 245 250 255  
 Val Pro Leu Leu Ile Val Asn Ala Phe Leu Val Leu Ile Thr Tyr Leu  
 260 265 270  
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp  
 275 280 285  
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu  
 290 295 300  
 Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu  
 305 310 315 320  
 Phe Ser Thr Met Pro His Tyr Asn Ala Met Glu Ala Thr Lys Ala Ile  
 325 330 335  
 Lys Pro Ile Leu Gly Asp Tyr Tyr Gln Phe Asp Gly Thr Pro Trp Tyr  
 340 345 350  
 Val Ala Met Tyr Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp  
 355 360 365  
 Arg Glu Gly Asp Lys Lys Gly Val Tyr Trp Tyr Asn Asn Lys Leu  
 370 375 380

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1394 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

09697379.10600

- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:  
(A) ORGANISM: Brassica napus
- (vii) IMMEDIATE SOURCE:  
(B) CLONE: pCF2-165D
- (ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 99..1250

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAGAGGAGAC AGAGACAGAG AGAGAGTTGA GAGAGCTCTC GTAGGTTATC GTATTAACGT	60
AATCTTCAAT CCCCCCTACG TCAGCCAGCT CAAGAAAC ATG GGT GCA GGT GGA	113
Met Gly Ala Gly Gly	
1 5	
AGA ATG CAA GTG TCT CCT CCC TCC AAA AAG TCT GAA ACC GAC AAC ATC	161
Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser Glu Thr Asp Asn Ile	
10 15 20	
AAG CGC GTA CCC TGC GAG ACA CCG CCC TTC ACT GTC GGA GAA CTC AAG	209
Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr Val Gly Glu Leu Lys	
25 30 35	
AAA GCA ATC CCA CCG CAC TGT TTC AAG CGC TCG ATC CCT CGC TCT TTC	257
Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe	
40 45 50	
TCC CAC CTC ATC TGG GAC ATC ATC ATA GCC TCC TGC TTC TAC TAC GTC	305
Ser His Leu Ile Trp Asp Ile Ile Ala Ser Cys Phe Tyr Tyr Val	
55 60 65	
GCC ACC ACT TAC TTC CCT CTC CTC CCT AAC CCT CTC TCC TAC TTC GCC	353
Ala Thr Thr Tyr Phe Pro Leu Leu Pro Asn Pro Leu Ser Tyr Phe Ala	
70 75 80 85	
TGG CCT CTC TAC TGG GCC TGC CAG GGC TGC GTC CTA ACC GGC GTC TGG	401
Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Val Trp	
90 95 100	
GTC ATA GCC CAC GAG TGC GGC CAC GCA GCC TTC AGC GAC TAC CAG TGG	449
Val Ile Ala His Glu Cys Gly His Ala Ala Phe Ser Asp Tyr Gln Trp	
105 110 115	
CTG GAC GAC ACC GTC GGC CTC ATC TTC CAC TCC TTC CTC CTC GTC CCT	497
Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser Phe Leu Leu Val Pro	
120 125 130	

09697379.102600

TAC TTC	TCC TGG	AAG TAC	AGT CAT	CGA CGC	CAC CAT	TCC AAC	ACT GGC	545
Tyr Phe	Ser Trp	Lys Tyr	Ser His	Arg Arg	His His	Ser Asn	Thr Gly	
135			140		145			
TCC CTC	GAG AGA	GAC GAA	GTG TTT	GTC CCA	AGA AGA	AGT CAG	ACA TCA	593
Ser Leu	Glu Arg	Asp Glu	Val Phe	Val Pro	Arg Arg	Ser Gln	Thr Ser	
150		155		160			165	
AGT GGT	ACG GCA	AGT ACC	TCA ACA	ACC TTT	GGA CGC	ACC GTG	ATG TTA	641
Ser Gly	Thr Ala	Ser Thr	Thr Ser	Thr Thr	Gly Arg	Thr Val	Met Leu	
		170			175		180	
ACG GTT	CAG TTC	ACT CTC	GGC TGG	CCT TTG	TAC TTA	GCC TTC	AAC GTC	689
Thr Val	Gln Phe	Thr Leu	Gly Trp	Pro Leu	Tyr Tyr	Leu Ala	Phe Asn	
	185			190			195	
TCG GGG	AGA CCT	TAC GAC	GGC GGC	TTC GCT	TGC CAT	TTT CAC	CCC AAC	737
Ser Gly	Arg Pro	Tyr Asp	Gly Gly	Phe Ala	Cys His	Phe His	Pro Asn	
	200		205			210		
GCT CCC	ATC TAC	AAC GAC	CGT GAG	CGT CTC	CAG ATA	TAC ATC	TCC GAC	785
Ala Pro	Ile Tyr	Asn Asp	Arg Glu	Arg Leu	Gln Ile	Tyr Ile	Ser Asp	
	215		220		225			
GCT GGC	ATC CTC	GCC GTC	TGC TAC	GGT CTG	CTA CCG	TAC GCT	GCT GTC	833
Ala Gly	Ile Leu	Ala Val	Cys Tyr	Gly Leu	Pro Pro	Tyr Ala	Ala Val	
230		235		240			245	
CAA GGA	GTT GCC	TCG ATG	GTC TGC	TTC CTA	CGA GTT	CCT CTT	CTG ATT	881
Gln Gly	Val Ala	Ser Met	Val Cys	Phe Leu	Arg Val	Pro Leu	Leu Ile	
	250			255			260	
GTC AAC	GGG TTC	TTA GTT	TTG ATC	ACT ACT	TAC TTG	CAG CAC	ACG CAT	929
Val Asn	Gly Phe	Leu Val	Leu Ile	Thr Thr	Tyr Leu	Gln His	Thr His	
	265			270			275	
TCC CTG	CCT CAC	TAT GAC	TCG TCT	GAG TGG	GAT TGG	TTG AGG	GGA GCT	977
Ser Leu	Pro His	Tyr Asp	Ser Ser	Glu Trp	Asp Trp	Leu Arg	Gly Ala	
	280		285			290		
TTG GCC	ACC GTT	GAC AGA	GAC TAC	GGA ATC	TTG AAC	CAA GGC	TTC CAC	1025
Leu Ala	Thr Val	Asp Arg	Tyr Tyr	Gly Ile	Leu Leu	Gln Gly	Phe His	
	295		300		305			
AAT ATC	ACG GAC	ACG CAC	GAG GCG	CAT CAC	CTG TTC	TCG ACC	ATG CCG	1073
Asn Ile	Thr Asp	Thr His	Glu Ala	His His	Leu Phe	Ser Thr	Met Pro	
310		315			320		325	
CAT TAT	CAT GCG	ATG GAA	GCT ACG	AAG GCG	ATA AAG	CCG ATA	CTG GGA	1121
His Tyr	His Ala	Met Glu	Ala Thr	Lys Lys	Ala Ile	Lys Pro	Ile Leu	
		330		335			340	
GAG TAT	TAT CAG	TTC GAT	GGG ACG	CCG GTG	GTT AAG	GCG ATG	TGG AGG	1169
Glu Tyr	Tyr Gln	Phe Asp	Gly Thr	Pro Val	Val Val	Lys Ala	Met Trp	
	345			350			355	

09597379-102600

GAG GCG AAG GAG TGT ATC TAT GTG GAA CCG GAC AGG CAA GGT GAG AAG 1217  
 Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Gln Gly Glu Lys  
           360  370

AAA GGT GTG TTC TGG TAC AAC AAT AAG TTA TGAAGCAAAG AAGAACTGA 1267  
 Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
           375  380

ACCTTTCTCT TCTATCAATT GTCTTTGTTT AAGAAGCTAT GTTCTGTGTT CAATAATCTT 1327

AATTATCCAT TTTGTGTGT TTTCTGACAT TTIGGCTAAA ATTATGTGAT GTTGAAGTT 1387

AGTGTCT 1394

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 383 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
   1  5  10  15

Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
   20  25  30

Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
   35  40  45

Ile Pro Arg Ser Phe Ser His Leu Ile Trp Asp Ile Ile Ile Ala Ser  
   50  55  60

Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro Asn Pro  
   65  70  75  80

Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val  
   85  90  95

Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His Ala Ala Phe  
   100  105  110

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser  
   115  120  125

Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His  
   130  135  140

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Arg  
   145  150  155  160

Arg Ser Gln Thr Ser Ser Gly Thr Ala Ser Thr Ser Thr Thr Phe Gly  
   165  170  175

09697379.102600



Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu Tyr  
 180 185  
 Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala Cys  
 195 200 205  
 His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu Gln  
 210 215 220  
 Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu Leu  
 225 230 235 240  
 Pro Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Leu Arg  
 245 250 255  
 Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr Leu  
 260 265 270  
 Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp Asp  
 275 280 285  
 Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu  
 290 295 300  
 Asn Gln Gly Phe His Asn Ile Thr Asp Thr His Glu Ala His His Leu  
 305 310 315 320  
 Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala Ile  
 325 330 335  
 Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val Val  
 340 345 350  
 Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro Asp  
 355 360 365  
 Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu  
 370 375 380

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1462 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Glycine max

09697379.102600

(vii) IMMEDIATE SOURCE:  
(B) CLONE: pSF2-165K

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 108..1247

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCATATACTA ATATTGCTT GTATTGATAG CCCCTCCGTT CCCAAGAGTA TAAAACTGCA	60
TCGAATAATA CAAGCCACTA GGCATGGGTC TAGCAAAGGA AACAAACA ATG GGA GGT	116
Met Gly Gly	
1	
AGA GGT CGT GTG GCC AAA GTG GAA GTT CAA GGG AAG AAG CCT CTC TCA	164
Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys Pro Leu Ser	
5 10 15	
AGG GTT CCA AAC ACA AAG CCA CCA TTC ACT GTT GGC CAA CTC AAG AAA	212
Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln Leu Lys Lys	
20 25 30 35	
GCA ATT CCA CCA CAC TGC TTT CAG CGC TCC CTC CTC ACT TCA TTC TCC	260
Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr Ser Phe Ser	
40 45 50	
TAT GTT GTT TAT GAC CTT TCA TTT GCC TTC ATT TTC TAC ATT GCC ACC	308
Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr Ile Ala Thr	
55 60 65	
ACC TAC TTC CAC CTC CTT CCT CAA CCC TTT TCC CTC ATT GCA TGG CCA	356
Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile Ala Trp Pro	
70 75 80	
ATC TAT TGG GTT CTC CAA GGT TGC CTT CTC ACT GGT GTG TGG GTG ATT	404
Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val Trp Val Ile	
85 90 95	
GCT CAC GAG TGT GGT CAC CAT GCC TTC AGC AAG TAC CAA TGG GTT GAT	452
Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln Trp Val Asp	
100 105 110 115	
GAT GTT GTG GGT TTG ACC CTT CAC TCA ACA CTT TTA GTC CCT TAT TTC	500
Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val Pro Tyr Phe	
120 125 130	
TCA TGG AAA ATA AGC CAT CGC CGC CAT CAC TCC AAC ACA GGT TCC CTT	548
Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu	
135 140 145	
GAC CGT GAT GAA GTG TTT GTC CCA AAA CCA AAA TCC AAA GTT GCA TGG	596
Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys Val Ala Trp	
150 155 160	

00697379.102600

TTT TCC AAG TAC TTA AAC AAC CCT CTA GGA AGG GCT GTT TCT CTT CTC	644
Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val Ser Leu Leu	
165 170 175	
GTC ACA CTC ACA ATA GGG TGG CCT ATG TAT TTA GCC TTC AAT GTC TCT	692
Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe Asn Val Ser	
180 185 190 195	
GGT AGA CCC TAT GAT AGT TTT GCA AGC CAC TAC CAC CCT TAT GCT CCC	740
Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro Tyr Ala Pro	
200 205 210	
ATA TAT TCT AAC CGT GAG AGG CTT CTG ATC TAT GTC TCT GAT GCT GCT	788
Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser Asp Val Ala	
215 220 225	
TTG TTT TCT GTG ACT TAC TCT CTC TAC CGT GTT GCA ACC CTG AAA GGG	836
Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr Leu Lys Gly	
230 235 240	
TTG GTT TGG CTG CTA TGT GTT TAT GGG GTG CCT TTG CTC ATT GTG AAC	884
Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu Ile Val Asn	
245 250 255	
GGT TTT CTT GTG ACT ATC ACA TAT TTG CAG CAC ACA CAC TTT GCC TTG	932
Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His Phe Ala Leu	
260 265 270 275	
CCT CAT TAC GAT TCA TCA GAA TGG GAC TGG CTG AAG GGA GCT TTG GCA	980
Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly Ala Leu Ala	
280 285 290	
ACT ATG GAC AGA GAT TAT GGG ATT CTG AAC AAG GTG TTT CAT CAC ATA	1028
Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His His Ile	
295 300 305	
ACT GAT ACT CAT GTG GCT CAC CAT CTC TTC TCT ACA ATG CCA CAT TAC	1076
Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr	
310 315 320	
CAT GCA ATG GAG GCA ACC AAT GCA ATC AAG CCA ATA TTG GGT GAG TAC	1124
His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu Gly Glu Tyr	
325 330 335	
TAC CAA TTT GAT GAC ACA CCA TTT TAC AAG GCA CTG TGG AGA GAA GCG	1172
Tyr Gln Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala	
340 345 350 355	
AGA GAG TGC CTC TAT GTG GAG CCA GAT GAA GGA ACA TCC GAG AAG GGC	1220
Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser Glu Lys Gly	
360 365 370	
GTG TAT TGG TAC AGG AAC AAG TAT TGATGGAGCA ACCAATGGGC CATAGTGGGA	1274
Val Tyr Trp Tyr Arg Asn Lys Tyr	
375 380	
GGTATGGAAG TTTTGTCTAG TATTAGTACA TAATTAGTAG AATGTTATAA ATAAGTGGAT	1334

00697379-102600

TTGCCGCGTA ATGACTTTGT GTGTATTGTG AACAGCTTG TTGCGATCAT GGTTATAATG 1394  
 TAAAAATAAT TCTGGTATTA ATTACATGTG GAAAGTGTTT TGCTTATAGC TTTCTGCCTA 1454  
 AAAAAAAA 1462

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 379 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Gly Gly Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys  
 1 5 10 15  
 Pro Leu Ser Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln  
 20 25 30  
 Leu Lys Lys Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr  
 35 40 45  
 Ser Phe Ser Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr  
 50 55 60  
 Ile Ala Thr Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile  
 65 70 75 80  
 Ala Trp Pro Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val  
 85 90 95  
 Trp Val Ile Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln  
 100 105 110  
 Trp Val Asp Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val  
 115 120 125  
 Pro Tyr Phe Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr  
 130 135 140  
 Gly Ser Leu Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys  
 145 150 155 160  
 Val Ala Trp Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val  
 165 170 175  
 Ser Leu Leu Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe  
 180 185 190  
 Asn Val Ser Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro  
 195 200 205

09697379.102600

Tyr Ala Pro Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser  
 210 215 220  
 Asp Val Ala Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr  
 225 230 235 240  
 Leu Lys Gly Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu  
 245 250 255  
 Ile Val Asn Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His  
 260 265 270  
 Phe Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly  
 275 280 285  
 Ala Leu Ala Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
 290 295 300  
 His His Ile Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met  
 305 310 315 320  
 Pro His Tyr His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu  
 325 330 335  
 Gly Glu Tyr Tyr Gln Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp  
 340 345 350  
 Arg Glu Ala Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser  
 355 360 365  
 Glu Lys Gly Val Tyr Trp Tyr Arg Asn Lys Tyr  
 370 375

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1790 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Zea mays
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pFad2#1
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 165..1328

09697379.102600

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGCCTCTCC	CCTCCTCTCT	CCCTGCAAAAT	CCTGCAGACA	CCACGCTCG	TTTTTCTCTC	60
CGGGACAGGA	GAAAAGGGGA	GAGAGAGGTG	AGCGCGGTG	TCGCCCCGAT	CTGCTCTGCC	120
CCGACGCAGC	TGTTACGACC	TCCTCAGTCT	CAGTCAGGAG	CAAG	ATG GGT GCC GGC	176
					Met Gly Ala Gly	
					1	
GGC AGG	ATG ACC	GAG AAG	GAG CGG	GAG AAG	CAG GAG CAG CTC	GCC CGA
Gly Arg	Met Thr	Glu Lys	Glu Arg	Glu Lys	Gln Glu Gln Leu	Ala Arg
5		10		15		20
GCT ACC	GGT GGC	GCC GCG	ATG CAG	CGG TCG	GTG GAG AAG	CCT CCG
Ala Thr	Gly Gly	Ala Ala	Met Gln	Arg Ser	Pro Val Glu Lys	Pro Pro
		25		30		35
TTC ACT	CTG GGT	CAG ATC	AAG AAG	GCC ATC	CCG CCA CAC	TGC TTC
Phe Thr	Leu Gly	Gln Ile	Lys Lys	Ala Ile	Pro Pro His	Cys Phe
		40		45		50
CGC TCG	GTG CTC	AAG TCC	TTC TCG	TAC GTG	GTC CAC GAC	CTG GTG
Arg Ser	Val Leu	Lys Ser	Phe Ser	Tyr Val	Val His Asp	Leu Val
		55		60		65
GCC GCG	GCG CTC	CTC TAC	TTC GCG	CTG GCC	ATC ATA CCG	GCG CTC
Ala Ala	Ala Leu	Leu Tyr	Phe Ala	Leu Ala	Ile Ile Pro	Ala Leu
		70		75		80
AGC CCG	CTC GCG	TAC GCC	GCC TGG	CCG CTG	TAC TGG	ATC GCG
Ser Pro	Leu Arg	Tyr Ala	Ala Ala	Trp Pro	Leu Tyr	Trp Ile
		85		90		95
TGC GTG	TGC ACC	GGC GTG	TGG GTC	ATC GCG	CAC GAG	TGC GGC
Cys Val	Cys Thr	Gly Val	Trp Val	Ile Ala	His Glu	Cys Gly
		105		110		115
GCC TTC	TCG GAC	TAC TCG	CTC CTG	GAC GAC	GTG GTC	GGC CTG
Ala Phe	Ser Asp	Tyr Ser	Leu Leu	Asp Asp	Val Val Gly	Leu Val
		120		125		130
CAC TCG	TCG CTC	ATG GTG	CCC TAC	TTC TCG	TGG AAG	TAC AGC
His Ser	Ser Leu	Met Val	Pro Tyr	Phe Ser	Trp Lys	Tyr Ser
		135		140		145
CGC CAC	CAC TCC	AAC ACG	GGG TCC	CTG GAG	CGC GAC	GAG GTG
Arg His	His Ser	Asn Thr	Gly Ser	Leu Glu	Arg Asp	Glu Val
		150		155		160
CCC AAG	AAG AAG	GAG GCG	CTG CCG	TGG TAC	ACC CCG	TAC GTG
Pro Lys	Lys Lys	Glu Ala	Leu Pro	Trp Tyr	Thr Pro	Tyr Val
		165		170		175
						180

009697379.102600

AAC CCG GTC GGC CGG GTG GTG CAC ATC GTG GTG CAG CTC ACC CTC GGG Asn Pro Val Gly Arg Val Val His Ile Val Val Gln Leu Thr Leu Gly 185 190	752
TGG CCG CTG TAC CTG GCG ACC AAC GCG TCG GGG CCG CCG TAC CCG CGC Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg Pro Tyr Pro Arg 200 205 210	800
TTC GCC TGC CAC TTC GAC CCC TAC GGC CCC ATC TAC AAC GAC CCG GAG Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr Asn Asp Arg Glu 215 220 225	848
CGC GCC CAG ATC TTC GTC TCG GAC GCC GGC GTC GTG GCC GTG GCG TTC Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val Ala Val Ala Phe 230 235 240	896
GGG CTG TAC AAG CTG GCG GCG GCG TTC GGG GTC TGG TGG GTG GTG CGC Gly Leu Tyr Lys Leu Ala Ala Phe Gly Val Trp Trp Val Val Arg 245 250 255 260	944
GTG TAC GCC GTG CCG CTG CTG ATC GTG AAC GCG TGG CTG GTG CTC ATC Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp Leu Val Leu Ile 265 270 275	992
ACC TAC CTG CAG CAC ACC CAC CCG TCG CTC CCC CAC TAC GAC TCG AGC Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser 280 285 290	1040
GAG TGG GAC TGG CTG GCG GGC GCG CTG GCC ACC ATG GAC CCG GAC TAC Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met Asp Arg Asp Tyr 295 300 305	1088
GGC ATC CTC AAC CGC GTG TTC CAC AAC ATC ACG GAC ACG CAC GTC GCG Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp Thr His Val Ala 310 315 320	1136
CAC CAC CTC TTC TCC ACC ATG CCG CAC TAC CAC GCC ATG GAG GCC ACC His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr 325 330 335 340	1184
AAG GCG ATC AGG CCC ATC CTC GGC GAC TAC TAC CAC TTC GAC CCG ACC Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His Phe Asp Pro Thr 345 350 355	1232
CCT GTC GCC AAG GCG ACC TGG GCG GAG GCC GGG GAA TGC ATC TAC GTC Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu Cys Ile Tyr Val 360 365 370	1280
GAG CCC GAG GAC CGC AAG GGC GTC TTC TGG TAC AAC AAG AAG TTC TAGCCGCCG Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn Lys Lys Phe 375 380 385	1335
CGCTGCGAGA GCTGAGGACG CTACCGTAGG AATGGGAGCA GAAACCAGGA GGAGGAGACG	1395
GTACTCGCCC CAAAGTCTCC GTCAACCTAT CTAATCGTTA GTCGTCAGTC TTTTAGACGG	1455
GAAGAGAGAT CATTGGGCA CAGAGACGAA GGCTTACTGC AGTGCCATCG CTAGAGCTGC	1515

009697379 106600

CATCAAGTAC AAGTAGGCAA ATTCGTCAAC TTAGTGTGTC CCATGTTGTT TTTCTTAGTC 1575  
 GTCCGCTGCT GTAGGCTTTC CGGCGGCGGT CGTTTGTGTG GTTGGCATCC GTGGCCATGC 1635  
 CTGTGCGTGC GTGGCCGCGC TTGTCGTGTG CGTCTGTCGT CGCCTGGGCG TCGTCTCTTC 1695  
 GTGCTCCCGG TGTGTTGTTG TAAACAAGA AGATGTTTTT TGGTGTCTTT GCGCGAATAA 1755  
 CAGATCGTCC GAACGAAAAA AAAAAAAAAA AAAAA 1790

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Gly Ala Gly Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu  
 1 5 10 15  
 Gln Leu Ala Arg Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val  
 20 25 30  
 Glu Lys Pro Pro Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro  
 35 40 45  
 His Cys Phe Glu Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His  
 50 55 60  
 Asp Leu Val Ile Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile  
 65 70 75 80  
 Pro Ala Leu Pro Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp  
 85 90 95  
 Ile Ala Gln Gly Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu  
 100 105 110  
 Cys Gly His His Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val  
 115 120 125  
 Gly Leu Val Leu His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys  
 130 135 140  
 Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp  
 145 150 155 160  
 Glu Val Phe Val Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro  
 165 170 175  
 Tyr Val Tyr Asn Asn Pro Val Gly Arg Val Val His Ile Val Val Gln  
 180 185 190

09697379.102600



Leu Thr Leu Gly Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg  
 195 200 205  
 Pro Tyr Pro Arg Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr  
 210 215 220  
 Asn Asp Arg Glu Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val  
 225 230 235 240  
 Ala Val Ala Phe Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp  
 245 250 255  
 Trp Val Val Arg Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp  
 260 265 270  
 Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His  
 275 280 285  
 Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met  
 290 295 300  
 Asp Arg Asp Tyr Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp  
 305 310 315 320  
 Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala  
 325 330 335  
 Met Glu Ala Thr Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His  
 340 345 350  
 Phe Asp Pro Thr Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu  
 355 360 365  
 Cys Ile Tyr Val Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn  
 370 375 380  
 Lys Lys Phe  
 385

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 673 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: *Ricinus communis*

009697379.102600

(vii) IMMEDIATE SOURCE:  
(B) CLONE: pRF2-1C

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 1..673

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGG GTG ATG GCG CAT GAT TGT GGG CAC CAT TTC AGT GAC TAT CAA	48
Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln	
1 5 10 15	
TTG CTT GAT GAT GTA GTT GGT CTT ATC CTA CAC TCC TGT CTC CTT GTC	96
Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val	
20 25 30	
CCT TAT TTT TCA TGG AAA CAC AGC CAT CGC CGA CAT CAT TCC AAC ACA	144
Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His Ser Asn Thr	
35 40 45	
GGG TCC CTG GAA CGG GAT GAA GTG TTT GTT CCC AAG AAG AAA TCT AGT	192
Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Ser Ser	
50 55 60	
ATC CGT TGG TAT TCC AAA TAC CTC AAC AAC CCT CCA GGT CGT ATC ATG	240
Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met	
65 70 75 80	
ACA ATT GCC GTC ACA CTT TCA CTT GGC TGG CCT CTG TAC CTA GCA TTC	288
Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe	
85 90 95	
AAT GTT TCA GGC AGG CCA TAT GAT CGG TTC GCC TGC CAC TAT GAC CCA	336
Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro	
100 105 110	
TAT GGC CCG ATC TAC AAT GAT CGC GAG CGA ATC GAG ATA TTC ATA TCA	384
Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser	
115 120 125	
GAT GCT GGT GTT CTT GCT GTC ACT TTT GGT CTC TAC CAA CTT GCT ATA	432
Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile	
130 135 140	
GCG AAG GGG CTT GCT TGG GTT GTC TGT GTA TAT GGA GTG CCA TTG TTG	480
Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu	
145 150 155 160	
GTG GTG AAT TCA TTC CTT GTT CTG ATC ACA TTT CTG CAG CAT ACT CAC	528
Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His	
165 170 175	
CCT GCA TTG CCA CAT TAT GAT TCG TCG GAG TGG GAC TGG CTA AGA GGA	576
Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly	
180 185 190	

09697379-102600

CAT AAC ATA ACG GAC ACT CAA GTA GCT CAC CAC CTT TTC ACC ATG CCC C 673  
His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
210 215 220

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

Trp 1	Val	Met	Ala	His 5	Asp	Cys	Gly	His	His 10	Ala	Phe	Ser	Asp	Tyr 15	Gln
Leu	Leu	Asp	Asp 20	Val	Val	Gly	Leu	Ile 25	Leu	His	Ser	Cys	Leu 30	Leu	Val
Pro	Tyr	Phe 35	Ser	Trp	Lys	His	Ser 40	His	Arg	Arg	His	His 45	Ser	Asn	Thr
Gly	Ser 50	Leu	Glu	Arg	Asp 55	Glu	Val	Phe	Val	Pro	Lys 60	Lys	Lys	Ser	Ser
Ile 65	Arg	Trp	Tyr	Ser 70	Lys	Tyr	Leu	Asn	Asn	Pro 75	Pro	Gly	Arg	Ile	Met 80
Thr	Ile	Ala	Val	Thr 85	Leu	Ser	Leu	Gly	Trp 90	Pro	Leu	Tyr	Leu	Ala 95	Phe
Asn	Val	Ser	Gly 100	Arg	Pro	Tyr	Asp	Arg 105	Phe	Ala	Cys	His	Tyr 110	Asp	Pro
Tyr	Gly	Pro 115	Ile	Tyr	Asn	Asp	Arg 120	Glu	Arg	Ile	Glu	Ile 125	Phe	Ile	Ser
Asp	Ala 130	Gly	Val	Leu	Ala 135	Val	Thr	Phe	Gly	Leu	Tyr 140	Gln	Leu	Ala	Ile
Ala 145	Lys	Gly	Leu	Ala 150	Trp	Val	Val	Cys	Val	Tyr 155	Gly	Val	Pro	Leu	Leu 160
Val	Val	Asn	Ser	Phe 165	Leu	Val	Leu	Ile	Thr 170	Phe	Leu	Gln	His	Thr 175	His
Pro	Ala	Leu	Pro 180	His	Tyr	Asp	Ser	Ser 185	Glu	Trp	Asp	Trp	Leu 190	Arg	Gly
Ala	Leu	Ala 195	Thr	Val	Asp	Arg	Asp 200	Tyr	Gly	Ile	Leu	Asn 205	Lys	Val	Phe

His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1369 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: *Ricinus communis*
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pRF197c-42
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 184..1347
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```
CGGCCGGGAT TCCGGTTTTC AACTAATTT GCAAAAAATG CATGATTCA CCTCAATCA      60
AACACCACAC CTTATAACTT AGTCTTAAGA GAGAGAGAGA GAGGAGACAT TTCTCTTCTC      120
TGAGATGAGC ACTTCTCTTC CAGACATCGA AGCCTCAGGA AAGTGCTTGA GAAGAGCTTG      180
AGA ATG GGA GGT GGT GGT CGC ATG TCT ACT GTC ATA ATC AGC AAC AAC      228
  Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn
    1             5             10             15
AGT GAG AAG AAA GGA GGA AGC AGC CAC CTG GAG CGA GCG CCG CAC ACG      276
Ser Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr
          20             25
AAG CCT CCT TAC ACA CTT GGT AAC CTC AAG AGA GCC ATC CCA CCC CAT      324
Lys Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His
          35             40             45
TGC TTT GAA CGC TCT TTT GTG CGC TCA TTC TCC AAT TTT GCC TAT AAT      372
Cys Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Ser Asn Phe Ala Tyr Asn
          50             55             60
TTC TGC TTA AGT TTT CTT TCC TAC TCG ATC GCC ACC AAC TTC TTC CCT      420
Phe Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro
          65             70             75
```

09697379.102600

TAC	ATC	TCT	TCT	CCG	CTC	TCG	TAT	GTC	GCT	TGG	CTG	GTT	TAC	TGG	CTC	468
Tyr	Ile	Ser	Ser	Pro	Leu	Ser	Tyr	Val	Ala	Trp	Leu	Val	Tyr	Trp	Leu	
80					85					90					95	
TTC	CAA	GGC	TGC	ATT	CTC	ACT	GGT	CTT	TGG	GTC	ATC	GGC	CAT	GAA	TGT	516
Phe	Gln	Gly	Cys	Ile	Leu	Thr	Gly	Leu	Trp	Val	Ile	Gly	His	Glu	Cys	
				100					105					110		
GGC	CAT	CAT	GCT	TTT	AGT	GAG	TAT	CAG	CTG	GCT	GAT	GAC	ATT	GTT	GGC	564
Gly	His	His	Ala	Phe	Ser	Glu	Tyr	Gln	Leu	Ala	Asp	Asp	Ile	Val	Gly	
			115					120					125			
CTA	ATT	GTC	CAT	TCT	GCA	CTT	CTG	GTT	CCA	TAT	TTT	TCA	TGG	AAA	TAT	612
Leu	Ile	Val	His	Ser	Ala	Leu	Leu	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Tyr	
		130					135					140				
AGC	CAT	CGC	CGC	CAC	CAT	TCT	AAC	ATA	GGA	TCT	CTC	GAG	CGA	GAC	GAA	660
Ser	His	Arg	Arg	His	Ser	Asn	Ile	Gly	Ser			Glu	Arg	Asp	Glu	
		145				150					155					
GTG	TTC	GTC	CCG	AAA	TCA	AAG	TCG	AAA	ATT	TCA	TGG	TAT	TCT	AAG	TAC	708
Val	Phe	Val	Pro	Lys	Ser	Lys	Ser	Lys	Ile	Ser	Trp	Tyr	Ser	Lys	Tyr	
160					165					170					175	
TTA	AAC	AAC	CCG	CCA	GGT	CGA	GTT	TTG	ACA	CTT	GCT	GCC	ACG	CTC	CTC	756
Leu	Asn	Asn	Pro	Pro	Gly	Arg	Val	Leu	Thr	Leu	Ala	Ala	Thr	Leu	Leu	
				180					185					190		
CTT	GGC	TGG	CCT	TTA	TAT	TTA	GCT	TTC	AAT	GTC	TCT	GGT	AGA	CCT	TAC	804
Leu	Gly	Trp	Pro	Leu	Tyr	Leu	Ala	Phe	Asn	Val	Ser	Gly	Arg	Pro	Tyr	
			195					200					205			
GAT	CGC	TTT	GCT	TGC	CAT	TAT	GAT	CCC	TAT	GGC	CCA	ATA	TTT	TCC	GAA	852
Asp	Arg	Phe	Ala	Cys	His	Tyr	Asp	Pro	Tyr	Gly	Pro	Ile	Phe	Ser	Glu	
		210					215					220				
AGA	GAA	AGG	CTT	CAG	ATT	TAC	ATT	GCT	GAC	CTC	GGA	ATC	TTT	GCC	ACA	900
Arg	Glu	Arg	Leu	Gln	Ile	Tyr	Ile	Ala	Asp	Leu	Gly	Ile	Phe	Ala	Thr	
		225				230					235					
ACG	TTT	GTG	CTT	TAT	CAG	GCT	ACA	ATG	GCA	AAA	GGG	TTG	GCT	TGG	GTA	948
Thr	Phe	Val	Leu	Tyr	Gln	Ala	Thr	Met	Ala	Lys	Gly	Leu	Ala	Trp	Val	
240					245					250					255	
ATG	CGT	ATC	TAT	GGG	GTG	CCA	TTG	CTT	ATT	GTT	AAC	TGT	TTC	CTT	GTT	996
Met	Arg	Ile	Tyr	Gly	Val	Pro	Leu	Leu	Ile	Val	Asn	Cys	Phe	Leu	Val	
				260					265					270		
ATG	ATC	ACA	TAC	TTG	CAG	CAC	ACT	CAC	CCA	GCT	ATT	CCA	CGC	TAT	GGC	1044
Met	Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Pro	Ala	Ile	Pro	Arg	Tyr	Gly	
			275					280					285			
TCA	TCG	GAA	TGG	GAT	TGG	CTC	CGG	GGA	GCA	ATG	GTG	ACT	GTC	GAT	AGA	1092
Ser	Ser	Glu	Trp	Asp	Trp	Leu	Arg	Gly	Ala	Met	Val	Thr	Val	Asp	Arg	
		290					295					300				

00697379 102600

GAT	TAT	GGG	GTG	TTG	AAT	AAA	GTA	TTC	CAT	AAC	ATT	GCA	GAC	ACT	CAT	1140
Asp	Tyr	Gly	Val	Leu	Asn	Lys	Val	Phe	His	Asn	Ile	Ala	Asp	Thr	His	
305					310					315						
GTA	GCT	CAT	CAT	CTC	TTT	GCT	ACA	GTG	CCA	CAT	TAC	CAT	GCA	ATG	GAG	1188
Val	Ala	His	His	Leu	Phe	Ala	Thr	Val	Pro	His	Tyr	His	Ala	Met	Glu	
320				325					330					335		
GCC	ACT	AAA	GCA	ATC	AAG	CCT	ATA	ATG	GGT	GAG	TAT	TAC	CGG	TAT	GAT	1236
Ala	Thr	Lys	Ala	Ile	Lys	Pro	Ile	Met	Gly	Glu	Tyr	Tyr	Arg	Tyr	Asp	
			340					345						350		
GGT	ACC	CCA	TTT	TAC	AAG	GCA	TTG	TGG	AGG	GAG	GCA	AAG	GAG	TGC	TTG	1284
Gly	Thr	Pro	Phe	Tyr	Lys	Ala	Leu	Trp	Arg	Glu	Ala	Lys	Glu	Cys	Leu	
		355					360						365			
TTC	GTC	GAG	CCA	GAT	GAA	GGA	GCT	CCT	ACA	CAA	GGC	GTT	TTC	TGG	TAC	1332
Phe	Val	Glu	Pro	Asp	Glu	Gly	Ala	Pro	Thr	Gln	Gly	Val	Phe	Trp	Tyr	
		370				375						380				
CGG	AAC	AAG	TAT	TAAAAAAGTG	TCATGTAGCC	TGCCG										1369
Arg	Asn	Lys	Tyr													
385																

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met	Gly	Gly	Gly	Gly	Arg	Met	Ser	Thr	Val	Ile	Ile	Ser	Asn	Asn	Ser	
1				5					10					15		
Glu	Lys	Lys	Gly	Gly	Ser	Ser	His	Leu	Glu	Arg	Ala	Pro	His	Thr	Lys	
			20					25					30			
Pro	Pro	Tyr	Thr	Leu	Gly	Asn	Leu	Lys	Arg	Ala	Ile	Pro	Pro	His	Cys	
		35				40						45				
Phe	Glu	Arg	Ser	Phe	Val	Arg	Ser	Phe	Ser	Asn	Phe	Ala	Tyr	Asn	Phe	
	50					55				60						
Cys	Leu	Ser	Phe	Leu	Ser	Tyr	Ser	Ile	Ala	Thr	Asn	Phe	Phe	Pro	Tyr	
	65				70					75				80		
Ile	Ser	Ser	Pro	Leu	Ser	Tyr	Val	Ala	Trp	Leu	Val	Tyr	Trp	Leu	Phe	
			85					90						95		
Gln	Gly	Cys	Ile	Leu	Thr	Gly	Leu	Trp	Val	Ile	Gly	His	Glu	Cys	Gly	
			100					105					110			

00697379.102600

His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly Leu  
 115 120  
 Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser  
 130 135 140  
 His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu Val  
 145 150 155 160  
 Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr Leu  
 165 170 175  
 Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu Leu  
 180 185 190  
 Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp  
 195 200 205  
 Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu Arg  
 210 215 220  
 Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr Thr  
 225 230 235 240  
 Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val Met  
 245 250 255  
 Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val Met  
 260 265 270  
 Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly Ser  
 275 280 285  
 Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg Asp  
 290 295 300  
 Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His Val  
 305 310 315 320  
 Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu Ala  
 325 330 335  
 Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp Gly  
 340 345 350  
 Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu Phe  
 355 360 365  
 Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr Arg  
 370 375 380  
 Asn Lys Tyr  
 385

09697379.102600

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
  - (A) NAME/KEY: misc\_feature
  - (B) LOCATION: 1..23
  - (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGGTATGCC AYGANTGYGG NCA

23

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 22 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (ix) FEATURE:
  - (A) NAME/KEY: misc\_feature
  - (B) LOCATION: 1..22
  - (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAARTGRTGG CACRTGNGTR TC

22

009201"62526960



## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 2973 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: *Arabidopsis thaliana*
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pAGF2-6
- (ix) FEATURE:
  - (A) NAME/KEY: exon
  - (B) LOCATION: 433..520
- (ix) FEATURE:
  - (A) NAME/KEY: intron
  - (B) LOCATION: 521..1654
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

```

ATTCGGTAAT TCCTACATAT TTTAGAGATT AGTTTGAGTT TCCATCCATA CTTTACTAGT    60
GATTATAAAT TTTAAATACG TACTTTTCGA CTATAAAGTG AAACATAAGTA AATTAGAACG    120
TGATATTAAA AAGTTAATGT TCACGTGTAT ATTTTTTTC AAGTAAAAA ATGGGTATT    180
TGCGGTAAT AAAAATACCA GATATTTTGA ATTGATTAAA AAGGTTGAAA TAAGAGAGGA    240
GGGGAAGAA AAGAAGGTGG GGGCCAGTA TGAAAGGGAA AGGTGTCATC AAATCATCTC    300
TCTCTCTCTC TACCTTCGAC CCACGGGCCG TGTCCATTTA AAGCCCTGTC TCTTGCCATT    360
CCCCATCTGA CCACCAGAAG AAGAGCCACA CACTCACAAA TAAAAAGAG AGAGAGAGAG    420
AGAGAGACAG AGAGAGAGAG AGATTCTGCG GAGGAGCTTC TTCTTCGTAG GGTGTTTCATC    480
GTTATTAAAC TTATCGCCCC TACGTCAGCT CCATCTCCAG GTCCGTCGCT TCTCTTCCAT    540
TTCTTCTCAT TTTGATTTT GATTCTTATT TCTTCCAGT AGCTCCTGCT CTGTGAATTT    600
CTCCGCTCAC GATAGATCTG CTTATATCC TTACATTCAA CCTTAGATCT GGTCTCGATT    660
CTCTGTTTCT CTGTTTTTTT CTITTGGTCG AGAATCTGAT GTTTGTTTAT GTTCTGTCAC    720
CATTAATAAT GATGAACCTC CTCATTCATA CAATGATTAG TTTCTCTCGT CTACCAAACG    780

```

09697379.102600

ATATGTTGCA	TTTTCACTTT	TCTCTTTTT	TTCTAAGATG	ATTGCTTTG	ACCAATTTGT	840
TTAGATCTTT	ATTTTATTTT	ATTTTCTGGT	GGGTTGGTGG	AAATTGAAAA	AAAAAAAAAA	900
AAAAGCATAA	ATTGTTATTT	GTTAATGTAT	TCATTTTTTG	GCTATTGTGT	CTGGGTAAAA	960
ATCTGCTTCT	ACTGTTGAAT	CTTTCCTGGA	TTTTTTACTC	CTATTGGGTT	TTTATAGTAA	1020
AAATACATAA	AAAAAGGAAA	ACAAAAGTTT	TATAGATTCT	CTTAAACCCC	TTACGATAAA	1080
AGTTGGAATC	AAAATAATTC	AGGATCAGAT	GCTCTTTGAT	TGATTACAGT	GCATTACAG	1140
TTGCATGGAA	AATTTTCTAG	ATCCGTCGTC	ACATTTTATT	TTCTGTTTAA	ATATCTAAAT	1200
CTGATATATG	ATGTCGACAA	ATTCTGGTGG	CTTATACATC	ACTTCAACTG	TTTTCTTTTG	1260
GCTTTGTTTG	TCAACTTGGT	TTTCAATACG	ATTTGTGATT	TCGATCGCTG	AATTTTTTAA	1320
ACAAGCAAAC	TGATGTTTAA	CACAAGCAAG	AGATGTGACC	TGCCTTATTA	ACATCGTATT	1380
ACTTACTACT	AGTCGTATT	TCAACGCAAT	CGTTTTTGTA	TTTCTCACAT	TATGCCGCTT	1440
CTCTACTCTT	TATTCCTTTT	GGTCCACGCA	TTTTCTATTT	GTGGCAATCC	CTTTCACAA	1500
CTGATTTCCC	ACTTTGGATC	ATTTGTCTGA	AGACTCTCTT	GAATCGTTAC	CACCTGTTTC	1560
TTGTGCATGC	TCTGTTTTTT	AGAATTAATG	ATAAACTAT	TCCATAGTCT	TGAGTTTTCA	1620
GCTTGTTGAT	TCTTTTGCTT	TTGGTTTTCT	GCAGAAACAT	GGGTGCAGGT	GGAAGAATGC	1680
CGGTTCCCTAC	TTCTTCCAAG	AAATCGGAAA	CCGACACCAC	AAAGCGTGTG	CCGTGCGAGA	1740
AACCGCCTTT	CTCGGTGGGA	GATCTGAAGA	AAGCAATCCC	GCGCATTTGT	TTCAAACGCT	1800
CAATCCCTCG	CTCTTCTCC	TACCTTATCA	GTGACATCAT	TATAGCCTCA	TGCTTCTACT	1860
ACGTGCGCAC	CAATTACTTC	TCTCTCCTCC	CTCAGCCTCT	CTCTTACTTG	GCTTGGCCAC	1920
TCTATTGGGC	CTGCAAGGC	TGTGTCCTAA	CTGGTATCTG	GGTCATAGCC	CACGAATGCG	1980
GTCACCACGC	ATTCAGCGAC	TACCAATGGC	TGGATGACAC	AGTTGGTCTT	ATCTTCCATT	2040
CCTTCCCTCT	CGTCCCTTAC	TTCTCCTGGA	AGTATAGTCA	TCGCCCTCAC	CATTCCAACA	2100
CTGATCCCT	CGAAAGAGAT	GAAGTATTTG	TCCCAAAGCA	GAAATCAGCA	ATCAAGTGGT	2160
ACGGGAAATA	CCTCAACAAC	CCTCTTGGAC	GCATCATGAT	GTTAACCCTG	CAGTTTGTCC	2220
TCGGGTGGCC	CTTGACTTTA	GCCTTTAAGC	TCTCTGGCAG	ACCGTATGAC	GGGTTTCGCTT	2280
GCCATTTCTT	CCCCAACGCT	CCCATCTACA	ATGACCGAGA	ACGCCTCCAG	ATATACCTCT	2340
CTGATCGCGG	TATTCTAGCC	GTCTGTTTTG	GTCTTTACCG	TTACGCTGCT	GCACAAGGGA	2400
TGGCCTCGAT	GATCTGCCTC	TACGGAGTAC	CGCTTCTGAT	AGTGAATGCG	TTCCTCGTCT	2460
TGATCACTTA	CTTGCGAGAC	ACTCATCCCT	CGTTCCTCA	CTACGATTCA	TCAGAGTGGG	2520

ACTGGCTCAG GGGAGCTTTG GCTACCGTAG ACAGAGACTA CGGAATCTTG AACAAAGGTG 2580  
 TCCACAACAT TACAGACACA CACGTGGCTC ATCACCTGTT CTCGACAATG CCGCATTATA 2640  
 ACGCAATGGA AGCTACAAAG GCGATAAAGC CAATTCTGGG AGACTATTAC CAGTTCGATG 2700  
 GAACACCGTG GTATGTGGCG ATGTATAGGG AGGCAAAGGA GTGTATCTAT GTAGAACCGG 2760  
 ACAGGGAAGG TGACAAGAAA GGTGTGTACT GGTACAACAA TAAGTTATGA GGATGATGGT 2820  
 GAAGAAATTG TCGACTTTTC TCTGTCTGT TTGTCITTTG TTAAAGAAGC TATGCTTCGT 2880  
 TTTAATAATC TTATTGTCCA TTTTGTGTG TTATGACATT TTGGCTGCTC ATTATGTTAT 2940  
 GTGGGAAGTT AGCGTTCAAA TGTTTTGGGT CGG 2973

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 1..23
- (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGGCATGTNG ARAANARRTG RTG 23

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

009697379.102600

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc feature

(B) LOCATION: 1..23

(D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGGCATGTRC TRAAANARRTG RTG

23

09697379.102600

CLAIMS

1. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a fatty acid desaturase or a fatty acid desaturase-related enzyme with an amino acid identity of 50% or greater to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
2. The isolated nucleic acid fragment of Claim 1 wherein the amino acid identity is 60% or greater to the polypeptide encoded by SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
3. The isolated nucleic acid fragment of Claim 1 wherein the nucleic acid identity is 90% or greater to SEQ ID NOS:1, 3, 5, 7, 9, 11, or 15.
4. The isolated nucleic acid fragment of Claim 1 wherein said fragment is isolated from an oil-producing plant species.
5. An isolated nucleic acid fragment comprising a nucleic acid sequence encoding a delta-12 fatty acid hydroxylase.
6. A chimeric gene capable of causing altered levels of ricinoleic acid in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of Claim 5, said fragment operably linked to suitable regulatory sequences.
7. A chimeric gene capable of causing altered levels of fatty acids in a transformed plant cell, said chimeric gene comprising a nucleic acid fragment of any of Claims 1, 2, 3, said fragment operably linked to suitable regulatory sequences.
8. Plants containing a chimeric gene of Claim 6 or Claim 7.
9. Oil obtained from seeds of the plants of Claim 8.
10. A method of producing seed oil containing altered levels of unsaturated fatty acids comprising:

09697379.102600

- (a) transforming a plant cell of an oil-producing species with a chimeric gene of Claim 5;
- (b) growing fertile plants from the transformed plant cells of step (a);
- 5 (c) screening progeny seeds from the fertile plants of step (b) for the desired levels of unsaturated fatty acids; and
- (d) processing the progeny seed of step (c) to obtain seed oil containing altered levels of
- 10 unsaturated fatty acids.
11. A method of molecular breeding to obtain altered levels of a fatty acid in seed oil of oil-producing plant species comprising:
- (a) making a cross between two varieites of
- 15 oil-producing species differing in the fatty acid trait;
- (b) making a Southern blot of restriction enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and
- (c) hybridizing the Southern blot with the
- 20 radiolabelled nucleic acid fragment of Claim 1.
12. A method of RFLP mapping comprising:
- (a) making a cross between two varieties of plants;
- (b) making a Southern blot of restriction
- 25 enzyme digested genomic DNA isolated from several progeny plants resulting from the cross of step (a); and
- (c) hybridizing the Southern blot with the radiolabelled nucleic acid fragments of Claim 1.
13. A method to isolate nucleic acid fragments
- 30 encoding fatty acid desaturases and related enzymes, comprising:
- (a) comparing SEQ ID NOS:2, 4, 6, 8, 10, or 12 and other fatty acid desaturase polypeptide sequences;

(b) identifying the conserved sequences of 4 or more amino acids obtained in step a;

(c) designing degenerate oligomers based on the conserved sequences identified in step b; and

- 5 (d) using the degenerate oligomers of step c to isolate sequences encoding fatty acid desaturases and desaturase-related enzymes by sequence dependent protocols.

14. An isolated nucleic acid fragment of Claim 1 comprising a nucleic acid sequence encoding a plant microsomal delta-12 fatty acid desaturase.

15. A method for altering fatty acids composition in seeds comprising:

- (a) making a cross between a mutant line with altered fatty acid composition with a plant containing the chimeric gene of Claim 7;

(b) growing fertile plants from seeds obtained from the cross; and

- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing altered fatty acid levels.

16. A method for reducing polyunsaturated fatty acids in rapeseed oil comprising:

- (a) making a cross between a rapeseed variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 7;

(b) growing fertile plants from seeds obtained from the cross; and

- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

17. The method of Claim 16 wherein the cross in

- (a) is between a progeny plant derived from a seed comprising the Brassica variety having an oleic acid

00697379-102600

content of about 69% to 77%, based upon total extractable oil and belonging to a line in which the said oleic acid content has been stabilized for both the generation to which the seed belongs and its parent generation.

5

18. A method for reducing saturated fatty acids in rapeseed seeds comprising:

- (a) making a cross between a rapeseed variety with increased oleic acid content with a plant containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

15

19. A method for reducing polyunsaturated fatty acids in soybean oil comprising:

- (a) making a cross between a soybean variety with increased oleic acid content or reduced linolenic acid content with a plant containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and
- (c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced polyunsaturated fatty acids.

25

20. A method for reducing saturated fatty acids in soybean seeds comprising:

- (a) making a cross between a soybean variety with increased oleic acid content with a plant containing the chimeric gene of Claim 7;
- (b) growing fertile plants from seeds obtained from the cross; and

30

009201-6226960



(c) screening progeny seeds from the fertile plants of step (b) for seeds containing reduced saturated fatty acids.

21. A *Brassica* sp. plant with seed palmitic acid  
5 of 2.7% or lower of total fatty acid.

22. A *Brassica* sp. plant with seed stearic acid of 1.1% or lower of total fatty acid.

23. A *Brassica* sp. plant with a combined seed palmitic acid and stearic acid content of 3.9% or lower  
10 of total fatty acids.

24. A soybean plant with seed palmitic acid of 6.7% or lower of total fatty acid.

25. A soybean plant with seed stearic acid of 2.1% or lower of total fatty acid.

15 26. A soybean plant with a combined seed palmitic acid and stearic acid content of 9.2% or lower total of fatty acids.

27. Oil obtained from the plants of Claims 21-26.

24 21

00697379.102600

TITLEGENES FOR MICROSOMAL DELTA-12 FATTY ACID  
DESATURASES AND RELATED ENZYMES FROM PLANTSABSTRACT OF THE DISCLOSURE

5       The preparation and use of nucleic acid fragments  
encoding fatty acid desaturase enzymes are described.  
The invention permits alteration of plant lipid  
composition. Chimeric genes incorporating such nucleic  
acid fragments with suitable regulatory sequences may be  
10       used to create transgenic plants with altered levels of  
unsaturated fatty acids.

15

20

25

30

35   BCS/bjm

00697379.102600

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: JONATHAN EDWARD LIGHTNER  
JOHN JOSEPH OKULEY

(ii) TITLE OF INVENTION: GENES FOR MICROSOMAL  
FATTY ACID DELTA-12  
DESATURASES AND RELATED  
ENZYMES FROM PLANTS

(iii) NUMBER OF SEQUENCES: 17

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: E. I. DU PONT DE NEMOURS  
AND COMPANY

(B) STREET: 1007 MARKET STREET

(C) CITY: WILMINGTON

(D) STATE: DELAWARE

(E) COUNTRY: U.S.A.

(F) ZIP: 19898

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: MacIntosh

(C) OPERATING SYSTEM: Microsoft Windows 95

(D) SOFTWARE: Microsoft Word Version 7.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: U.S. 07/977,339

(B) FILING DATE: 17-NOV-1992

(A) APPLICATION NUMBER: U.S. 09/133,962

(B) FILING DATE: 14-AUG-1998

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Christenbury, Lynne M..

(B) REGISTRATION NUMBER: 30,971

(C) REFERENCE/DOCKET NUMBER: BB1043 US DIV

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (302) 992-5481

(B) TELEFAX: (302) 892-1026

(C) TELEX: 835420

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1372 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Arabidopsis thaliana*

(vii) IMMEDIATE SOURCE:  
 (B) CLONE: p92103

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 93..1244

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AGAGAGAGAG ATTCTGCGGA GGAGCTTCTT CTCGCTAGGG TGTCATCGT TATTAACGTT  
 60

ATCGCCCCTA CGTCAGCTCC ATCTCCAGAA AC ATG GGT GCA GGT GGA AGA ATG  
 113

Met Gly Ala Gly Gly Arg Met  
 1 5

CCG GTT CCT ACT TCT TCC AAG AAA TCG GAA ACC GAC ACC ACA AAG CGT  
 161

Pro Val Pro Thr Ser Ser Lys Lys Ser Glu Thr Asp Thr Thr Lys Arg  
 10 15 20

GTG CCG TGC GAG AAA CCG CCT TTC TCG GTG GGA GAT CTG AAG AAA GCA  
 209

Val Pro Cys Glu Lys Pro Pro Phe Ser Val Gly Asp Leu Lys Lys Ala  
 25 30 35

ATC CCG CCG CAT TGT TTC AAA CGC TCA ATC CCT CGC TCT TTC TCC TAC  
 257

Ile Pro Pro His Cys Phe Lys Arg Ser Ile Pro Arg Ser Phe Ser Tyr  
 40 45 50 55

CTT ATC AGT GAC ATC ATT ATA GCC TCA TGC TTC TAC TAC GTC GCC ACC  
 305

Leu Ile Ser Asp Ile Ile Ile Ala Ser Cys Phe Tyr Tyr Val Ala Thr  
 60 65 70

AAT TAC TTC TCT CTC CTC CCT CAG CCT CTC TCT TAC TTG GCT TGG CCA  
 353

Asn Tyr Phe Ser Leu Leu Pro Gln Pro Leu Ser Tyr Leu Ala Trp Pro  
 75 80 85

CTC TAT TGG GCC TGT CAA GGC TGT GTC CTA ACT GGT ATC TGG GTC ATA  
 401

```

Leu Tyr Trp Ala Cys Gln Gly Cys Val Leu Thr Gly Ile Trp Val Ile
    90                      95                      100

GCC CAC GAA TGC GGT CAC CAC GCA TTC AGC GAC TAC CAA TGG CTG GAT
449
Ala His Glu Cys Gly His His Ala Phe Ser Asp Tyr Gln Trp Leu Asp
105                      110                      115

GAC ACA GTT GGT CTT ATC TTC CAT TCC TTC CTC CTC GTC CCT TAC TTC
497
Asp Thr Val Gly Leu Ile Phe His Ser Phe Leu Leu Val Pro Tyr Phe
120                      125                      130                      135

TCC TGG AAG TAT AGT CAT CGC CGT CAC CAT TCC AAC ACT GGA TCC CTC
545
Ser Trp Lys Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu
140                      145                      150

GAA AGA GAT GAA GTA TTT GTC CCA AAG CAG AAA TCA GCA ATC AAG TGG
593
Glu Arg Asp Glu Val Phe Val Pro Lys Gln Lys Ser Ala Ile Lys Trp
155                      160                      165

TAC GGG AAA TAC CTC AAC AAC CCT CTT GGA CGC ATC ATG ATG TTA ACC
641
Tyr Gly Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ile Met Met Leu Thr
170                      175                      180

GTC CAG TTT GTC CTC GGG TGG CCC TTG TAC TTA GCC TTT AAC GTC TCT
689
Val Gln Phe Val Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser
185                      190                      195

GGC AGA CCG TAT GAC GGG TTC GCT TGC CAT TTC TTC CCC AAC GCT CCC
737
Gly Arg Pro Tyr Asp Gly Phe Ala Cys His Phe Phe Pro Asn Ala Pro
200                      205                      210                      215

ATC TAC AAT GAC CGA GAA CGC CTC CAG ATA TAC CTC TCT GAT GCG GGT
785
Ile Tyr Asn Asp Arg Glu Arg Leu Gln Ile Tyr Leu Ser Asp Ala Gly
220                      225                      230

ATT CTA GCC GTC TGT TTT GGT CTT TAC CGT TAC GCT GCT GCA CAA GGG
833
Ile Leu Ala Val Cys Phe Gly Leu Tyr Arg Tyr Ala Ala Ala Gln Gly
235                      240                      245

ATG GCC TCG ATG ATC TGC CTC TAC GGA GTA CCG CTT CTG ATA GTG AAT
881
Met Ala Ser Met Ile Cys Leu Tyr Gly Val Pro Leu Leu Ile Val Asn
250                      255                      260

GCG TTC CTC GTC TTG ATC ACT TAC TTG CAG CAC ACT CAT CCC TCG TTG
929
Ala Phe Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu

```

```

265                270                275
CCT CAC TAC GAT TCA TCA GAG TGG GAC TGG CTC AGG GGA GCT TTG GCT
977
Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala
280                285                290                295

ACC GTA GAC AGA GAC TAC GGA ATC TTG AAC AAG GTG TTC CAC AAC ATT
1025
Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile
300                305                310

ACA GAC ACA CAC GTG GCT CAT CAC CTG TTC TCG ACA ATG CCG CAT TAT
1073
Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr
315                320                325

AAC GCA ATG GAA GCT ACA AAG GCG ATA AAG CCA ATT CTG GGA GAC TAT
1121
Asn Ala Met Glu Ala Thr Lys Ala Ile Lys Pro Ile Leu Gly Asp Tyr
330                335                340

TAC CAG TTC GAT GGA ACA CCG TGG TAT GTA GCG ATG TAT AGG GAG GCA
1169
Tyr Gln Phe Asp Gly Thr Pro Trp Tyr Val Ala Met Tyr Arg Glu Ala
345                350                355

AAG GAG TGT ATC TAT GTA GAA CCG GAC AGG GAA GGT GAC AAG AAA GGT
1217
Lys Glu Cys Ile Tyr Val Glu Pro Asp Arg Glu Gly Asp Lys Lys Gly
360                365                370                375

GTG TAC TGG TAC AAC AAT AAG TTA TGAGCATGAT GGTGAAGAAA TTGTCGACCT
1271
Val Tyr Trp Tyr Asn Asn Lys Leu
380

TTCTCTTGTC TGT TTGCTT TTGTTAAAGA AGCTATGCTT CGTTTAAATA ATCTTATTGT
1331

CCATTTTGTG GTGTTATGAC ATTTTGGCTG CTCATTATGT T
1372

```

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 383 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Gly Ala Gly Gly Arg Met Pro Val Pro Thr Ser Ser Lys Lys Ser
1          5          10          15

```



(2) INFORMATION FOR SEO ID NO:3:

(A) LENGTH: 1426 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: sing

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Brassica napus*

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 130..1284

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Page 6



Ile	Ala	Ser	Cys	Phe	Tyr	Tyr	Val	Ala	Thr	Thr	Tyr	Phe	Pro	Leu	Leu		
			65					70					75				
CCT	CAC	CCT	CTC	TCC	TAC	TTC	GCC	TGG	CCT	CTC	TAC	TGG	GCC	TGC	CAG	408	
Pro	His	Pro	Leu	Ser	Tyr	Phe	Ala	Trp	Pro	Leu	Tyr	Trp	Ala	Cys	Gln		
		80					85					90					
GGC	TGC	GTC	CTA	ACC	GGC	GTC	TGG	GTC	ATA	GCC	CAC	GAG	TGC	GGC	CAC	456	
Gly	Cys	Val	Leu	Thr	Gly	Val	Trp	Val	Ile	Ala	His	Glu	Cys	Gly	His		
	95					100					105						
CAC	GCC	TTC	AGC	GAC	TAC	CAG	TGG	CTG	GAC	GAC	ACC	GTC	GGC	CTC	ATC	504	
His	Ala	Phe	Ser	Asp	Tyr	Gln	Trp	Leu	Asp	Asp	Thr	Val	Gly	Leu	Ile		
110					115					120					125		
TTC	CAC	TCC	TTC	CTC	CTC	GTC	CCT	TAC	TTC	TCC	TGG	AAG	TAC	AGT	CAT	552	
Phe	His	Ser	Phe	Leu	Leu	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Tyr	His	His		
				130					135					140			
CGA	CGC	CAC	CAT	TCC	AAC	ACT	GGC	TCC	CTC	GAG	AGA	GAC	GAA	GTG	TTT	600	
Arg	Arg	His	His	Ser	Asn	Thr	Gly	Ser	Leu	Glu	Arg	Asp	Glu	Val	Phe		
				145				150					155				
GTC	CCC	AAG	AAG	AAG	TCA	GAC	ATC	AAG	TGG	TAC	GGC	AAG	TAC	CTC	AAC	648	
Val	Pro	Lys	Lys	Lys	Ser	Asp	Ile	Lys	Trp	Tyr	Gly	Lys	Tyr	Leu	Asn		
		160					165					170					
AAC	CCT	TTG	GGA	CGC	ACC	GTG	ATG	TTA	ACG	GTT	CAG	TTC	ACT	CTC	GGC	696	
Asn	Pro	Leu	Gly	Arg	Thr	Val	Met	Leu	Thr	Val	Gln	Phe	Thr	Leu	Gly		
		175				180					185						
TGG	CCT	TTG	TAC	TTA	GCC	TTC	AAC	GTC	TCG	GGG	AGA	CCT	TAC	GAC	GGC	744	
Trp	Pro	Leu	Tyr	Leu	Ala	Phe	Asn	Val	Ser	Gly	Arg	Pro	Tyr	Asp	Gly		
190					195					200					205		
GGC	TTC	GCT	TGC	CAT	TTC	CAC	CCC	AAC	GCT	CCC	ATC	TAC	AAC	GAC	CGT	792	
Gly	Phe	Ala	Cys	His	Phe	His	Pro	Asn	Ala	Pro	Ile	Tyr	Asn	Asp	Arg		
				210					215					220			
GAG	CGT	CTC	CAG	ATA	TAC	ATC	TCC	GAC	GCT	GGC	ATC	CTC	GCC	GTC	TGC	840	
Glu	Arg	Leu	Gln	Ile	Tyr	Ile	Ser	Asp	Ala	Gly	Ile	Leu	Ala	Val	Cys		
			225					230					235				
TAC	GGT	CTC	TAC	CGC	TAC	GCT	GCT	GTC	CAA	GGA	GTT	GCC	TCG	ATG	GTC	888	
Tyr	Gly	Leu	Tyr	Arg	Tyr	Ala	Ala	Val	Gln	Gly	Val	Ala	Ser	Met	Val		
		240				245						250					
TGC	TTC	TAC	GGA	GTT	CCT	CTT	CTG	ATT	GTC	AAC	GGG	TTC	TTA	GTT	TTG	936	
Cys	Phe	Tyr	Gly	Val	Pro	Leu	Leu	Ile	Val	Asn	Gly	Phe	Leu	Val	Leu		
		255				260					265						
ATC	ACT	TAC	TTG	CAG	CAC	ACG	CAT	CCT	TCC	CTG	CCT	CAC	TAT	GAC	TCG	984	
Ile	Thr	Tyr	Leu	Gln	His	Thr	His	Pro	Ser	Leu	Pro	His	Tyr	Asp	Ser		
270					275					280				285			
TCT	GAG	TGG	GAT	TGG	TTG	AGG	GGA	GCT	TTG	GCC	ACC	GTT	GAC	AGA	GAC	1032	

Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp  
 290 295 300

TAC GGA ATC TTG AAC AAG GTC TTC CAC AAT ATC ACG GAC ACG CAC GTG 1080  
 Tyr Gly Ile Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val  
 305 310 315

GCG CAT CAC CTG TTC TCG ACC ATG CCG CAT TAT CAT GCG ATG GAA GCT 1128  
 Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala  
 320 325 330

ACG AAG GCG ATA AAG CCG ATA CTG GGA GAG TAT TAT CAG TTC GAT GGG 1176  
 Thr Lys Ala Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly  
 335 340 345

ACG CCG GTG GTT AAG GCG ATG TGG AGG GAG GCG AAG GAG TGT ATC TAT 12  
 24  
 Thr Pro Val Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr  
 350 355 360 365

GTG GAA CCG GAC AGG CAA GGT GAG AAG AAA GGT GTG TTC TGG TAC AAC 12  
 72  
 Val Glu Pro Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn  
 370 375 380

AAT AAG TTA TGA AGCAAAGAAG AAAGTGAACC TTCTCTTCT ATGATTGTCT 13  
 24  
 Asn Lys Leu

TTGTTTAAGA AGCTATGTTT CTGTTTCAAT AATCTTAATT ATCCATTTTG TTGTGTTTTC 13  
 84

TGACATTTTG GCTAAAATTA TGTGATGTTG GAAGTTAGTG TC 14  
 26

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 384 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Gly Ala Gly Gly Arg Met Gln Val Ser Pro Pro Ser Lys Lys Ser  
 1 5 10 15

Glu Thr Asp Asn Ile Lys Arg Val Pro Cys Glu Thr Pro Pro Phe Thr  
 20 25 30

Val Gly Glu Leu Lys Lys Ala Ile Pro Pro His Cys Phe Lys Arg Ser  
 35 40 45

```

Ile Pro Arg Ser Phe Ser Tyr Leu Ile Trp Asp Ile Ile Ile Ala Ser
 50                      55                      60

Cys Phe Tyr Tyr Val Ala Thr Thr Tyr Phe Pro Leu Leu Pro His Pro
 65                      70                      75                      80

Leu Ser Tyr Phe Ala Trp Pro Leu Tyr Trp Ala Cys Gln Gly Cys Val
      85                      90                      95

Leu Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His Ala Phe
      100                      105                      110

Ser Asp Tyr Gln Trp Leu Asp Asp Thr Val Gly Leu Ile Phe His Ser
      115                      120                      125

Phe Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg Arg His
      130                      135                      140

His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys
      145                      150                      155                      160

Lys Lys Ser Asp Ile Lys Trp Tyr Gly Lys Tyr Leu Asn Asn Pro Leu
      165                      170                      175

Gly Arg Thr Val Met Leu Thr Val Gln Phe Thr Leu Gly Trp Pro Leu
      180                      185                      190

Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp Gly Gly Phe Ala
      195                      200                      205

Cys His Phe His Pro Asn Ala Pro Ile Tyr Asn Asp Arg Glu Arg Leu
      210                      215                      220

Gln Ile Tyr Ile Ser Asp Ala Gly Ile Leu Ala Val Cys Tyr Gly Leu
      225                      230                      235                      240

Tyr Arg Tyr Ala Ala Val Gln Gly Val Ala Ser Met Val Cys Phe Tyr
      245                      250                      255

Gly Val Pro Leu Leu Ile Val Asn Gly Phe Leu Val Leu Ile Thr Tyr
      260                      265                      270

Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser Glu Trp
      275                      280                      285

Asp Trp Leu Arg Gly Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile
      290                      295                      300

Leu Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His
      305                      310                      315                      320

Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala
      325                      330                      335

Ile Lys Pro Ile Leu Gly Glu Tyr Tyr Gln Phe Asp Gly Thr Pro Val
      340                      345                      350

```

09697379.102600

```

Val Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Ile Tyr Val Glu Pro
    355                      360                      365
Asp Arg Gln Gly Glu Lys Lys Gly Val Phe Trp Tyr Asn Asn Lys Leu
    370                      375                      380

```

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1462 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Glycine max

## (vii) IMMEDIATE SOURCE:

- (B) CLONE: pSF2-165K

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 108..1247

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

CCATATACTA ATATTTGCTT GTATTGATAG CCCCTCCGTT CCCAAGAGTA TAAAACTGCA
    60

```

```

TCGAATAATA CAAGCCACTA GGCATGGGTC TAGCAAAGGA AACAAACA ATG GGA GGT
    116

```

```

Met Gly Gly
    1

```

```

AGA GGT CGT GTG GCC AAA GTG GAA GTT CAA GGG AAG AAG CCT CTC TCA
    164

```

```

Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys Pro Leu Ser
    5                      10                      15

```

```

AGG GTT CCA AAC ACA AAG CCA CCA TTC ACT GTT GGC CAA CTC AAG AAA
    212

```

```

Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln Leu Lys Lys
    20                      25                      30                      35

```

```

GCA ATT CCA CCA CAC TGC TTT CAG CGC TCC CTC CTC ACT TCA TTC TCC
    260

```

```

Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr Ser Phe Ser
    40                      45                      50

```

TAT GTT GTT TAT GAC CTT TCA TTT GCC TTC ATT TTC TAC ATT GCC ACC  
 308  
 Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr Ile Ala Thr  
 55 60 65

ACC TAC TTC CAC CTC CTT CCT CAA CCC TTT TCC CTC ATT GCA TGG CCA  
 356  
 Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile Ala Trp Pro  
 70 75 80

ATC TAT TGG GTT CTC CAA GGT TGC CTT CTC ACT GGT GTG TGG GTG ATT  
 404  
 Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val Trp Val Ile  
 85 90 95

GCT CAC GAG TGT GGT CAC CAT GCC TTC AGC AAG TAC CAA TGG GTT GAT  
 452  
 Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln Trp Val Asp  
 100 105 110 115

GAT GTT GTG GGT TTG ACC CTT CAC TCA ACA CTT TTA GTC CCT TAT TTC  
 500  
 Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val Pro Tyr Phe  
 120 125 130

TCA TGG AAA ATA AGC CAT CGC CGC CAT CAC TCC AAC ACA GGT TCC CTT  
 548  
 Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu  
 135 140 145

GAC CGT GAT GAA GTG TTT GTC CCA AAA CCA AAA TCC AAA GTT GCA TGG  
 596  
 Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys Val Ala Trp  
 150 155 160

TTT TCC AAG TAC TTA AAC AAC CCT CTA GGA AGG GCT GTT TCT CTT CTC  
 644  
 Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val Ser Leu Leu  
 165 170 175

GTC ACA CTC ACA ATA GGG TGG CCT ATG TAT TTA GCC TTC AAT GTC TCT  
 692  
 Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe Asn Val Ser  
 180 185 190 195

GGT AGA CCC TAT GAT AGT TTT GCA AGC CAC TAC CAC CCT TAT GCT CCC  
 740  
 Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro Tyr Ala Pro  
 200 205 210

ATA TAT TCT AAC CGT GAG AGG CTT CTG ATC TAT GTC TCT GAT GTT GCT  
 788  
 Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser Asp Val Ala  
 215 220 225

TTG TTT TCT GTG ACT TAC TCT CTC TAC CGT GTT GCA ACC CTG AAA GGG

00697379.102600

836  
 Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr Leu Lys Gly  
           230                          235                          240

TTG GTT TGG CTG CTA TGT GTT TAT GGG GTG CCT TTG CTC ATT GTG AAC  
 884  
 Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu Ile Val Asn  
           245                          250                          255

GGT TTT CTT GTG ACT ATC ACA TAT TTG CAG CAC ACA CAC TTT GCC TTG  
 932  
 Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His Phe Ala Leu  
           260                          265                          270                          275

CCT CAT TAC GAT TCA TCA GAA TGG GAC TGG CTG AAG GGA GCT TTG GCA  
 980  
 Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly Ala Leu Ala  
                           280                          285                          290

ACT ATG GAC AGA GAT TAT GGG ATT CTG AAC AAG GTG TTT CAT CAC ATA  
 1028  
 Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe His His Ile  
                           295                          300                          305

ACT GAT ACT CAT GTG GCT CAC CAT CTC TTC TCT ACA ATG CCA CAT TAC  
 1076  
 Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr  
                           310                          315                          320

CAT GCA ATG GAG GCA ACC AAT GCA ATC AAG CCA ATA TTG GGT GAG TAC  
 1124  
 His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu Gly Glu Tyr  
           325                          330                          335

TAC CAA TTT GAT GAC ACA CCA TTT TAC AAG GCA CTG TGG AGA GAA GCG  
 1172  
 Tyr Gln Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala  
           340                          345                          350                          355

AGA GAG TGC CTC TAT GTG GAG CCA GAT GAA GGA ACA TCC GAG AAG GGC  
 1220  
 Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser Glu Lys Gly  
                           360                          365                          370

GTG TAT TGG TAC AGG AAC AAG TAT TGATGGAGCA ACCAATGGGC CATAGTGGGA  
 1274  
 Val Tyr Trp Tyr Arg Asn Lys Tyr  
                           375                          380

GTTATGGAAG TTTTGTGTCATG TATTAGTACA TAATTAGTAG AATGTTATAA ATAAGTGGAT  
 1334

TTGCCGCGTA ATGACTTTGT GTGTATTGTG AAACAGCTTG TTGCGATCAT GGTATAATG  
 1394

TAAAAATAAT TCTGGTATTA ATTACATGTG GAAAGTGTC TGCTTATAGC TTTCTGCCTA

1454

AAAAAAAAA

1462

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 379 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Gly Gly Arg Gly Arg Val Ala Lys Val Glu Val Gln Gly Lys Lys
 1          5          10          15
Pro Leu Ser Arg Val Pro Asn Thr Lys Pro Pro Phe Thr Val Gly Gln
          20          25          30
Leu Lys Lys Ala Ile Pro Pro His Cys Phe Gln Arg Ser Leu Leu Thr
          35          40          45
Ser Phe Ser Tyr Val Val Tyr Asp Leu Ser Phe Ala Phe Ile Phe Tyr
          50          55          60
Ile Ala Thr Thr Tyr Phe His Leu Leu Pro Gln Pro Phe Ser Leu Ile
          65          70          75          80
Ala Trp Pro Ile Tyr Trp Val Leu Gln Gly Cys Leu Leu Thr Gly Val
          85          90          95
Trp Val Ile Ala His Glu Cys Gly His His Ala Phe Ser Lys Tyr Gln
          100          105          110
Trp Val Asp Asp Val Val Gly Leu Thr Leu His Ser Thr Leu Leu Val
          115          120          125
Pro Tyr Phe Ser Trp Lys Ile Ser His Arg Arg His His Ser Asn Thr
          130          135          140
Gly Ser Leu Asp Arg Asp Glu Val Phe Val Pro Lys Pro Lys Ser Lys
          145          150          155          160
Val Ala Trp Phe Ser Lys Tyr Leu Asn Asn Pro Leu Gly Arg Ala Val
          165          170          175
Ser Leu Leu Val Thr Leu Thr Ile Gly Trp Pro Met Tyr Leu Ala Phe
          180          185          190
Asn Val Ser Gly Arg Pro Tyr Asp Ser Phe Ala Ser His Tyr His Pro
          195          200          205
Tyr Ala Pro Ile Tyr Ser Asn Arg Glu Arg Leu Leu Ile Tyr Val Ser

```

210	215	220
Asp Val Ala Leu Phe Ser Val Thr Tyr Ser Leu Tyr Arg Val Ala Thr 225 230 235 240		
Leu Lys Gly Leu Val Trp Leu Leu Cys Val Tyr Gly Val Pro Leu Leu 245 250 255		
Ile Val Asn Gly Phe Leu Val Thr Ile Thr Tyr Leu Gln His Thr His 260 265 270		
Phe Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Lys Gly 275 280 285		
Ala Leu Ala Thr Met Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe 290 295 300		
His His Ile Thr Asp Thr His Val Ala His His Leu Phe Ser Thr Met 305 310 315 320		
Pro His Tyr His Ala Met Glu Ala Thr Asn Ala Ile Lys Pro Ile Leu 325 330 335		
Gly Glu Tyr Tyr Glu Phe Asp Asp Thr Pro Phe Tyr Lys Ala Leu Trp 340 345 350		
Arg Glu Ala Arg Glu Cys Leu Tyr Val Glu Pro Asp Glu Gly Thr Ser 355 360 365		
Glu Lys Gly Val Tyr Trp Tyr Arg Asn Lys Tyr 370 375		

## (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1790 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (A) ORGANISM: Zea mays
- (vii) IMMEDIATE SOURCE:
  - (B) CLONE: pFad2#1
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 165..1328



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGGCCTCTCC CCTCCTCTCT CCCTGCAAAT CCTGCAGACA CCACCGCTCG TTTTCTCTCT  
60

CGGGACAGGA GAAAAGGGGA GAGAGAGGTG AGGCGCGGTG TCCGCCCGAT CTGCTCTGCC  
120

CCGACGCAGC TGTTACGACC TCCTCAGTCT CAGTCAGGAG CAAG ATG GGT GCC GGC  
176

Met Gly Ala Gly  
1

GGC AGG ATG ACC GAG AAG GAG CGG GAG AAG CAG GAG CAG CTC GCC CGA  
224

Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu Gln Leu Ala Arg  
5 10 15 20

GCT ACC GGT GGC GCC GCG ATG CAG CGG TCG CCG GTG GAG AAG CCT CCG  
272

Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val Glu Lys Pro Pro  
25 30 35

TTC ACT CTG GGT CAG ATC AAG AAG GCC ATC CCG CCA CAC TGC TTC GAG  
320

Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro His Cys Phe Glu  
40 45 50

CGC TCG GTG CTC AAG TCC TTC TCG TAC GTG GTC CAC GAC CTG GTG ATC  
368

Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His Asp Leu Val Ile  
55 60 65

GCC GCG GCG CTC CTC TAC TTC GCG CTG GCC ATC ATA CCG GCG CTC CCA  
416

Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile Pro Ala Leu Pro  
70 75 80

AGC CCG CTC CGC TAC GCC GCC TGG CCG CTG TAC TGG ATC GCG CAG GGG  
464

Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp Ile Ala Gln Gly  
85 90 95 100

TGC GTG TGC ACC GGC GTG TGG GTC ATC GCG CAC GAG TGC GGC CAC CAC  
512

Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu Cys Gly His His  
105 110 115

GCC TTC TCG GAC TAC TCG CTC CTG GAC GAC GTG GTC GGC CTG GTG CTG  
560

Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val Gly Leu Val Leu  
120 125 130

CAC TCG TCG CTC ATG GTG CCC TAC TTC TCG TGG AAG TAC AGC CAC CGG  
608

His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys Tyr Ser His Arg

135	140	145
CGC CAC CAC TCC AAC ACG GGG TCC CTG GAG CGC GAC GAG GTG TTC GTG		
656		
Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp Glu Val Phe Val		
150	155	160
CCC AAG AAG AAG GAG GCG CTG CCG TGG TAC ACC CCG TAC GTG TAC AAC		
704		
Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro Tyr Val Tyr Asn		
165	170	180
AAC CCG GTC GGC CGG GTG GTG CAC ATC GTG GTG CAG CTC ACC CTC GGG		
752		
Asn Pro Val Gly Arg Val Val His Ile Val Val Gln Leu Thr Leu Gly		
	185	195
TGG CCG CTG TAC CTG GCG ACC AAC GCG TCG GGG CGG CCG TAC CCG CGC		
800		
Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg Pro Tyr Pro Arg		
	200	210
TTC GCC TGC CAC TTC GAC CCC TAC GGC CCC ATC TAC AAC GAC CGG GAG		
848		
Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr Asn Asp Arg Glu		
	215	225
CGC GCC CAG ATC TTC GTC TCG GAC GCC GGC GTC GTG GCC GTG GCG TTC		
896		
Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val Ala Val Ala Phe		
	230	240
GGG CTG TAC AAG CTG GCG GCG GCG TTC GGG GTC TGG TGG GTG GTG CGC		
944		
Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp Trp Val Val Arg		
245	250	260
GTG TAC GCC GTG CCG CTG CTG ATC GTG AAC GCG TGG CTG GTG CTC ATC		
992		
Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp Leu Val Leu Ile		
	265	275
ACC TAC CTG CAG CAC ACC CAC CCG TCG CTC CCC CAC TAC GAC TCG AGC		
1040		
Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His Tyr Asp Ser Ser		
	280	290
GAG TGG GAC TGG CTG CGC GGC GCG CTG GCC ACC ATG GAC CGC GAC TAC		
1088		
Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met Asp Arg Asp Tyr		
	295	305
GGC ATC CTC AAC CGC GTG TTC CAC AAC ATC ACG GAC ACG CAC GTC GCG		
1136		
Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp Thr His Val Ala		
	310	320

CAC CAC CTC TTC TCC ACC ATG CCG CAC TAC CAC GCC ATG GAG GCC ACC  
1184  
His His Leu Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr  
325 330 335 340  
AAG GCG ATC AGG CCC ATC CTC GGC GAC TAC TAC CAC TTC GAC CCG ACC  
1232  
Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His Phe Asp Pro Thr  
345 350 355  
CCT GTC GCC AAG GCG ACC TGG CGC GAG GCC GGG GAA TGC ATC TAC GTC  
1280  
Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu Cys Ile Tyr Val  
360 365 370  
GAG CCC GAG GAC CGC AAG GGC GTC TTC TGG TAC AAC AAG AAG TTC TAGCCGCCG  
C 1335  
Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn Lys Lys Phe  
375 380 385  
CGCTCGCAGA GCTGAGGACG CTACCGTAGG AATGGGAGCA GAAACCAGGA GGAGGAGACG  
1395  
GTACTCGCCC CAAAGTCTCC GTCAACCTAT CTAATCGTTA GTCGTCAGTC TTTTAGACGG  
1455  
GAAGAGAGAT CATTTGGGCA CAGAGACGAA GGCTTACTGC AGTGCCATCG CTAGAGCTGC  
1515  
CATCAAGTAC AAGTAGGCAA ATTCGTCAAC TTAGTGTGTC CCATGTTGTT TTTCTTAGTC  
1575  
GTCCGCTGCT GTAGGCTTTC CGGCGGCGGT CGTTTGTGTG GTTGGCATCC GTGGCCATGC  
1635  
CTGTGCGTGC GTGGCCGCGC TTGTCTGTGT CGTCTGTCGT CGCGTTGGCG TCGTCTCTTC  
1695  
GTGCTCCCCG TGTGTTGTTG TAAACAAGA AGATGTTTTC TGGTGCTTTT GGCGGAATAA  
1755  
CAGATCGTCC GAACGAAAAA AAAAAAAAAA AAAAA  
1790

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 387 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Gly Ala Gly Gly Arg Met Thr Glu Lys Glu Arg Glu Lys Gln Glu  
 1 5 10 15  
 Gln Leu Ala Arg Ala Thr Gly Gly Ala Ala Met Gln Arg Ser Pro Val  
 20 25 30  
 Glu Lys Pro Pro Phe Thr Leu Gly Gln Ile Lys Lys Ala Ile Pro Pro  
 35 40 45  
 His Cys Phe Glu Arg Ser Val Leu Lys Ser Phe Ser Tyr Val Val His  
 50 55 60  
 Asp Leu Val Ile Ala Ala Ala Leu Leu Tyr Phe Ala Leu Ala Ile Ile  
 65 70 75 80  
 Pro Ala Leu Pro Ser Pro Leu Arg Tyr Ala Ala Trp Pro Leu Tyr Trp  
 85 90 95  
 Ile Ala Gln Gly Cys Val Cys Thr Gly Val Trp Val Ile Ala His Glu  
 100 105 110  
 Cys Gly His His Ala Phe Ser Asp Tyr Ser Leu Leu Asp Asp Val Val  
 115 120 125  
 Gly Leu Val Leu His Ser Ser Leu Met Val Pro Tyr Phe Ser Trp Lys  
 130 135 140  
 Tyr Ser His Arg Arg His His Ser Asn Thr Gly Ser Leu Glu Arg Asp  
 145 150 155 160  
 Glu Val Phe Val Pro Lys Lys Lys Glu Ala Leu Pro Trp Tyr Thr Pro  
 165 170 175  
 Tyr Val Tyr Asn Asn Pro Val Gly Arg Val Val His Ile Val Val Gln  
 180 185 190  
 Leu Thr Leu Gly Trp Pro Leu Tyr Leu Ala Thr Asn Ala Ser Gly Arg  
 195 200 205  
 Pro Tyr Pro Arg Phe Ala Cys His Phe Asp Pro Tyr Gly Pro Ile Tyr  
 210 215 220  
 Asn Asp Arg Glu Arg Ala Gln Ile Phe Val Ser Asp Ala Gly Val Val  
 225 230 235 240  
 Ala Val Ala Phe Gly Leu Tyr Lys Leu Ala Ala Ala Phe Gly Val Trp  
 245 250 255  
 Trp Val Val Arg Val Tyr Ala Val Pro Leu Leu Ile Val Asn Ala Trp  
 260 265 270  
 Leu Val Leu Ile Thr Tyr Leu Gln His Thr His Pro Ser Leu Pro His  
 275 280 285  
 Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Leu Ala Thr Met  
 290 295 300

Asp Arg Asp Tyr Gly Ile Leu Asn Arg Val Phe His Asn Ile Thr Asp  
 305 310 315 320  
 Thr His Val Ala His His Leu Phe Ser Thr Met Pro His Tyr His Ala  
 325 330 335  
 Met Glu Ala Thr Lys Ala Ile Arg Pro Ile Leu Gly Asp Tyr Tyr His  
 340 345 350  
 Phe Asp Pro Thr Pro Val Ala Lys Ala Thr Trp Arg Glu Ala Gly Glu  
 355 360 365  
 Cys Ile Tyr Val Glu Pro Glu Asp Arg Lys Gly Val Phe Trp Tyr Asn  
 370 375 380  
 Lys Lys Phe  
 385

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 673 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Ricinus communis*

- (vii) IMMEDIATE SOURCE:  
 (B) CLONE: pRF2-1C

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..673

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGG GTG ATG GCG CAT GAT TGT GGG CAC CAT GCC TTC AGT GAC TAT CAA  
 49  
 Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln  
 1 5 10 15  
 TTG CTT GAT GAT GTA GTT GGT CTT ATC CTA CAC TCC TGT CTC CTT GTC  
 96  
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val  
 20 25 30  
 CCT TAT TTT TCA TGG AAA CAC AGC CAT CGC CGA CAT CAT TCC AAC ACA

144  
Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr  
35 40 45

GGG TCC CTG GAA CGG GAT GAA GTG TTT GTT CCC AAG AAG AAA TCT AGT  
192  
Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Lys Ser Ser  
50 55 60

ATC CGT TGG TAT TCC AAA TAC CTC AAC AAC CCT CCA GGT CGT ATC ATG  
240  
Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met  
65 70 75 80

ACA ATT GCC GTC ACA CTT TCA CTT GGC TGG CCT CTG TAC CTA GCA TTC  
288  
Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe  
85 90 95

AAT GTT TCA GGC AGG CCA TAT GAT CGG TTC GCC TGC CAC TAT GAC CCA  
336  
Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro  
100 105 110

TAT GGC CCG ATC TAC AAT GAT CGC GAG CGA ATC GAG ATA TTC ATA TCA  
384  
Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser  
115 120 125

GAT GCT GGT GTT CTT GCT GTC ACT TTT GGT CTC TAC CAA CTT GCT ATA  
432  
Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile  
130 135 140

GCG AAG GGG CTT GCT TGG GTT GTC TGT GTA TAT GGA GTG CCA TTG TTG  
480  
Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu  
145 150 155 160

GTG GTG AAT TCA TTC CTT GTT CTG ATC ACA TTT CTG CAG CAT ACT CAC  
528  
Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His  
165 170 175

CCT GCA TTG CCA CAT TAT GAT TCG TCG GAG TGG GAC TGG CTA AGA GGA  
576  
Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly  
180 185 190

GCT CTA GCA ACT GTT GAC AGA GAT TAC GGG ATC TTG AAC AAG GTG TTC  
624  
Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
195 200 205

CAT AAC ATA ACG GAC ACT CAA GTA GCT CAC CAC CTT TTC ACC ATG CCC C  
673

His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 224 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Trp Val Met Ala His Asp Cys Gly His His Ala Phe Ser Asp Tyr Gln  
 1 5 10 15  
 Leu Leu Asp Asp Val Val Gly Leu Ile Leu His Ser Cys Leu Leu Val  
 20 25 30  
 Pro Tyr Phe Ser Trp Lys His Ser His Arg Arg His His Ser Asn Thr  
 35 40 45  
 Gly Ser Leu Glu Arg Asp Glu Val Phe Val Pro Lys Lys Ser Ser  
 50 55 60  
 Ile Arg Trp Tyr Ser Lys Tyr Leu Asn Asn Pro Pro Gly Arg Ile Met  
 65 70 75 80  
 Thr Ile Ala Val Thr Leu Ser Leu Gly Trp Pro Leu Tyr Leu Ala Phe  
 85 90 95  
 Asn Val Ser Gly Arg Pro Tyr Asp Arg Phe Ala Cys His Tyr Asp Pro  
 100 105 110  
 Tyr Gly Pro Ile Tyr Asn Asp Arg Glu Arg Ile Glu Ile Phe Ile Ser  
 115 120 125  
 Asp Ala Gly Val Leu Ala Val Thr Phe Gly Leu Tyr Gln Leu Ala Ile  
 130 135 140  
 Ala Lys Gly Leu Ala Trp Val Val Cys Val Tyr Gly Val Pro Leu Leu  
 145 150 155 160  
 Val Val Asn Ser Phe Leu Val Leu Ile Thr Phe Leu Gln His Thr His  
 165 170 175  
 Pro Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp Trp Leu Arg Gly  
 180 185 190  
 Ala Leu Ala Thr Val Asp Arg Asp Tyr Gly Ile Leu Asn Lys Val Phe  
 195 200 205  
 His Asn Ile Thr Asp Thr Gln Val Ala His His Leu Phe Thr Met Pro  
 210 215 220

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1369 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Ricinus communis*

(vii) IMMEDIATE SOURCE:

(B) CLONE: pRF197c-42

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 184..1347

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CGGCCGGGAT TCCGGTTTTC ACTACTAATTT GCAAAAAATG CATGATTTC CCTCAAATCA  
 60

AACACCACAC CTTATAACTT AGTCTTAAGA GAGAGAGAGA GAGGAGACAT TTCTCTTCTC  
 120

TGAGATGAGC ACTTCTCTTC CAGACATCGA AGCCTCAGGA AAGTGCTTGA GAAGAGCTTG  
 180

AGA ATG GGA GGT GGT GGT CGC ATG TCT ACT GTC ATA ATC AGC AAC AAC  
 228

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn  
 1 5 10 15

AGT GAG AAG AAA GGA GGA AGC AGC CAC CTG GAG CGA GCG CCG CAC ACG  
 276

Ser Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr  
 20 25 30

AAG CCT CCT TAC ACA CTT GGT AAC CTC AAG AGA GCC ATC CCA CCC CAT  
 324

Lys Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro Pro His  
 35 40 45

TGC TTT GAA CGC TCT TTT GTG CGC TCA TTC TCC AAT TTT GCC TAT AAT  
 372

Cys Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn  
 50 55 60

TTC TGC TTA AGT TTT CTT TCC TAC TCG ATC GCC ACC AAC TTC TTC CCT



420  
 Phe Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro  
 65 70 75  
 TAC ATC TCT TCT CCG CTC TCG TAT GTC GCT TGG CTG GTT TAC TGG CTC  
 468  
 Tyr Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu  
 80 85 90 95  
 TTC CAA GGC TGC ATT CTC ACT GGT CTT TGG GTC ATC GGC CAT GAA TGT  
 516  
 Phe Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys  
 100 105 110  
 GGC CAT CAT GCT TTT AGT GAG TAT CAG CTG GCT GAT GAC ATT GTT GGC  
 564  
 Gly His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly  
 115 120 125  
 CTA ATT GTC CAT TCT GCA CTT CTG GTT CCA TAT TTT TCA TGG AAA TAT  
 612  
 Leu Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr  
 130 135 140  
 AGC CAT CGC CGC CAC CAT TCT AAC ATA GGA TCT CTC GAG CGA GAC GAA  
 660  
 Ser His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu  
 145 150 155  
 GTG TTC GTC CCG AAA TCA AAG TCG AAA ATT TCA TGG TAT TCT AAG TAC  
 708  
 Val Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr  
 160 165 170 175  
 TTA AAC AAC CCG CCA GGT CGA GTT TTG ACA CTT GCT GCC ACG CTC CTC  
 756  
 Leu Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu  
 180 185 190  
 CTT GGC TGG CCT TTA TAT TTA GCT TTC AAT GTC TCT GGT AGA CCT TAC  
 804  
 Leu Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr  
 195 200 205  
 GAT CGC TTT GCT TGC CAT TAT GAT CCC TAT GGC CCA ATA TTT TCC GAA  
 852  
 Asp Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu  
 210 215 220  
 AGA GAA AGG CTT CAG ATT TAC ATT GCT GAC CTC GGA ATC TTT GCC ACA  
 900  
 Arg Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr  
 225 230 235  
 ACG TTT GTG CTT TAT CAG GCT ACA ATG GCA AAA GGG TTG GCT TGG GTA  
 948

Thr Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val  
240 245 250 255

ATG CGT ATC TAT GGG GTG CCA TTG CTT ATT GTT AAC TGT TTC CTT GTT  
996

Met Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val  
260 265 270

ATG ATC ACA TAC TTG CAG CAC ACT CAC CCA GCT ATT CCA CGC TAT GGC  
1044

Met Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly  
275 280 285

TCA TCG GAA TGG GAT TGG CTC CGG GGA GCA ATG GTG ACT GTC GAT AGA  
1092

Ser Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg  
290 295 300

GAT TAT GGG GTG TTG AAT AAA GTA TTC CAT AAC ATT GCA GAC ACT CAT  
1140

Asp Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His  
305 310 315

GTA GCT CAT CAT CTC TTT GCT ACA GTG CCA CAT TAC CAT GCA ATG GAG  
1188

Val Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu  
320 325 330 335

GCC ACT AAA GCA ATC AAG CCT ATA ATG GGT GAG TAT TAC CGG TAT GAT  
1236

Ala Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp  
340 345 350

GGT ACC CCA TTT TAC AAG GCA TTG TGG AGG GAG GCA AAG GAG TGC TTG  
1284

Gly Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu  
355 360 365

TTC GTC GAG CCA GAT GAA GGA GCT CCT ACA CAA GGC GTT TTC TGG TAC  
1332

Phe Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr  
370 375 380

CGG AAC AAG TAT TAAAAAAGTG TCATGTAGCC TGCCG  
1369

Arg Asn Lys Tyr  
385

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 387 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

Met Gly Gly Gly Gly Arg Met Ser Thr Val Ile Ile Ser Asn Asn Ser
 1           5           10           15

Glu Lys Lys Gly Gly Ser Ser His Leu Glu Arg Ala Pro His Thr Lys
 20           25           30

Pro Pro Tyr Thr Leu Gly Asn Leu Lys Arg Ala Ile Pro His Cys
 35           40           45

Phe Glu Arg Ser Phe Val Arg Ser Phe Ser Asn Phe Ala Tyr Asn Phe
 50           55           60

Cys Leu Ser Phe Leu Ser Tyr Ser Ile Ala Thr Asn Phe Phe Pro Tyr
 65           70           75           80

Ile Ser Ser Pro Leu Ser Tyr Val Ala Trp Leu Val Tyr Trp Leu Phe
 85           90           95

Gln Gly Cys Ile Leu Thr Gly Leu Trp Val Ile Gly His Glu Cys Gly
100           105           110

His His Ala Phe Ser Glu Tyr Gln Leu Ala Asp Asp Ile Val Gly Leu
115           120           125

Ile Val His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Tyr Ser
130           135           140

His Arg Arg His His Ser Asn Ile Gly Ser Leu Glu Arg Asp Glu Val
145           150           155           160

Phe Val Pro Lys Ser Lys Ser Lys Ile Ser Trp Tyr Ser Lys Tyr Leu
165           170           175

Asn Asn Pro Pro Gly Arg Val Leu Thr Leu Ala Ala Thr Leu Leu Leu
180           185           190

Gly Trp Pro Leu Tyr Leu Ala Phe Asn Val Ser Gly Arg Pro Tyr Asp
195           200           205

Arg Phe Ala Cys His Tyr Asp Pro Tyr Gly Pro Ile Phe Ser Glu Arg
210           215           220

Glu Arg Leu Gln Ile Tyr Ile Ala Asp Leu Gly Ile Phe Ala Thr Thr
225           230           235           240

Phe Val Leu Tyr Gln Ala Thr Met Ala Lys Gly Leu Ala Trp Val Met
245           250           255

Arg Ile Tyr Gly Val Pro Leu Leu Ile Val Asn Cys Phe Leu Val Met
260           265           270

Ile Thr Tyr Leu Gln His Thr His Pro Ala Ile Pro Arg Tyr Gly Ser
275           280           285

```

Ser Glu Trp Asp Trp Leu Arg Gly Ala Met Val Thr Val Asp Arg Asp  
 290 295 300

Tyr Gly Val Leu Asn Lys Val Phe His Asn Ile Ala Asp Thr His Val  
 305 310 315 320

Ala His His Leu Phe Ala Thr Val Pro His Tyr His Ala Met Glu Ala  
 325 330 335

Thr Lys Ala Ile Lys Pro Ile Met Gly Glu Tyr Tyr Arg Tyr Asp Gly  
 340 345 350

Thr Pro Phe Tyr Lys Ala Leu Trp Arg Glu Ala Lys Glu Cys Leu Phe  
 355 360 365

Val Glu Pro Asp Glu Gly Ala Pro Thr Gln Gly Val Phe Trp Tyr Arg  
 370 375 380

Asn Lys Tyr  
 385

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 1..23
- (D) OTHER INFORMATION: /product="synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGGGTATGCC AYGANTGYGG NCA

23

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) TRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc\_feature

(B) LOCATION: 1..22

(D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAARTGRTGG CACRTGNGTR TC

22

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2973 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Arabidopsis thaliana

(vii) IMMEDIATE SOURCE:

(B) CLONE: pAGF2-6

(ix) FEATURE:

(A) NAME/KEY: exon

(B) LOCATION: 433..520

(ix) FEATURE:

(A) NAME/KEY: intron

(B) LOCATION: 521..1654

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATTGCGTAAT TCCTACATAT TTTAGAGATT AGTTTGAGTT TCCATCCATA CTTTACTAGT  
60

GATTATAAAT TTAAATACG TACTTTTTCGA CTATAAAGTG AACTAAGTA AATTAGAACG 1  
20

TGATATTAAA AAGTTAATGT TCACTGTTAT ATTTTTCCTA CAAGTAAAAA ATGGGTTATT 1  
80

TGCGGTAAT AAAAATACCA GATATTTTGA ATTGATTAAA AAGTTTGAAA TAAGAGAGGA 2

40

```

GGGGAAAGAA AAGAAGGTGG GGGCCCAGTA TGAAAGGGAA AGGTGTCATC AAATCATCTC 3
00

TCTCTCTCTC TACCTTCGAC CCACGGGCCG TGTCCTTTA AAGCCCTGTC TCTTGCCATT 3
60

CCCCATCTGA CCACCAGAAG AAGAGCCACA CACTCACAAA TTA AAAAGAG AGAGAGAGAG 4
20

AGAGAGACAG AGAGAGAGAG AGATTCTGCG GAGGAGCTTC TTCTTCGTAG GGTGTTTCATC 4
80

GTTATTAAACG TTATCGCCCC TACGTCAGCT CCATCTCCAG GTCCGTCGCT TCTCTCCAT 5
40

TTCTTCTCAT TTTCGATTTT GATTCTTATT TCTTCCAGT AGCTCCTGCT CTGTGAATTT 6
00

CTCCGCTCAC GATAGATCTG CTTATACTCC TTACATTCAA CCTTAGATCT GGTCTCGATT 6
60

CTCTGTTTCT CTGTTTTTTT CTTTGGTTCG AGAATCTGAT GTTGTTTAT GTTCTGTCAC 7
20

CATTAATAAT GATGAACCTCT CTCATTCATA CAATGATTAG TTCTCTCGT CTACCAAACG 7
80

ATATGTTGCA TTTTCACTTT TCTTCTTTT TTCTAAGATG ATTTGCTTTG ACCAATTTGT 8
40

TTAGATCTTT ATTTTATTTT ATTTTCTGGT GGGTTGGTGG AAATTGAAAA AAAAAAAAAA 9
00

AAAAGCATAA ATTGTTATTT GTTAATGTAT TCATTTTTTG GCTATTTGTT CTGGGTAAAA 9
60

ATCTGCTTCT ACTGTTGAAT CTTTCTGGA TTTTTTACTC CTATTGGGT TTTATAGTAA 10
20

AAATACATAA TAAAAGGAAA ACAAAGTTT TATAGATTCT CTAAACCCC TTACGATAAA 10
80

AGTTGGAATC AAAATAATTC AGGATCAGAT GCTCTTTGAT TGATTAGAT GCGATTACAG 11
40

TTGCATGGAA AATTTTCTAG ATCCGTCGTC ACATTTTATT TTCTGTTAA ATATCTAAAT 12
00

CTGATATATG ATGTCGACAA ATTCTGGTGG CTTATACATC ACTTCAACTG TTTTCTTTTG 12
60

GCTTTGTTTG TCAACTGGT TTTCAATACG ATTTGTGATT TCATCGCTG AATTTTTAAT 13
20

```

ACAAGCAAAC TGATGTTAAC CACAAGCAAG AGATGTGACC TGCCTTATTA ACATCGTATT 13  
 80  
 ACTTACTACT AGTCGTATTC TCAACGCAAT CGTTTTTGTA TTTCTCACAT TATGCCGCTT 14  
 40  
 CTCTACTCTT TATTCCTTTT GGTCCACGCA TTTTCTATTT GTGGCAATCC CTTTCACAAC 15  
 00  
 CTGATTTCCC ACTTTGGATC ATTTGTCTGA AGACTCTCTT GAATCGTTAC CACTTGTTTC 15  
 60  
 TTGTGCATGC TCTGTTTTTT AGAATTAATG ATAAACTAT TCCATAGTCT TGAGTTTTCA 16  
 20  
 GCTTGTGAT TCTTTTGCTT TTGGTTTTCT GCAGAAACAT GGGTGCAGGT GGAAGAATGC 16  
 80  
 CGGTTCTTAC TTCTTCCAAG AAATCGGAAA CCGACACCAC AAAGCGTGTG CCGTGCAGAGA 17  
 40  
 AACCGCCTTT CTCGGTGGGA GATCTGAAGA AAGCAATCCC GCCGCATTGT TTCAAACGCT 18  
 00  
 CAATCCCTCG CTCTTTCTCC TACCTTATCA GTGACATCAT TATAGCCTCA TGCTTCTACT 18  
 60  
 ACGTCGCCAC CAATTACTTC TCTCTCCTCC CTCAGCCTCT CTCTTACTTG GCTTGGCCAC 19  
 20  
 TCTATTGGGC CTGTCAAGGC TGTGTCCTAA CTGGTATCTG GGTCATAGCC CACGAATGCG 19  
 80  
 GTCACCACGC ATTCAGCGAC TACCAATGGC TGGATGACAC AGTTGGTCTT ATCTTCCATT 20  
 40  
 CCTTCCTCCT CGTCCCTTAC TTCTCCTGGA AGTATAGTCA TCGCCGTCAC CATTCCAACA 21  
 00  
 CTGGATCCCT CGAAAGAGAT GAAGTATTG TCCCAAAGCA GAAATCAGCA ATCAAGTGGT 21  
 60  
 ACGGGAAATA CCTCAACAAC CCTCTTGGAC GCATCATGAT GTTAACCGTC CAGTTTGTC 22  
 20  
 TCGGGTGGCC CTTGTACTTA GCCTTAAACG TCTCTGGCAG ACCGTATGAC GGGTTTCGCTT 22  
 80  
 GCCATTTCTT CCCCACGCT CCCATCTACA ATGACCGAGA ACGCCTCCAG ATATACCTCT 23  
 40  
 CTGATGCGGG TATTCTAGCC GTCTGTTTTG GTCTTTACCG TTACGCTGCT GCACAAGGGA 24  
 00  
 TGGCCTCGAT GATCTGCCTC TACGGAGTAC CGCTTCTGAT AGTGAATGCG TTCCTCGTCT 24  
 60

009201"62326600

TGATCACTTA CTTGCAGCAC ACTCATCCCT CGTTGCCTCA CTACGATTCA TCAGAGTGGG 25  
 20  
 ATGGGCTCAG GGGAGCTTTG GCTACCGTAG ACAGAGACTA CGGAATCTTG AACAAAGGTGT 25  
 80  
 TCCACAACAT TACAGACACA CACGTGGCTC ATCACCTGTT CTCGACAATG CCGCATTATA 26  
 40  
 ACGCAATGGA AGTACAAAG GCGATAAAGC CAATTCTGGG AGACTATTAC CAGTTCGATG 27  
 00  
 GAACACCGTG GTATGTGGCG ATGTATAGGG AGGCAAAGGA GTGTATCTAT GTAGAACCGG 27  
 60  
 ACAGGGAAGG TGACAAGAAA GGTGTGTACT GGTACAACAA TAAGTTATGA GGATGATGGT 28  
 20  
 GAAGAAATTG TCGACTTTTC TCTTGTCTGT TTGTCTTTTG TTAAAGAAGC TATGCTTCGT 28  
 80  
 TTTAATAATC TTATTGTCCA TTTTGTGTGT TTATGACATT TTGGCTGCTC ATTATGTTAT 29  
 40  
 GTGGGAAGTT AGCGTTCAAA TGTTTTGGGT CGG 29  
 73

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 23 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (ix) FEATURE:
- (A) NAME/KEY: misc\_feature
  - (B) LOCATION: 1..23
  - (D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGGCATGTNG ARAANARRTG RTG 23

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:



(A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: misc\_feature  
(B) LOCATION: 1..23  
(D) OTHER INFORMATION: /product=  
"synthetic  
oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGGCATGTRC TRAANARRTG RTG

23

EXPRESS MAIL LABEL NO. EL073740691US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN THE APPLICATION OF:

E. I. DUPONT DE NEMOURS AND COMPANY

CASE NO.: BB1043 US NA DIV

APPLICATION NO.: UNKNOWN

GROUP ART UNIT: UNKNOWN

FILED: CONCURRENTLY HERewith

EXAMINER: UNKNOWN

FOR: GENES FOR MICROSOMAL DELTA-12  
FATTY ACID DESATURASES AND  
RELATED ENZYMES FROM PLANTS


Assistant Commissioner for Patents  
Washington, DC 20231

Sir:

**DECLARATION IN ACCORDANCE WITH 37 CFR 1.821**

I hereby state that the content of the paper and computer readable copies of the Sequence Listing, submitted in accordance with 37 CFR 1.821(c) and (e), respectively are the same.

Respectfully submitted,



LYNNE M. CHRISTENBURY  
ATTORNEY FOR APPLICANTS  
REGISTRATION NO. 30,971  
TELEPHONE: 302-992-5481  
FACSIMILE: 302-892-1026

Dated: October 26, 2000

09697379-102600

DECLARATION and  
POWER OF ATTORNEY

As a below-named inventor, I declare that I believe that I am the original, first and sole inventor (if only one name is listed below) or an original first and joint inventor (if plural names are listed below) of the subject matter which is claimed in the specification identified below and in which a patent is sought on the invention as titled therein. I hereby state that I have reviewed and understand the contents of said specification including the claims, as amended by any amendment referred to herein. I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 CFR §1.56.

Full Name of Inventor <sup>(1)</sup>	Last Name	First Name	Middle Name
	<b>LIGHTNER</b>	<b>JONATHAN</b>	<b>EDWARD</b>
Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
	<b>MARIETTA</b>	<b>PENNSYLVANIA</b>	<b>U.S.A.</b>
Post Office Address	Post Office Address	City	State or Country
	<b>R.D.2 BOX 101A</b>	<b>AIRVILLE</b>	<b>PENNSYLVANIA</b>
			Zip Code
			<b>17302</b>
Full Name of Inventor <sup>(2)</sup>	Last Name	First Name	Middle Name
	<b>OKULEY</b>	<b>JOHN</b>	<b>JOSEPH</b>
Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
	<b>COLUMBUS</b>	<b>OHIO</b>	<b>U.S.A.</b>
Post Office Address	Post Office Address	City	State or Country
	<b>217 FALLIS ROAD</b>	<b>COLUMBUS</b>	<b>OHIO</b>
			Zip Code
			<b>43214</b>
Full Name of Inventor <sup>(3)</sup>	Last Name	First Name	Middle Name
Residence & Citizenship	City	State or Foreign Country	Country of Citizenship
Post Office Address	Post Office Address	City	State or Country
			Zip Code



Title of the Invention: **Genes for Microsomal Delta-12 Fatty Acid Desaturases and Related Enzymes from Plants**

- ☒ Patent specification attached.
- ☐ Patent specification was filed on \_\_\_\_\_ as U.S. application Serial No. \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).
- ☐ No earlier-filed foreign applications of which priority benefit is claimed.
- ☐ Required information as to earlier-filed foreign applications of which priority benefit is claimed appears on page 2 attached.
- ☐ No earlier-filed U.S. applications of which priority benefit is claimed.
- ☒ Required information as to earlier-filed U.S. applications of which priority benefit is claimed appears on page 2 attached.

**POWER OF ATTORNEY:** The power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith is hereby granted to the following ☒ attorney(s) ☐ agent(s):

Name	Registration No.:
<b>Barbara C. Siegell</b>	<b>30,684</b>
<b>Hoge T. Sutherland</b>	<b>20,010</b>
Send correspondence and direct telephone calls to:	E. I. du Pont de Nemours & Co. Legal - Patents Wilmington, DE 19898
<b>Barbara C. Siegell</b>	Tel. No. <b>(302) 992-4931</b>

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor <sup>(1)</sup>	Signature of Inventor <sup>(4)</sup>	Signature of Inventor <sup>(3)</sup>
		
Date <b>20 JUNE 1994</b>	Date <b>Sept 24, 1994</b>	Date

09697375-102600

☐ Required information as to earlier-filed foreign applications of which priority benefit is claimed.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

Country	Application No.	Filing Date	Priority Claimed (Yes-No)
---------	-----------------	-------------	---------------------------

☒ Required information as to earlier-filed U.S. applications of which priority benefit is claimed.

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112. I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in 37 CFR §1.56, which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Application Serial No.	U.S. Filing Date	Status (patented, pending or abandoned)
07/977,339	11/17/92	Abandoned
PCT/US93/09987	10/15/93	Pending
U.S. National Application filed 6/17/94		